Revised Draft

Site Characterization and Focused Feasibility Study for the RVAAP-51 Dump Along Paris-Windham Road

Ravenna Army Ammunition Plant Ravenna, Ohio

Contract No. W912QR-08-D-0008 Delivery Order No. 0014

Prepared for:



US Army Corps of Engineers_®

United States Army Corps of Engineers Louisville District 600 Martin Luther King, Jr. Place Louisville, Kentucky 40202

Prepared by:



Science Applications International Corporation 151 Lafayette Drive Oak Ridge, Tennessee 37831

April 5, 2012

DISCLAIMER STATEMENT

This report is a work prepared for the United States Government by Science Applications International Corporation. In no event shall either the United States Government or Science Applications International Corporation have any responsibility of liability for any consequences of any, use, misuse, inability to use, or reliance on the information contained herein, nor does either warrant or otherwise represent in any way the accuracy, adequacy, efficacy, or applicability of the contents hereof.

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any benafty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

			f information if it does not displa HE ABOVE ADDRESS.	ay a currently valid (OMB control	number.		
1. REPORT DAT			ORT TYPE			3. DATES COVERED (From - To)		
	04-2012	·	Technica	ા		April 2012		
4. TITLE AND S	UBTITLE	_		-	5a. CO	NTRACT NUMBER		
Revised Draft					Cartes et Na. W0120D0 00 D 0000			
Site Characteriz	zation and Foo	used Feasibili	ty Study			Contract No. W912QR0-08-D-0008		
for the RVAAF					5b. GR	ANT NUMBER		
Ravenna Army	Ammunition	Plant				NA		
Ravenna, Ohio					5c. PRO	OGRAM ELEMENT NUMBER		
,						NA		
6. AUTHOR(S)					E4 DB0	DJECT NUMBER		
Jill Kovalchik					Ju. 1110	Delivery Order 0014		
Barney Cornab	v. PhD					•		
Sharon Robers	<i>J</i> , 1123				be. TAS	SK NUMBER		
						NA		
					5f. WO	RK UNIT NUMBER		
						NA		
7 DEDECORATE	0.000.48117.471	O B L B L A B # E / O \ A	ND ADDDEGG(FG)			1 8. PERFORMING ORGANIZATION		
			ND ADDRESS(ES)			REPORT NUMBER		
SAIC Engineer 8866 Commons	nc.				3827.20120405.001			
Twinsburg, OH						3027.20120403.001		
I willsburg, Off	1 44007							
9. SPONSORIN	G/MONITORING	AGENCY NAN	IE(S) AND ADDRESS(ES)	10. SPONSOR/MONITOR'S ACRON			
USACE - Louis	sville District				CELRL-ED-EE			
U.S. Army Cor		s, Louisville I	District					
600 Martin Lut						11. SPONSOR/MONITOR'S REPORT		
P.O. Box 50						NUMBER(S)		
Louisville, Ken	tucky 40202-0	0059			USACE			
12. DISTRIBUTION	ON/AVAILABIL	TY STATEMEN	Т					
Reference Distr	ribution Page							
13. SUPPLEMEN	ITADV NOTES							
	IIANT NOTES							
None								
14. ABSTRACT								
This Site Chara	cterization/Fo	cused Feasibil	ity Study presents phys	sical character	istics and	I nature and extent of contamination;		
						ssments; identifies response actions,		
						address chemicals of concern (COCs) for		
						ive to meet the remedial action objective at		
this AOC. The	recommended	alternative for	this AOC is "Alternati	ive 2: Land U	se Contro	ols."		
15. SUBJECT TI								
Investigation, c	haracterizatio	n, feasibility st	udy, remediation					
16. SECURITY O	N A COLEIO A TIO	N OE:	17. LIMITATION OF	18 MIIMDED	100 814	ME OF RESPONSIBLE PERSON		
	b. ABSTRACT	c. THIS PAGE	ABSTRACT	OF	I 198. NA	Joan Cullen		
	S. ADOTHACT	o. IIIIo FAGE	G + D	PAGES	19h TE	EPHONE NUMBER (Include area code)		
U	U	U	SAR	192	55. 161	502.315.6344		

PLACE HOLDER FOR:

Documentation of Ohio EPA Approval of Final Document

(Documentation to be provided once approval is issued.)

Revised Draft

Site Characterization and Focused Feasibility Study for the RVAAP-51 Dump Along Paris-Windham Road

Ravenna Army Ammunition Plant Ravenna, Ohio 44266

Contract No. W912QR-08-D-0008 Delivery Order No. 0014

Prepared For:

U.S. Army Corps of Engineers 600 Martin Luther King, Jr. Place Louisville, Kentucky 40202

Prepared By:

Science Applications International Corporation 151 Lafayette Drive Oak Ridge, Tennessee 37831

April 5, 2012

CONTRACTOR STATEMENT OF INDEPENDENT TECHNICAL REVIEW

Science Applications International Corporation (SAIC) has completed the Site Characterization and Focused Feasibility Study for the RVAAP-51 Dump Along Paris-Windham Road at the Ravenna Army Ammunition Plant, Ravenna, Ohio. Notice is hereby given that an independent technical review has been conducted that is appropriate to the level of risk and complexity inherent in the project. During the independent technical review, compliance with established policy principles and procedures, utilizing justified and valid assumptions, was verified. This included review of data quality objectives; technical assumptions; methods, procedures, and materials to be used; the appropriateness of data used and level of data obtained; and reasonableness of the results, including whether the product meets the customer's needs consistent with law and existing United States Army Corps of Engineers (USACE) policy.

Jee Kovalchite	1/7/11
Jill Kovalchik	Date
Study/Design Team Leader	
5125/h	1/7/11
Sharon Stoller	Date
Independent Technical Review Team Leader	
Harry Fatkin	1/7/11 Date
•	Date
Human Health Risk Assessment Reviewer	
Michael S. Buta	1/7/11
Mike Barta	Date
Ecological Risk Assessment Reviewer	
Dal Thomas	1/7/11
Jed Thomas	
Feasibility Study Reviewer	

Significant concerns and the explanation of the resolution are as follows:

Internal SAIC Independent Technical Review was conducted on the Preliminary Draft version of this document. Subsequent versions of this document (e.g., Draft and Final) incorporated changes based on the technical reviews of USACE, the Ohio Army National Guard, and the Ohio Environmental Protection Agency. Internal SAIC Independent Technical Review comments are recorded on a Document Review Record per SAIC quality assurance procedure QAAP 3.1. This Document Review Record is maintained in the project file. Changes to the report addressing the comments have been verified by the Study/Design Team Leader. As noted above, all concerns resulting from independent technical review of the project have been considered.

W. Lewin Jago	1/10/11
Kevin Jago	Date
Principal w/ A-E firm	

DOCUMENT DISTRIBUTION

for the

Revised Draft

Site Characterization and Focused Feasibility Study for the RVAAP-51 Dump Along Paris-Windham Road Ravenna Army Ammunition Plant Ravenna, Ohio

	Number of	Number of
Name/Organization	Printed Copies	Electronic Copies
Kimberly Harriz, NGB	0	1
Katie Tait, OHARNG	1	1
Brian Tucker, Ohio EPA – Central Office	1	1
Todd Fisher, Ohio EPA – NEDO	1	1
Eileen Mohr, Ohio EPA – NEDO	1	1
Mark Patterson, RVAAP Facility Manager	2	2
Glen Beckham, USACE – Louisville District	1	1
Joan Cullen, USACE – Louisville District	1	1
Angela Schmidt, USACE – Louisville District	1	1
Mark Eldridge, USAEC	0	1
Kevin Jago, SAIC	1	1
Jed Thomas, SAIC	1	1
REIMS	0	1
SAIC Project File W912QR-05-F-0033	1	1
SAIC Central Records Facility	0	1

NEDO = Northeast District Office

NGB = National Guard Bureau

OHARNG = Ohio Army National Guard

Ohio EPA = Ohio Environmental Protection Agency

REIMS = Ravenna Environmental Information Management System

RVAAP = Ravenna Army Ammunition Plant

SAIC = Science Applications International Corporation

USACE = United States Army Corps of Engineers

USAEC = United States Army Environmental Command

TABLE OF CONTENTS

2			
3	LIST	OF TABLES	iv
4	LIST	OF FIGURES	iv
5	LIST	OF PHOTOGRAPHS	v
6	LIST	OF APPENDICES	
7		DNYMS AND ABBREVIATIONS	
8	1.0	INTRODUCTION	1- 1
9	1.1	PURPOSE	1-1
10	1.2	SCOPE	
11	1.3	REPORT ORGANIZATION	1-4
12	2.0	BACKGROUND INFORMATION	2- 1
13	2.1	RAVENNA ARMY AMMUNITION PLANT FACILITY DESCRIPTION	
14	2.2	DUMP ALONG PARIS-WINDHAM ROAD DESCRIPTION	
15	3.0	SUMMARY OF HISTORICAL DATA AND OCCURRENCE AND	
16		DISTRIBUTION OF CONTAMINATION	
17	3.1	PREVIOUS INVESTIGATIONS AND ACTIVITIES	
18	3.	1.1 Relative Risk Site Evaluation	
19		1.2 Decision Document for a Removal Action at the Paris-Windham Road Dumpsite	
20	3.	1.3 Limited Remedial Design/Remedial Action	
21		3.1.3.1 Limited Remedial Design/Remedial Action Sampling Results	
22		3.1.3.2 Limited Remedial Design/Remedial Action Conclusions	
23	3.2	NATURE AND EXTENT OF CONTAMINATION	
24		2.1 Site-Related Contaminants	
25	3.	2.2 Occurrence and Distribution of Contaminants	
26 27		3.2.2.1 Soil	
2 <i>1</i> 28	3	2.3 Soil to Groundwater Leaching Screen	
20 29		2.4 Conceptual Site Model	
30	٥.	3.2.4.1 Primary and Secondary Sources	
31		3.2.4.2 Migration Pathways and Receptors	
32		3.2.4.3 Uncertainties and Data Gaps	
_		•	
33	4.0	HUMAN HEALTH RISK ASSESSMENT AND ECOLOGICAL RISK	
34		ASSESSMENT	
35	4.1	CONCEPTUAL SITE EXPOSURE MODEL	4-1
36	4.2	DATA EVALUATION FOR HUMAN HEALTH AND ECOLOGICAL RISK	
37		ASSESSMENTS	
38		2.1 Data Aggregate – Soil	
39		2.2 Data Aggregate – Surface Water	
40	4.3	HUMAN HEALTH RISK ASSESSMENT	
41		3.1 Identify Media of Concern	
42 42	4.	3.2 Identify Chemicals of Potential Concern	
43		4.3.2.1 Chemicals of Potential Concern in the Fill Area Exposure Unit for Soil	4-10

1		TABLE OF CONTENTS (CONTINUED)	
2			
3		4.3.2.2 Chemicals of Potential Concern in the Surface Area Exposure Unit for Soil	
4		4.3.2.3 Chemicals of Potential Concern in Incremental Sampling Method Soil Samples	
5		4.3.2.4 Chemicals of Potential Concern in Surface Water	
6	4.3		
7	4.3		4-12
8		4.3.4.1 Selection of Appropriate Facility-Wide Cleanup Goals for the Dump Along	
9		Paris-Windham Road	
10		4.3.4.2 Exposure Point Concentrations for Comparison to Facility-Wide Cleanup Goals	4-14
11		4.3.4.3 Identification of Dump Along Paris-Windham Road Chemicals of Concern: Range	
12		Maintenance Soldier Scenario	4-14
13		4.3.4.4 Identification of Dump Along Paris-Windham Road Chemicals of Concern: Trespasser	4.1
14		Scenario	4-1:
15		4.3.4.5 Identification of Dump Along Paris-Windham Road Chemicals of Concern:	4.1.
16	4.2	Resident Farmer Scenario	
17 18	4.3	,	
19		4.3.5.1 Uncertainty in Estimating Potential Exposure	
20		4.3.5.2 Uncertainty in Use of Facility-Wide Cleanup Goals	
21	4.3	•	
22	4.3		
23	4.4		
24	4.4		
25	4.4	· · · · · ·	
26		4.4.3.1 AOC Description and Land Use	
27		4.4.3.2 Evidence of Chemical Contamination	
28		4.4.3.3 Ecological Significance	
29		4.4.3.4 Evaluation of Chemical Contamination and Ecological Significance	
30		4.4.3.5 Summary and Recommendations of Scoping Level Ecological Risk Assessment	
31	4.4	•	
32	5.0	REMEDIAL ACTION OBJECTIVE	5-
33	5.1	REMEDIAL ACTION OBJECTIVE	
34	5.2	REASONABLE AND ANTICIPATED FUTURE LAND USE	5-
35	5.3	FACILITY-WIDE CLEANUP GOALS	5-2
36	6.0	APPLICABLE OR RELEVANT AND APPROPRIATE REQUIREMENTS	
37	6.1	INTRODUCTION	
38	6.2	POTENTIAL APPLICABLE OR RELEVANT AND APPROPRIATE REQUIREMENTS	6
39	6.3	POTENTIAL CHEMICAL-SPECIFIC APPLICABLE OR RELEVANT AND	
40		APPROPRIATE REQUIREMENTS	6-4
41	6.4	POTENTIAL ACTION-SPECIFIC APPLICABLE OR RELEVANT AND	_
42	- -	APPROPRIATE REQUIREMENTS	6-4
43	6.5	POTENTIAL LOCATION-SPECIFIC APPLICABLE OR RELEVANT AND	_
44		APPROPRIATE REQUIREMENTS	6-:
45			

TABLE OF CONTENTS (CONTINUED)

7-1
7-1
8-1
8-1
8-1
9-1
9-1
9-2
9-2
9-3
9-4
9-4
9-4
9-4
9-5
9-5
9-5
9-5
9-5
9-5
9-6
9-6
9-6
9-6
9-7
9-7
9-7
9-7
10-1
10-1
10-1
11-1
11-1
11-1
12-1

39

1		LIST OF TABLES	
2	Table 3-1.	Results of Limited "RD/RA" Confirmatory Surface Soil Discrete Samples –	
3		Inorganic Chemicals	3-5
4	Table 3-2.	Results of Limited "RD/RA" Confirmatory Surface Soil and Dry Sediment Discret	te
5		Samples – Organic Chemicals	3-6
6	Table 3-3.	Results of Limited "RD/RA" Confirmatory Surface Water Discrete Samples	3-7
7	Table 3-4.	Results of Limited "RD/RA" Confirmatory Dry Sediment Discrete Samples –	
8		Inorganic Chemicals	3-8
9	Table 3-5.	Results of Limited "RD/RA" Contingency Incremental Sampling Method	
10		Surface Soil Samples	3-9
11	Table 3-6.	Soil SRCs	3-11
12	Table 3-7.	Surface Water SRCs	3-12
13	Table 3-8.	Results of Contaminant Migration Soil to Groundwater Screening	3-14
14	Table 4-1.	Risk Assessment Datasets for Soil	4-5
15	Table 4-2.	Risk Assessment Dataset for Surface Water	4-6
16	Table 4-3.	Summary of COPCs	4-10
17	Table 4-4.	FWCUGs for COPCs in Soil	4-13
18	Table 4-5.	FWCUGs for COPCs in Surface Water	4-14
19	Table 4-6.	Summary of COCs and FWCUGs	4-20
20	Table 4-7.	Summary of COPECs for Surface Soil at the Fill Area EU	4-26
21	Table 4-8.	Summary of COPECs for Surface Soil at the Surface Area EU	4-27
22	Table 4-9.	Summary of COPECs for Surface Water	4-28
23	Table 4-10.	Summary of COPEC Concentrations for Dry Sediment and Surface Water at and	
24		in the Vicinity of the Dump Along Paris-Windham Road	4-36
25	Table 4-11.	Comparison of Five Assessment Techniques at Sampling Stations Near the	
26		Dump Along Paris-Windham Road	4-39
27	Table 4-12.	Survey of Proximity to the AOC of Various Ecological Resources	4-40
28	Table 5-1.	Chemicals of Concern and Cleanup Goals by Media and Receptor	5-2
29	Table 6-1.	Potential Action ARARs for Disposal of RCRA Hazardous Waste	6-5
30	Table 9-1.	Comparison of Alternatives by Evaluation Criteria	9-7
31			
32		LIST OF FIGURES	
33	Figure 2-1.	General Location of RVAAP	2-3
34	•	Location of Dump Along Paris-Windham Road Within RVAAP	
35	•	Dump Along Paris-Windham Road	
36	-	Limited "RD/RA" Location Map	
37	•	Conceptual Site Exposure Model	
38	_	Risk Assessment Input to Support Remediation Decisions	
39	-	Exposure Units at the Dump Along Paris-Windham Road	
40	_	Natural Resources (OHARNG 2008) Inside and Near Habitat Area at the	
41	-	Dump Along Paris-Windham Road	4-31

1		LIST OF PHOTOGRAPHS
2		
3	Photograph 4-1.	Looking South along Paris-Windham Road; Green Ash, American Elm, and
4		Hackberry Temporary Flooded Forest Alliance in Background4-32
5	Photograph 4-2.	Drainage Swale with no Standing Water Along the Northwest Portion of the
6		AOC; Green Ash, American Elm, and Hackberry Temporary Flooded Forest
7		Alliance in Background4-32
8		
9		LIST OF APPENDICES
10		
11	Appendix A.	Photographs
12	Appendix B.	Human Health Risk Assessment Supporting Data
13	Appendix C.	Ecological Risk Assessment Information and Data
14	Appendix D.	Detailed Cost Estimates

ACRONYMS AND ABBREVIATIONS

ACM Asbestos-Containing Material

amsl Above Mean Sea Level

AOC Area of Concern

ARAR Applicable or Relevant and Appropriate Requirement

bgs Below Ground Surface

BRAC Base Realignment and Closure C&D Construction and Demolition

CERCLA Comprehensive Environmental Response, Compensation, and Liability Act

CFR Code of Federal Regulations

CMCOPC Contaminant Migration Chemical of Potential Concern

COC Chemical of Concern

COPC Chemical of Potential Concern

COPEC Chemical of Potential Ecological Concern

CSM Conceptual Site Model

CUG Cleanup Goal

DD Decision Document

DERR Division of Emergency and Remedial Response

DFFO Director's Final Findings and Orders

ERA Ecological Risk Assessment
EPC Exposure Point Concentration
ESA Endangered Species Act
ESL Ecological Screening Level
ESV Ecological Screening Value

EU Exposure Unit

FFS Focused Feasibility Study

FS Feasibility Study

FWCUG Facility-Wide Cleanup Goal

FWERWP Facility-Wide Ecological Risk Work Plan

FWHHRAM Facility-Wide Human Health Risk Assessor Manual

GRA General Response Action

HHRA Human Health Risk Assessment

HO Hazard Quotient

IBI Index of Biotic Integrity

ICI Invertebrate Community Index

INRMP Integrated Natural Resources Management Plan

IRP Installation Restoration Program ISM Incremental Sampling Method

LUC Land Use Control

MCL Maximum Contaminant Level MDC Maximum Detected Concentration

MDL Method Detection Limit
MFL Million Fibers per Liter
MKM MKM Engineers, Inc.

ACRONYMS AND ABBREVIATIONS (CONTINUED)

NCP National Oil and Hazardous Substances Pollution Contingency Plan

NFA No Further Action

O&M Operation and Maintenance OAC Ohio Administrative Code OHARNG Ohio Army National Guard

Ohio EPA Ohio Environmental Protection Agency

ORAM Ohio Rapid Assessment Method
PAH Polycyclic Aromatic Hydrocarbon
PBT Persistent, Bioaccumulative, and Toxic

PCB Polychlorinated Biphenyl
PMP Property Management Plan

PP Proposed Plan

PRG Preliminary Remediation Goal

QA Quality Assurance
QC Quality Control

QHEI Qualitative Habitat Evaluation Index

RA Remedial Action

RAFLU Reasonable and Anticipated Future Land Use

RAO Remedial Action Objective

RCRA Resource Conservation and Recovery Act

RD Remedial Design
RI Remedial Investigation

RM River Mile

ROD Record of Decision

RRSE Relative Risk Site Evaluation
RSL Regional Screening Level

RVAAP Ravenna Army Ammunition Plant

SAIC Science Applications International Corporation

SC Site Characterization SDZ Safety Danger Zone SOR Sum-of-Ratios

SRC Site-related Contaminant SRV Sediment Reference Value SSL Soil Screening Level

SVOC Semi-volatile Organic Compound T&E Threatened and Endangered

TAL Target Analyte List

TEC Threshold Effects Concentration

TNT 2,4,6-Trinitrotoluene TOC Total Organic Carbon

TR Target Risk

TSCA Toxic Substances Control Act

UCL₉₅ 95% Upper Confidence Limit of the Mean

USACHPPM United States Army Center for Health Promotion and Preventive Medicine

ACRONYMS AND ABBREVIATIONS (CONTINUED)

USACE United States Army Corps of Engineers

USEPA Unites States Environmental Protection Agency

USFS United States Forest Service VOC Volatile Organic Compound

WOE Weight-of-Evidence

WP Work Plan

WQS Water Quality Standard WWH Warm Water Habitat

1.0 Introduction

- 1 Science Applications International Corporation (SAIC) has been contracted by the United States
- 2 Army Corps of Engineers (USACE), Louisville District to perform a Site Characterization
- 3 (SC)/Focused Feasibility Study (FFS) for the Dump Along Paris-Windham Road at the Ravenna
- 4 Army Ammunition Plant (RVAAP). The Dump Along Paris-Windham Road is designated as Area of
- 5 Concern (AOC) RVAAP-51 in the RVAAP Installation Restoration Program (IRP).

6

- 7 This work is being performed under Contract Number W912QR-08-D-0008, Delivery Order Number
- 8 0014, issued by USACE, Louisville District on June 16, 2009, and in compliance with the Site
- 9 Characterization and Focused Feasibility Study Work Plan for the RVAAP-51 Dump Along Paris-
- Windham Road (USACE 2010c) [herein referred to as the SC/FFS Work Plan (WP)]. In addition,
- planning and performance of all elements of this work are in accordance with the requirements of the
- Ohio Environmental Protection Agency (Ohio EPA) Director's Final Findings and Orders (DFFO)
- 13 dated June 10, 2004 (Ohio EPA 2004).

14 15

1.1 PURPOSE

16 17

18

19

20

21

22

23

24

25

26

27

28

This SC/FFS characterizes the AOC [following a limited remedial design/remedial action (RD/RA)] and identifies the final RA alternatives for soil at the Dump Along Paris-Windham Road. The limited "RD/RA," as titled by the U.S. Army Base Realignment and Closure (BRAC) Division, was performed in 2003. Activities included the removal of surface debris, excavation of transite along the embankment to the extent practicable (without undermining Paris-Windham Road), confirmatory sampling to evaluate the success of the RA, and placement of a protective soil and vegetation cover over portions of the AOC. The limited "RD/RA" terminology has been retained within this SC/FFS to be consistent with historical documents; however, the Ohio EPA commented the "RD/RA" should be considered an interim removal action and not a final remedy (MKM 2004). The limited "RD/RA" did not evaluate the nature and extent of contamination or identify chemicals of potential concern (COPCs) or chemicals of concern (COCs). This SC/FFS completes these tasks and evaluates the remedial alternatives, as required, to address impacts to environmental media in accordance with the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) process.

293031

The SC portion of this document:

- Describes the AOC conditions;
- Summarizes historical data:
- Evaluates the nature and extent of contamination at the AOC;
- Presents a conceptual site model (CSM);
- Evaluates contaminant risk to human and ecological receptors;
- Determines COPCs and COCs; and
- Identifies applicable facility-wide cleanup goals (FWCUGs).

Groundwater data do not exist for the vicinity of this AOC; therefore, only a qualitative evaluation of potential impacts of residual soil contaminants on groundwater quality is included in the SC portion of this document. The U.S. Army will address groundwater at this AOC under a future decision for the RVAAP Facility-Wide Groundwater AOC (RVAAP-66). Surface water at the AOC occurs intermittently in a drainage swale located at the base of the slope face of the dump. As discussed in Sections 1.2 and 4.0, surface water within this AOC is evaluated in the SC; however, no further action (NFA) is recommended based on assessment of human health and ecological risks. Therefore, the FFS portion of this document evaluates remedial alternatives to obtain a final remedy for soil at the Dump Along Paris-Windham Road.

For the purposes of this SC/FFS, the term "surface soil" includes dry sediment. Dry sediment refers to unconsolidated inorganic and organic material within conveyances, ditches, or low-lying areas that occasionally may be covered with water, usually following a precipitation event or due to snowmelt. Dry sediment is not covered with water for extended periods and typically is dry within 7 days (USACE 2008). It does not function as a permanent habitat for aquatic organisms; although, it may serve as a natural medium for the growth of terrestrial organisms. Dry sediment is addressed in the same manner as surface soil [0-1 ft below ground surface (bgs)] in terms of contaminant nature and extent, fate and transport, and risk exposure models. The definitions and terminology usages for dry sediment within this SC/FFS are consistent with the *Facility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition Plant, Ravenna, Ohio* (USACE 2010a) (herein referred to as the FWCUG Report).

An FFS develops a remedial action objective (RAO) to protect receptors from impacted environmental media and COCs and identifies applicable or relevant and appropriate requirements (ARARs). In accordance with the RAO, a focused technology screening is performed and remedial alternatives are developed to provide options to reduce risks to the environment and human health.

The recommended alternative listed in the FFS will be presented in a Proposed Plan (PP) for public review and comment. Public comments will be considered in the final selection of a remedy, which will be documented in a Record of Decision (ROD). Responses to public comments will be addressed in the responsiveness summary of the ROD.

1.2 SCOPE

This SC/FFS uses historical data to characterize the nature and extent of contaminants in soil (inclusive of dry sediment) and intermittent surface water at the Dump Along Paris-Windham Road. Risks to human and ecological receptors posed by contaminants in soil and surface water are assessed, and potential impacts to groundwater from residual soil contaminants are evaluated. The assessments of human health and ecological risks for surface water (Section 4.0) address temporal exposures due to the fact that surface water is present at the AOC on an intermittent basis. No human health COCs were identified for surface water, and the ecological risk assessment (ERA) recommended NFA with respect to ecological receptors. Based on the risk assessment results, the FFS does not include remedial alternatives for surface water. Potential final remedies for soil are

presented along with a detailed analysis and comparative evaluation. This document provides the information and decisions necessary for a subsequent PP and ROD to address soil media at the AOC.

2 3 4

1

The Ohio Army National Guard (OHARNG) has established a Reasonable and Anticipated Future Land Use (RAFLU) for the Dump Along Paris-Windham Road as follows (OHARNG 2008b):

567

• Dump Along Paris-Windham Road (RVAAP-51) Restricted Access OHARNG Military Use and Training Land Use.

8 9 10

• Representative Receptor – National Guard Range Maintenance Soldier

11 12

13

14

1516

17

18

19

20

21

22

23

2425

26

27

RVAAP is a controlled-access facility that is fenced and patrolled by security personnel. Full-time OHARNG, BRAC, and contractor staff work at the facility. Military training and operations are conducted at the facility. The AOC is located in the eastern central portion of the facility. The AOC is not currently used for military training activities but may receive periodic foot traffic. The OHARNG projected future land use for the AOC is Restricted Access due to residual asbestos at the AOC. The most representative receptor is the National Guard Range Maintenance Soldier. This anticipated future land use, in conjunction with the evaluation of agricultural-residential land uses and associated receptors, form the basis for identifying COCs in this RI. Residential land use, specifically the Resident (adult and child) Farmer scenario, is included in the human health risk assessment (HHRA) (Section 4.3) to evaluate COCs for unrestricted land use at the AOC as required by the CERCLA process and as outlined in the Facility-Wide Human Health Risk Assessor Manual, Amendment 1 (USACE 2005b) (herein referred to as the FWHHRAM); however, the topography of the area (i.e., steep slope and floodplain), precludes Residential Land Use. As described in the approved SC/FFS WP (USACE 2010c) and further discussed in the FFS (Section 5.0), a remedial alternative based on Residential Land Use is not evaluated due to these location and physical characteristics of the AOC, as well as waste remaining in place [construction and demolition (C&D) material, including transite, glass, concrete, brick, metal, and wood debris].

282930

31

32

33

Because the AOC is located immediately adjacent to a primary road, trespassers may potentially visit the AOC; therefore, Adult and Juvenile Trespassers are also considered in the HHRA. The exposure assumptions for the Range Maintenance Soldier are also protective of the Adult and Child trespasser. Per guidelines in the FWCUG Report (USACE 2010a), the application of these receptor scenarios to the Dump Along Paris-Windham Road is described in more detail in Section 4.3.

343536

The following key points relate to the scope of this SC/FFS:

3738

39

• This SC/FFS includes an evaluation of contaminant nature and extent and incorporates existing data. Based on the results of the SC, any remaining data gaps are identified and additional sampling, if required, is recommended.

40 41 42

43

• Data previously collected for the limited "RD/RA" were of good quality; however, the data screening processes employed in the limited "RD/RA" (MKM 2004) were not in conformance

with current RVAAP protocols. This SC/FFS includes an HHRA (Section 4.3), which follows the processes outlined in the FWCUG Report (USACE 2010a), identifies COPCs, COCs, and applicable FWCUGs. This SC/FFS also includes an ERA (Section 4.4), which follows a unified approach of methods integrating U.S. Army, Ohio EPA, and United States Environmental Protection Agency (USEPA) guidance.

6 7

8

9

• Residual fragments of transite covered in place during the limited "RD/RA" are qualitatively evaluated in this SC/FFS with respect to the potential for human exposure (e.g. friable or non-friable asbestos and any mitigating effect of the soil/vegetation cover placed over the dump following the limited "RD/RA").

101112

1.3 REPORT ORGANIZATION

13 14

15

16

This report is organized in accordance with USEPA guidance for CERCLA remedial investigations (RIs) and feasibility studies (FSs) and the proposed outline included in Section 7.0 of the approved SC/FFS WP (USACE 2010c). This report combines an SC and FFS for the Dump Along Paris-Windham Road and is organized as follows:

17 18

• Section 1.0 provides an introduction.

202122

Section 2.0 presents facility and AOC background information.

2324

• Section 3.0 summarizes historical data and evaluates the occurrence and distribution of contaminants at the Dump Along Paris-Windham Road.

25

• Section 4.0 presents the HHRA and ERA.

2728

• Section 5.0 identifies the RAO.

29

• Section 6.0 discusses ARARs.

31

• Section 7.0 presents the limited technology types and process options for RAs.

3334

• Section 8.0 discusses the development of remedial alternatives for the AOC.

35

• Section 9.0 presents detailed and comparative analyses of the remedial alternatives.

37

• Section 10.0 summarizes the partnering and public involvement activities.

39

• Section 11.0 presents the conclusions.

41

• Section 12.0 provides the references used in this report.

- 1 Appendices A through D provide information supporting the evaluations presented within this
- 2 SC/FFS and are organized as follows:

3

- Appendix A contains photographs.
 - Appendix B presents HHRA supporting data.
- 6 Appendix C presents ERA supporting data.
- 7 Appendix D includes the cost evaluations.

2.1 RAVENNA ARMY AMMUNITION PLANT FACILITY DESCRIPTION

When the RVAAP IRP began in 1989, RVAAP was identified as a 21,419-acre facility. The property boundary was resurveyed by OHARNG over a 2-year period (2002 and 2003), and the total acreage of the property was found to be 21,683 acres.

As of June 2010, a total of 20,423 acres of the former 21,683-acre RVAAP has been transferred to the National Guard Bureau and subsequently licensed to OHARNG for use as a military training site (Camp Ravenna Joint Military Training Center). These transferred portions are now referred to as Camp Ravenna. The current RVAAP consists of 1,260 acres in various parcels throughout Camp Ravenna.

Camp Ravenna is in northeastern Ohio within Portage and Trumbull counties, approximately 3 miles east-northeast of the city of Ravenna and approximately 1 mile northwest of the city of Newton Falls (Figure 2-1). The RVAAP portions of the property are solely located within Portage County. Camp Ravenna/RVAAP is a parcel of property approximately 11 miles long and 3.5 miles wide, bounded by State Route 5, the Michael J. Kirwan Reservoir, and the CSX System Railroad on the south; Garret, McCormick, and Berry roads on the west; the Norfolk Southern Railroad on the north; and State Route 534 on the east. Camp Ravenna is surrounded by several communities: Windham on the north; Garrettsville 6 miles to the northwest; Newton Falls 1 mile to the southeast; Charlestown to the southwest; and Wayland 3 miles to the south.

The entire 21,683-acre parcel was an industrial facility that was government-owned, contractor-operated when RVAAP was operational (Camp Ravenna did not exist at that time). The RVAAP IRP encompasses investigation and cleanup of past activities over the entire 21,683 acres of the former RVAAP. References to RVAAP in this document are considered to be inclusive of the historical extent of RVAAP, which is inclusive of the combined acreages of the current Camp Ravenna and RVAAP, unless otherwise specifically stated.

Industrial operations at the former RVAAP consisted of 12 munitions-assembly facilities referred to as "load lines." Load Lines 1 through 4 were used to melt and load 2,4,6-trinitrotoluene (TNT) and Composition B into large-caliber shells and bombs. The operations on the load lines produced explosive dust, spills, and vapors that collected on the floors and walls of each building. Periodically, the floors and walls were cleaned with water and steam. Following cleaning, the wastewater containing TNT and Composition B was known as "pink water" for its characteristic color. Pink water was collected in concrete holding tanks, filtered, and pumped into unlined ditches for transport to earthen settling ponds. Load Lines 5 through 11 were used to manufacture fuzes, primers, and boosters. Potential contaminants in these load lines include lead compounds, mercury compounds, and explosives. Load Line 12 was used from 1946 to 1949 to produce ammonium nitrate for explosives and fertilizers, and portions of the AOC were later used for weapons demilitarization.

In 1950, the facility was placed on standby status, and operations were limited to renovation, demilitarization, normal maintenance of equipment, and munitions storage. Production activities were resumed from July 1954 to October 1957 and again from May 1968 to August 1972. In addition to production missions, various demilitarization activities were conducted at facilities constructed at Load Lines 1, 2, 3, and 12. Demilitarization activities included disassembly of munitions and explosives melt-out and recovery operations using hot water and steam processes. Periodic demilitarization of various munitions continued through 1992.

In addition to production and demilitarization activities at the load lines, other facilities at RVAAP include AOCs that were used for the burning, demolition, and testing of munitions. These burning and demolition grounds consist of large parcels of open space or abandoned quarries. Potential contaminants at these AOCs include explosives, propellants, inorganic chemicals, and waste oils. Other types of AOCs present at RVAAP include landfills, an aircraft fuel tank testing facility, and various general industrial support and maintenance facilities.

2.2 DUMP ALONG PARIS-WINDHAM ROAD DESCRIPTION

The Dump Along Paris-Windham Road is located in the east-central portion of RVAAP, along a steep embankment on the west side of Paris-Windham Road between the bridge over Sand Creek and the intersection of Paris-Windham Road with Remalia Road (Figure 2-2). The AOC was used as an open dump for a variety of miscellaneous C&D material, including asbestos-containing material (ACM) (e.g. transite roofing and siding), laboratory bottles and drums, concrete, brick, glass, scrap metal, fencing, and wood debris. There are no records indicating the quantities of material dumped at the AOC or the dates of operation.

The former dump was approximately 400 ft long by 30 ft wide and slopes east to west, away from Paris-Windham Road (Figure 2-3). The slope face ranges 40 to 60 degrees from horizontal. There are no structures or dwellings on the AOC.

Sand Creek is located to the west and north at distances ranging from approximately 30 ft (north end of the AOC) to 170 ft (south-central portion of the AOC). Surface water runoff follows the topography and flows in a westerly direction through a drainage swale at the base of the dump slope, entering Sand Creek. Surface water within the drainage swale occurs intermittently during and after rainfall events and periods of snow melt. During an August 2009 walkover, SAIC noted the sediment in the drainage swale had high moisture content, but no standing water was observed. During a November 2011 walkover following an extended rainfall event, water was observed. The Sand Creek floodplain occupies the land between the dump and Sand Creek.




Figure 2-1. General Location of RVAAP




Figure 2-2. Location of Dump Along Paris-Windham Road Within RVAAP




Figure 2-3. Dump Along Paris-Windham Road

3.0 SUMMARY OF HISTORICAL DATA AND OCCURRENCE AND DISTRIBUTION OF CONTAMINATION

3.1 Previous Investigations and Activities

2 3

Previous investigative activities at the Dump Along Paris-Windham Road include a Relative Risk Site Evaluation (RRSE) in 1998, environmental sampling conducted by USACE, Louisville District in 2001, and confirmatory/contingency sampling performed during the 2003 limited "RD/RA." The investigations and results are summarized in the following sections.

3.1.1 Relative Risk Site Evaluation

The United States Army Center for Health Promotion and Preventive Medicine (USACHPPM) conducted an RRSE for newly added AOCs at RVAAP in 1998. Thirteen AOCs, including the Dump Along Paris-Windham Road, were evaluated. Three surface soil samples and one sediment sample was collected from the AOC on October 19, 1998 and analyzed for semi-volatile organic compounds (SVOCs), explosives, and inorganic chemicals. No groundwater or surface water samples were collected. The RRSE was summarized in the *Relative Risk Site Evaluation for Newly Added Sites* (USACHPPM 1998).

The RRSE found the AOC contained C&D debris, including ACM (e.g., transite roofing and siding) and inorganic contaminants. The study identified potential human and ecological receptors for surface soil and sediment contamination and assumed complete exposure pathways because there were no access controls (e.g., fence) in place and due to the AOC's proximity to Sand Creek. As a result, the RRSE score for this AOC was "High." Data collected during the 1998 RRSE are not assessed qualitatively or quantitatively in this SC/FFS because these data were minimal Level III data, as defined by the USEPA, and were not intended to be used as definitive evidence of contamination presence or absence or to support quantitative health risk assessment (USACHPPM 1998). Additionally, these data were collected five years prior to the limited "RD/RA" and do not reflect current conditions at the AOC.

3.1.2 Decision Document for a Removal Action at the Paris-Windham Road Dumpsite

In 2003, USACE, Louisville District prepared a *Decision Document for a Removal Action at Paris-Windham Road Dumpsite (RVAAP-51)* (USACE 2003a). The Decision Document (DD) is included in Appendix E of the *Final Report for Remedial Design/Removal Action Plan at Paris-Windham Road Dump* (MKM 2004). As stated in the DD, chemicals in soil include benz(a)anthracene, benzo(a)pyrene, benzo(b)fluoranthene, indeno(1,2,3-cd)pyrene, and inorganic chemicals. The DD reported the principal contaminants with potential impact to human health were benz(a)anthracene, benzo(a)pyrene, benzo(b)fluoranthene, and indeno(1,2,3-cd)pyrene. Chemicals with potential impact to ecological receptors were cadmium, polychlorinated biphenyls (PCBs), benz(a)anthracene, benzo(a)pyrene, benzo(b)fluoranthene, chrysene, and indeno(1,2,3-cd)pyrene.

- 1 The DD outlined four potential remedial alternatives for the AOC: (1) no action; (2) land use controls
- 2 (LUCs); (3) multi-layer cap and LUCs; and (4) removal/disposal of solvent drums, gas cylinders,
- 3 laboratory bottles, and miscellaneous debris with confirmation sampling. Following a public meeting
- 4 and 30-day open comment period, Alternative 4 (Removal/Disposal of Solvent Drums, Gas
- 5 Cylinders, Lab Bottles, and Miscellaneous Debris with Confirmation Sampling) was selected for
 - implementation under a limited "RD/RA."

3.1.3 Limited Remedial Design/Remedial Action

The 2003 limited "RD/RA" activities are summarized in the *Final Report for Remedial Design/Remedial Action Plan at Paris-Windham Road Dump* (MKM 2004). The limited "RD/RA" was conducted in accordance with CERCLA to mitigate risks related to potential contact with exposed waste material. The limited "RD/RA" was not intended to be a final remedy, and (as noted in Section 1.0) the U.S. Army planned for future evaluation of the need for additional characterization and RAs under an SC/FFS and the completion of the CERCLA process (USACE 2010c).

On April 19, 2003, the limited "RD/RA" was initiated at the Dump Along Paris-Windham Road. Initial AOC preparation and mobilization activities included an ordnance and explosive survey. The limited "RD/RA" removal activities consisted of removing all existing surface debris, limited removal of subsurface debris, transportation and disposal of debris, performing confirmation sampling, and AOC restoration. A combined total of 300.66 tons of surface and subsurface debris was removed from the AOC. During the debris removal operations, subsurface pockets of buried transite debris were exposed at several different locations at the AOC. Although removal of subsurface debris was not included in the original limited "RD/RA" scope, visible subsurface transite debris was excavated to the extent possible without undermining and compromising the integrity of Paris-Windham Road (MKM 2004).

The majority of the subsurface transite removed during the limited "RD/RA" was concentrated at the southern end of the AOC; one small pocket of transite debris was located near the central portion of the AOC. Test pits were excavated in 10-ft intervals along the extent of the AOC to ensure all subsurface transite was located. Where transite debris was encountered in the test pits, it was excavated to the extent possible without compromising the integrity of Paris-Windham Road.

Upon completion of the debris removal operations and prior to application of the soil cover, confirmation and contingency samples were collected to evaluate the success of the limited "RD/RA" and provide data for future evaluation of a final remedy. The dump area was divided into 10 equally sized grids, each measuring approximately 40 ft by 20 ft, to facilitate collection of discrete and incremental sampling method (ISM) soil samples (Figure 3-1).

Confirmation sampling activities included collecting 1 discrete surface (0-1 ft bgs) soil sample from each of the 10 grids. Additionally, six confirmatory co-located discrete sediment and surface water samples were collected. Five of these sample locations (PWsw/PWsd-002 through PWsw/PWsd-006) were located within the adjacent Sand Creek floodplain in the intermittent drainage swale between the

dump and Sand Creek, which contained water at that time. One sample location (PWsw/PWsd-001) was located on the northern end of the AOC, outside of the drainage swale (Figure 3-1).

Confirmation samples were analyzed for target analyte list (TAL) metals and asbestos. In addition, 10% of the samples (one sample from Grid 9) were analyzed for a full suite of parameters, including explosives, SVOCs, cyanide, volatile organic compounds (VOCs), propellants, pesticides, PCBs, and asbestos. A full suite of analyses was also performed for sediment/surface water sampling location PWsd/PWsd-004 (Figure 3-1). All six sediment confirmation samples were also analyzed for grain size, and four of the six samples were analyzed for total organic carbon (TOC) (PWsd-001 through PWsd-004). The remaining two sediment samples were not analyzed for TOC due to an error on the chain of custody (MKM 2004).

During confirmatory sampling activities, additional transite debris was found in the excavated areas on the southern portion of the AOC. These small fragments had not been visible during the removal action but were exposed following a heavy rain event. As cited in the *Final Report for Remedial Design/Remedial Action Plan at Paris-Windham Road Dump*, RVAAP stakeholders and the Akron Regional Air Quality Management District agreed to proceed with AOC restoration activities because further excavation had the potential to undermine and compromise the integrity of Paris-Windham Road (MKM 2004). The transite material was subsequently covered in place during AOC restoration activities.

Based on the results of confirmatory sampling and due to the presence of detected SVOCs, MKM Engineers, Inc. (MKM) collected an ISM contingency sample in September 2003 from an approximate 5-ft by 5-ft area surrounding soil sample location PWss-009. In November 2003, a second ISM contingency sample was collected across the 10 sampling grids prior to placement of a final soil cover and AOC restoration.

The excavation area was restored to grade in November 2003 using a combination of clean, hard fill and soil backfill from an Ohio EPA-approved source. Approximately 480 tons of non-contaminated concrete demolition material of various sizes obtained from a stockpile at Load Line 6 was used to create a layer of clean, hard fill for stability in excavated areas, followed by approximately 2 ft (277 tons) of soil backfill material for cover. The area was then seeded and mulched.

3.1.3.1 Limited Remedial Design/Remedial Action Sampling Results

This section summarizes the 2003 sampling results by environmental media at the Dump Along Paris-Windham Road. During preparation of the SC/FFS WP, conditions within the drainage swale at the base of the dump were evaluated, including an AOC walkover. Based on available information summarized below, sediment samples within the drainage swale are considered "dry" sediment in accordance with RVAAP guidelines and are treated as surface soil in the SC (USACE 2010c).

• Surface water in the swale occurs only during occasional storms, periods of snowmelt, or overflow conditions from Sand Creek.

• Samples were collected using a hand trowel in 2003, which, in accordance with the RVAAP Facility-Wide Sampling and Analysis Plan for Environmental Investigations (USACE 2001a), applies when the water depth is less than 6 inches.

3

1

2

• During an August 2009 AOC walkover, SAIC observed no standing water in the swale.

567

• During a November 2011 AOC walkover, SAIC observed water in the drainage swale following an extended precipitation event.

8 9

Thirteen SVOCs, 21 inorganic chemicals, and PCB-1254 were detected in discrete soil samples collected in April 2003 (Tables 3-1 and 3-2). Asbestos was not detected in any of the 10 shallow soil samples.

13

Fifteen inorganic chemicals and nitrocellulose were detected in surface water samples collected in April 2003 (Table 3-3). Asbestos was not detected in any of the six surface water samples.

- One VOC (acetone), 7 SVOCs, 21 inorganic chemicals, PCB-1254, and nitrocellulose were detected in dry sediment samples collected in April 2003 (Tables 3-2 and 3-4). Grain size in the six dry sediment samples was classified as coarse to medium sand. Four dry sediment samples were analyzed for TOC (PWsd-001 through PWsd-004). TOC concentrations in these four samples ranged from 10,000 mg/kg to 34,000 mg/kg. Twenty-three SVOCs were detected in the two contingency
- 22 ISM soil samples collected in the fall of 2003 (Table 3-5).

Table 3-1. Results of Limited "RD/RA" Confirmatory Surface Soil Discrete Samples – Inorganic Chemicals

Station			RVAAP	PWss-001	PWss-002	PWss-003	PWss-004	PWss-005	PWss-006	PWss-007	PWss-008	PWss-009	PWss-010
			Surface Soil	PWss-001-	PWss-002-	PWss-003-	PWss-004-	PWss-005-	PWss-006-	PWss-007-	PWss-008-	PWss-009-	PWss-010-
Sample ID	CAS		Background	0001-SO	0001-SO	0001-SO	0001-SO	0001-SO	0001-SO	0001-SO	0001-SO	0001-SO	0001-SO
Date	Number	Units	Criteria	04/28/03	04/28/03	04/28/03	04/28/03	04/29/03	04/29/03	04/29/03	04/29/03	04/28/03	04/28/03
Inorganic Chemicals													
Aluminum	7429905	mg/kg	17,700	7,500 =	18,000 =	7,000 =	5,600 =	11,000 =	6,500 =	6,500 =	8,600 =	8,600 =	7,700 =
Antimony	7440360	mg/kg	0.96	0.6 =	0.42 U	0.36 U	0.39 U	0.34 U	0.36 U	0.33 U	0.34 U	0.36 U	0.49 =
Arsenic	7440382	mg/kg	15.4	10 =	11 =	11 =	8.5 =	12 =	9.7 =	9.2 =	13 =	12 =	13 =
Barium	7440393	mg/kg	88.4	46 =	150 =	50 =	43 =	180 =	51 =	47 =	49 =	56 =	78 =
Beryllium	7440417	mg/kg	0.88	0.42 J	1.9 =	0.4 J	0.34 J	1.2 =	0.35 J	0.34 J	0.54 =	0.45 =	0.47 =
Cadmium	7440439	mg/kg	0	0.1 J	0.3 =	0.22 J	0.24 U	0.2 U	0.23 U	0.21 U	0.2 U	0.22 U	0.23 U
Calcium ^a	7440702	mg/kg	15,800	2,500 =	55,000 =	2,700 =	3,100 =	39,000 =	2,500 =	1,500 =	4,300 =	2,000 =	1,800 =
Chromium	7440473	mg/kg	17.4	14 =	9.7 =	12 =	8.1 =	8.3 =	9.5 =	10 =	11 =	11 =	12 =
Cobalt	7440484	mg/kg	10.4	6.7 =	4.7 =	6.6 =	5.1 =	4.3 =	5.6 =	6.3 =	6.6 =	7.1 =	7.5 =
Copper	7440508	mg/kg	17.7	16 =	15 =	16 =	9.7 =	19 =	9.6 =	9.3 =	18 =	14 =	16 =
Iron ^a	7439896	mg/kg	23,100	18,000 =	13,000 =	18,000 =	12,000 =	22,000 =	14,000 =	15,000 =	22,000 =	17,000 =	17,000 =
Lead	7439921	mg/kg	26.1	16 =	29 =	15 =	17 =	19 =	15 =	19 =	14 =	14 =	22 =
Magnesium ^a	7439954	mg/kg	3,030	2,100 =	10,000 =	1,800 =	1,400 =	6,100 =	1,500 =	1,600 =	1,900 =	1,800 =	1,800 =
Manganese	7439965	mg/kg	1,450	270 =	1,900 =	520 =	380 =	880 =	410 =	390 =	530 =	490 =	790 =
Mercury	7439976	mg/kg	0.036	0.06 =	0.078 =	0.064 =	0.047 =	0.043 =	0.048 =	0.025 =	0.025 =	0.039 =	0.06 =
Nickel	7440020	mg/kg	21.1	19 =	12 =	15 =	10 =	10 =	11 =	13 =	16 =	21 =	22 =
Potassium ^a	7440097	mg/kg	927	910 =	1,400 =	860 =	780 =	1,100 =	760 =	740 =	970 =	890 =	1,200 =
Silver	7440224	mg/kg	0	0.39 J	0.66 U	0.57 U	0.6 U	0.51 U	0.58 U	0.53 U	0.51 U	0.55 U	0.57 U
Sodium ^a	7440235	mg/kg	123	170 =	480 =	180 =	120 =	380 =	130 =	130 =	190 =	180 =	160 =
Vanadium	7440622	mg/kg	31.1	13 =	9.8 =	13 =	9.5 =	10 =	11 =	11 =	15 =	14 =	14 =
Zinc	7440666	mg/kg	61.8	97 =	78 =	70 =	52 =	100 =	59 =	50 =	63 =	62 =	88 =
						Miscel	laneous						
Asbestos	1332214	Percent	None	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1

Source: Final Report for Remedial Design/Remedial Action Plan at Paris-Windham Road Dump (MKM 2004)

Note: All constituents with at least one detection are shown. All asbestos results are shown as reported

^aEssential human nutrient; not evaluated as a site-related contaminant

Bold text indicates the concentration exceeds background concentration

CAS = Chemical Abstract Service Data Qualifiers: RA = Remedial Action

"=" = Detected at the concentration shown

J = Estimated concentration

U = Not detected at the concentration shown

RVAAP = Ravenna Army Ammunition Plant

RD = Remedial Design

Table 3-2. Results of Limited "RD/RA" Confirmatory Surface Soil and Dry Sediment Discrete Samples – Organic Chemicals

Station			RVAAP	PWss-009	PWsd-004				
Sample ID	CAS		Background	PWss-009-0001-SO	PWsd-004-0001-SD				
Date	Number	Units	Criteria	04/28/03	04/29/03				
		Volatile	Organic Compo	unds					
Acetone	67641	mg/kg	None	0.062 U	0.041 =				
Semi-volatile Organic Compounds									
Acenaphthylene	208968	mg/kg	None	0.13 J	0.83 U				
Anthracene	120127	mg/kg	None	0.12 J	0.83 U				
Benz(a)anthracene	56553	mg/kg	None	1.0 =	0.25 J				
Benzo(a)pyrene	50328	mg/kg	None	1.3 =	0.33 J				
Benzo(b)fluoranthene	205992	mg/kg	None	1.2 =	0.39 J				
Benzo(ghi)perylene	191242	mg/kg	None	0.75 =	0.83 U				
Benzo(k)fluoranthene	207089	mg/kg	None	1.4 =	0.33 J				
Chrysene	218019	mg/kg	None	1.1 =	0.33 J				
Dibenz(<i>a</i> , <i>h</i>)anthracene	53703	mg/kg	None	0.24 J	0.83 U				
Fluoranthene	206440	mg/kg	None	1.7 =	0.44 J				
Indeno(1,2,3-cd)pyrene	193395	mg/kg	None	0.75 =	0.83 U				
Phenanthrene	85018	mg/kg	None	0.32 J	0.83 U				
Pyrene	129000	mg/kg	None	1.4 =	0.44 J				
Polychlorinated Biphenyls									
PCB-1254	11097691	0.23 =	0.086 =						
		N	<i>Iiscellaneous</i>						
Nitrocellulose	9004700	mg/kg	None	NA	2 J				

Source: Final Report for Remedial Design/Remedial Action Plan at Paris-Windham Road Dump (MKM 2004)

Note: All constituents with at least one detection are shown

Data Qualifiers:

"=" = Detected at the concentration shown

J = Estimated concentration

U = Not detected at the concentration shown

CAS = Chemical Abstract Service

NA = Not analyzed

None = No background concentration; all detected values are considered above background concentration

PCB = Polychlorinated Biphenyl

RA = Remedial Action

RD = Remedial Design

RVAAP = Ravenna Army Ammunition Plant

Dump Along Paris-Windham Road

Station			RVAAP Surface	PWsw-001	PWsw-002	PWsw-003	PWsw-004	PWsw-005	PWsw-006
			Water	PWsw-001-	PWsw-002-	PWsw-003-	PWsw-004-	PWsw-005-	PWsw-006-
Sample ID	CAS		Background	0001-SW	0001-SW	0001-SW	0001-SW	0001-SW	0001-SW
Date	Number	Units	Criteria	04/29/03	04/29/03	04/29/03	04/29/03	04/29/03	04/29/03
				Inorga	nic Chemicals				
Aluminum	7429905	mg/L	3.37	0.091 J	0.065 J	0.066 J	0.28 =	0.081 J	0.1 J
Arsenic	7440382	mg/L	0.0032	0.0028 =	0.0058	0.0052 =	0.0074 =	0.0082 =	0.0041 =
Barium	7440393	mg/L	0.0475	0.035 =	0.065 =	0.063 =	0.12 =	0.059 =	0.055 =
Calciuma	7440702	mg/L	41.4	60 =	40 =	40 =	52 =	34 =	23 =
Cobalt	7440484	mg/L	0	0.0013 J	0.05 U	0.005 U	0.0011 J	0.001 J	0.015 J
Copper	7440508	mg/L	0.0079	022 J	022 J	022 J	039 J	024 J	024 J
Iron	7439896	mg/L	2.56	4.3 =	3.7 =	3.9 =	5.3 =	9.4 =	5.1 =
Lead	7439921	mg/L	0	0.002 U	0.002 U	0.002 U	0.0027 =	0.0019 J	0.002 U
Magnesium ^a	7439954	mg/L	10.8	10 =	9.8 =	9.8 =	12 =	8.3 =	6 =
Manganese	7439965	mg/L	0.391	0.32 =	0.27 =	0.26 =	0.51 =	0.47 =	0.56 =
Mercury	7439976	mg/L	0	0.0007 J	0.00009 J	0.00009 J	0.0002 U	0.0001 J	0.00008 J
Nickel	7440020	mg/L	0	0.002 J	0.01 U	0.01 U	0.0024 J	0.0072 J	0.0075 J
Potassium ^a	7440097	mg/L	3.17	1.7 =	4.8 =	4.8 =	5.4 =	4.4 =	3.5 =
Sodium ^a	7440235	mg/L	21.3	8.8 =	8.9 =	8.5 =	9.9 =	5.8 =	4.2 =
Zinc	7440666	mg/L	0.042	0.02 =	0.02 U	0.02 U	0.024 =	0.02 U	0.017 J
				Mis	cellaneous				
Asbestos (total fibers)	1332214	MFL	None	5.900 U	13.02 U	7.812 U	13.02 U	39.06 U	7.812 U
Nitrocellulose	9004700	mg/L	None	NA	NA	NA	0.094 J	NA	NA

Source: Report for Remedial Design/Remedial Action Plan at Paris-Windham Road Dump (MKM 2004)

Note: All constituents with at least one detection are shown

^aEssential human nutrient; not evaluated as a site-related contaminant

Data Qualifiers:

"=" = Detected at the concentration shown

J = Estimated concentration

U = Not detected at the concentration shown

Bold text indicates the concentration exceeds background concentration

CAS = Chemical Abstract Service

MFL = Million Fibers Per Liter

NA = Not analyzed

None = No background concentration; all detected values are considered above background concentration

RA = Remedial Action

RD = Remedial Design

RVAAP = Ravenna Army Ammunition Plant

Station			RVAAP	PWsd-001	PWsd-002	PWsd-003	PWsd-004	PWsd-005	PWsd-006
			Surface Soil	PWsd-001-	PWsd-002-	PWsd-003-	PWsd-004-	PWsd-005-	PWsd-006-
Sample ID	CAS		Background	0001-SD	0001-SD	0001-SD	0001-SD	0001-SD	0001-SD
Date	Number	Units	Criteria	04/29/03	04/29/03	04/29/03	04/29/03	04/29/03	04/29/03
				Inor	ganic Chemicals				
Aluminum	7429905	mg/kg	17,700	8,000 =	9,000 =	7,100 =	8,400 =	9,900 =	7,600 =
Antimony	7440360	mg/kg	0.96	0.47 U	0.72 U	0.59 U	0.73 U	0.73 U	0.46 U
Arsenic	7440382	mg/kg	15.4	6.1 =	2.6 =	5.1 =	3.8 =	8.4 =	5.6 =
Barium	7440393	mg/kg	88.4	53 =	140 =	61 =	110 =	77 =	64 =
Beryllium	7440417	mg/kg	0.88	0.44 J	0.49 J	0.47 J	0.54 J	0.61 J	0.52 J
Cadmium	7440439	mg/kg	0	0.31 U	0.59 =	0.36 U	0.43 U	0.46 U	0.27 U
Calcium ^a	7440702	mg/kg	15,800	2,000 =	4,400 =	2,500 =	4,000 =	1,900 =	1,700 =
Chromium	7440473	mg/kg	17.4	13 =	14 =	12 =	14 =	15 =	17 =
Cobalt	7440484	mg/kg	10.4	5.8 =	5.5 =	5.5 =	6.7 =	6.1 =	5.7 =
Copper	7440508	mg/kg	17.7	17 =	24 =	21 =	25 =	27 =	23 =
Iron	7439896	mg/kg	23,100	14,000 =	12,000 =	18,000 =	15,000 =	15,000 =	17,000 =
Lead	7439921	mg/kg	26.1	19 =	25 =	19 =	18 =	16 =	20 =
Magnesium ^a	7439954	mg/kg	3,030	2,200 =	2,800 =	2,000 =	2,600 =	2,700 =	2,300 =
Manganese	7439965	mg/kg	1,450	99 =	150 =	97 =	120 =	100 =	150 =
Mercury	7439976	mg/kg	0.036	0.059 =	0.08 =	0.058 =	0.073 =	0.077 =	0.05 =
Nickel	7440020	mg/kg	21.1	17 =	24 =	19 =	23 =	37 =	23 =
Potassium ^a	7440097	mg/kg	927	890 =	1,300 =	1,000 =	1,600 =	1,700 =	1,900 =
Silver	7440224	mg/kg	0	0.79 U	1.1 U	0.89 U	1.1 U	1.1 U	0.66 U
Sodium ^a	7440235	mg/kg	123	150 J	210 U	210 =	200 J	230 U	190 =
Vanadium	7440622	mg/kg	31.1	13 =	13 =	14 =	15 =	18 =	14 =
Zinc	7440666	mg/kg	61.8	81 =	120 =	75 =	88 =	99 =	90 =
	Miscellaneous								
Asbestos	1332214	Percent	None	<1	<1	<1	<1	<1	<1

Source: Report for Remedial Design/Remedial Action Plan at Paris-Windham Road Dump (MKM 2004)

Note: All constituents with at least one detection are shown. All asbestos results are shown.

Data Qualifiers:

"=" = Detected at the concentration shown

J = Estimated concentration

U = Not detected at the concentration shown

Bold text indicates the concentration exceeds background concentration

CAS = Chemical Abstract Service

RA = Remedial Action

RD = Remedial Design

RVAAP = Ravenna Army Ammunition Plant

^aEssential human nutrient; not evaluated as a site-related contaminant

Location				Grid 9	Grids 1 through 10
Location	-			PWss-	PWss-
			RVAAP	CONT1-	CONT2-
Sample ID	CAS		Background	00010-SO	00010-SO
Date	Number	Units	Criteria	09/30/03	10/28/03
Date			anic Compound	************	10/20/03
1,4-Dichlorobenzene	106467	mg/kg	None	0.18 U	0.23 =
2-Methylnaphthalene	91576	mg/kg	None	0.0055 J	0.064 =
Acenaphthene	83329	mg/kg	None	0.035 U	0.12 =
Acenaphthylene	208968	mg/kg	None	0.056 =	0.12 =
Anthracene	120127	mg/kg	None	0.041 =	0.22 =
Benz(a)anthracene	56553	mg/kg	None	0.36 =	1.0 =
Benzo(a)pyrene	50328	mg/kg	None	0.46 =	1.4 =
Benzo(b)fluoranthene	205992	mg/kg	None	0.5 =	1.4 =
Benzo(ghi)perylene	191242	mg/kg	None	0.3 =	0.79 =
Benzo(k)fluoranthene	207089	mg/kg	None	0.45 =	1.4 =
Bis(2-ethylhexyl)phthalate	117817	mg/kg	None	0.18 U	0.025 J
Carbazole	86748	mg/kg	None	0.18 U	0.19 =
Chrysene	218019	mg/kg	None	0.41 =	1.2 =
Di-n-butyl phthalate	84742	mg/kg	None	0.18 U	0.041 J
Dibenz(a,h)anthracene	53703	mg/kg	None	0.14 =	0.36 =
Dibenzofuran	132649	mg/kg	None	0.0064 J	0.051 J
Diethyl phthalate	84662	mg/kg	None	0.0093 J	0.067 U
Fluoranthene	206440	mg/kg	None	0.67 =	2.9 =
Fluorene	86737	mg/kg	None	0.011 J	0.1 =
Indeno(1,2,3-cd)pyrene	193395	mg/kg	None	0.31 =	0.7 =
Naphthalene	91203	mg/kg	None	0.035 U	0.039 =
Phenanthrene	85018	mg/kg	None	0.16 =	1.1 =
Pyrene	129000	mg/kg	None	0.62 =	2.0 =

Source: Final Report for Remedial Design/Remedial Action Plan at Paris-Windham Road Dump (MKM 2004) Data Qualifiers:

CAS = Chemical Abstract Service

None = No background concentration; all detected values are considered above background concentration

RA = Remedial Action

RD = Remedial Design

RVAAP = Ravenna Army Ammunition Plant

3

3.1.3.2 Limited Remedial Design/Remedial Action Conclusions

5 6

7

8

9

10

11

The limited "RD/RA" compared results of the 2003 sampling to RVAAP facility-wide background concentrations for inorganic chemicals and USEPA Region 9 preliminary remediation goals (PRGs) in use at the time (MKM 2004). The data screening indicated elevated concentrations of arsenic above its background concentration and/or PRGs in soil, dry sediment, and surface water. Elevated concentrations of SVOCs also were detected in the soil and dry sediment (Grid 9 and dry sediment location PWsd-004). No SVOCs were detected in the surface water sample collected at PWsw-004.

12 Asbestos was below laboratory reporting limits in all soil, dry sediment, and surface water samples.

[&]quot;=" = Detected at the concentration shown

J = Estimated concentration

U = Not detected at the concentration shown

The results of ISM surface soil sampling verified that elevated levels of the following SVOCs were present in soil prior to placement of the soil cover: benz(a)anthracene, benzo(a)pyrene, benzo(b)fluoranthene, indeno(1,2,3-cd) pyrene, and dibenz(a,h)anthracene.

Based on the results of the limited "RD/RA" confirmation samples, MKM recommended an evaluation of risk for the AOC, followed by regulatory AOC closure or additional remedial efforts, as necessary.

3.2 NATURE AND EXTENT OF CONTAMINATION

This section presents the nature and extent of contamination at the Dump Along Paris-Windham Road based upon sampling conducted in 2003 after the removal action. As discussed in Section 3.1.1, data collected during the 1998 RRSE are not included in the current AOC characterization because these data were not intended to be used as definitive evidence of contamination presence or absence or to support quantitative health risk assessment, and they do not reflect current conditions at the AOC.

Available soil data were screened with respect to potential leaching and impacts to groundwater. Numerical modeling for soil leaching or contaminant migration in groundwater was not conducted for this SC as no groundwater data are currently available for the AOC for model validation purposes. A CSM is provided to discuss contaminant sources, migration pathways, and potential receptors.

3.2.1 Site-Related Contaminants

The purpose of identifying site-related contaminants (SRCs) is to determine the presence or absence of contamination that is site-related and above naturally occurring levels. The SRC screening process includes three steps, as outlined in the FWCUG Report (USACE 2010a).

Background screening. The maximum detected concentrations (MDCs) of naturally occurring
inorganic chemicals were compared to the facility-wide background concentrations for RVAAP,
published in the *Phase II Remedial Investigation Report for the Winklepeck Burning Grounds*(USACE 2001b). Inorganic chemicals detected above background concentrations were retained
as SRCs.

• Screening of essential human nutrients. Chemicals considered essential nutrients (e.g., calcium, chloride, iodine, iron, magnesium, potassium, phosphorous, and sodium) are an integral part of the human food supply and are often added to foods as supplements. USEPA recommends these chemicals not be evaluated provided they are present at low concentrations (i.e., only slightly elevated above naturally occurring levels) and toxic only at very high doses (i.e., much higher than those that could be associated with contact at the AOC) (USEPA 1989). Essential nutrients detected near or below their recommended daily allowance/recommended daily intake-based screening levels were eliminated as SRCs.

• **Frequency-of-detection screening.** Analytes detected in less than 5% of the samples may be subject to a weight-of-evidence (WOE) evaluation and may be screened out from further consideration. This frequency-of-detection screen only applies to datasets containing 20 or more samples. No frequency-of-detection screening was performed for this SC/FFS because fewer than 20 discrete samples were available for each of the datasets. Frequency-of-detection screening was not used for ISM samples.

6 7 8

9

10

1

2

3

4

5

SRC screening was conducted separately for discrete and ISM sample results. The SRCs identified for the Dump Along Paris-Windham Road are summarized in Tables 3-6 and 3-7 (soil and surface water, respectively). Appendix Tables B-1 through B-4, present summary statistics and results of the SRC screening process for data included for evaluation in the SC/FFS.

11 12 13

Table 3-6. Soil SRCs

Ι	Discrete Samples	ISM	M Samples		
Inc	organic Chemicals	Inorg	Inorganic Chemicals		
Aluminum	Lead		NA		
Barium	Manganese				
Beryllium	Mercury				
Cadmium	Nickel				
Copper	Silver				
	Zinc				
	<u>SVOCs</u>	S	VOCs		
Acenaphthylene		1,4-Dichlorobenzene	Chrysene		
Anthracene		2-Methylnaphthalene	Dibenz(a,h)anthracene		
Benz(a)anthracene		Acenaphthene	Dibenzofuran		
Benzo(a)pyrene		Acenaphthylene	Diethyl phthalate		
Benzo(b)fluoranthe	ene	Anthracene	Di-n-butyl phthalate		
Benzo(ghi)perylene	e	Benz(a)anthracene	Fluoranthene		
Benzo(k)fluoranthe	ene	Benzo(a)pyrene	Fluorene		
Chrysene		Benzo(b)fluoranthene	Indeno(1,2,3-cd)pyrene		
Dibenz(a,h)anthrac	ene	Benzo(ghi)perylene	Naphthalene		
Fluoranthene		Benzo(k)fluoranthene	Phenanthrene		
Indeno(1,2,3-cd)py	rene	Bis(2-	Pyrene		
Phenanthrene		ethylhexyl)phthalate			
Pyrene	Pyrene				
Pesticides/PCBs					
PCB-1254					
Explosives/Propellants					
Nitrocellulose					
	VOCs				
Acetone					

 $ISM = Incremental \ Sampling \ Method$

NA = Not analyzed

PCB = Polychlorinated Biphenyl

SRC = Site-related Contaminant

SVOC = Semi-volatile Organic Compound

VOC = Volatile Organic Compound

Surface Water					
Inorganic Chemicals	Explosives/Propellants				
Arsenic	Nitrocellulose				
Barium					
Cobalt					
Lead					
Manganese					
Mercury					
Nickel					

SRC = Site-related Contaminant

3.2.2 Occurrence and Distribution of Contaminants

3.2.2.1 Soil

 Eleven inorganic chemicals were identified as SRCs in soil: aluminum, barium, beryllium, cadmium, copper, lead, manganese, mercury, nickel, silver, and zinc. All but three were detected at concentrations above background concentrations at soil sample location PWss-02. Concentrations of mercury and zinc exceeding background concentrations were detected consistently throughout the AOC (13 of 16 and 14 of 16 samples, respectively). Copper and nickel also were frequently detected at concentrations exceeding background concentrations, particularly within the drainage swale (five of six and four of six discrete samples, respectively, from the drainage swale).

Twenty-three SVOCs were identified as SRCs in soil; 13 of those were detected above background concentrations in the April 2003 discrete samples. Ten additional SRCs were identified in the ISM samples collected later that year. Initial SVOC detections in the soil sample collected from Grid 9 suggested the location may have contained a localized release of SVOCs. The contingency ISM sample collected from Grid 9 supports this concept because concentrations of SVOCs are lower in the ISM sample than in the original April grab sample. However, results of the contingency ISM sample collected from Grids 1 through 10 indicate detectable SVOCs, primarily polycyclic aromatic hydrocarbons (PAHs), were present in soil throughout the AOC prior to placement of the soil cover.

Only one VOC (acetone) was detected in sample PWsd-004 collected from the drainage swale; no VOCs were detected in soil sample PWss-09. Additionally, nitrocellulose was detected in sample PWsd-004.

3.2.2.2 Surface Water

Seven inorganic chemicals were identified as SRCs in surface water at the AOC: arsenic, barium, cobalt, lead, manganese, mercury, and nickel. Mercury was detected in all six samples at concentrations exceeding background concentrations; arsenic and barium were each detected in five of six samples at concentrations exceeding background concentrations; and cobalt and mercury were both detected in four of six samples at concentrations exceeding background concentrations. In

general, surface water from locations PWsw-002 and PWsw-003 displayed the lowest concentrations of inorganic chemicals, with two exceptions: the highest detected concentrations of barium and mercury were detected at PWsw-002 and PWsw-003.

No VOCs or SVOCs were detected in surface water sample PWsw-004. Nitrocellulose was detected in this surface water sample as well as in the corresponding dry sediment sample. Asbestos was not detected in any of the surface water samples.

3.2.3 Soil to Groundwater Leaching Screen

To evaluate potential impacts to groundwater from contaminants in soil (inclusive of dry sediment), the April 2003 dataset was compared to the USEPA regional screening level (RSL) (USEPA 2010). When available, the maximum contaminant level (MCL)-based soil screening level (SSL) was used; for analytes without an MCL-based SSL, the risk-based SSL was used. Table 3-8 presents the results of this comparison. Six SVOCs, four inorganic chemicals, and one PCB were identified as contaminant migration chemicals of potential concern (CMCOPCs). Barium, lead, and manganese had the highest frequency of SSL exceedances; however, the SSLs for these three inorganic chemicals are less than their respective RVAAP surface soil background concentrations.

3.2.4 Conceptual Site Model

3.2.4.1 Primary and Secondary Sources

The primary source of contamination at the Dump Along Paris-Windham Road was exposed waste material. However, as part of the 2003 limited "RD/RA," approximately 300 tons of debris was removed, and a minimum 2-ft-thick soil cover was placed over the remaining waste. The soil cover isolates waste and prevents direct exposure. The soil and vegetative cover also prevents direct contact of waste with surface water runoff and helps to limit infiltration of rainfall and snow melt.

Secondary contaminant sources include dry sediment and runoff accumulation points along the drainage swale at the base of the dump. The drainage swale was not excavated or capped with clean soil during the limited "RD/RA" (Figure 3-1). The drainage swale is estimated to be 15 ft wide by 400 ft long (approximately 0.15 acres). In the swale, surface water is present during occasional storms or periods of snow melt or during overflow conditions from nearby Sand Creek. Prior to capping the dump, surface runoff potentially carried contaminants sorbed to particulates and/or contaminants in the dissolved phase to the drainage swale. Percolating rainfall also may have contributed to migration of contaminants from the dump to the drainage swale. Thus, contaminants in surface water and dry sediment in the drainage swale represent secondary sources. Installation of the soil cap minimized direct contact between surface water and waste and reduced infiltration rates through waste material; therefore, the process for continuing contaminant migration to and deposition in the drainage swale has been largely mitigated.

	CAS	Freq of	Maximum Detected	USEPA SSL ^a	SSL	CMCOPC?c	Number >SSL/ Total	
Analyte	Number	Detect	(mg/kg)	(mg/kg)	Type ^b	(yes/no)	Analyses	
V	I.	Volatile	Organic Com	, ,	, <u>, , , , , , , , , , , , , , , , , , </u>	,		
Acetone	67-64-1	1/2	0.041	4.5	Risk	No	0 / 2	
	Semi-volatile Organic Compounds							
Acenaphthylene 208-96-8 1 / 2 0.13 22 Risk No 0 / 2								
Anthracene	120-12-7	1 / 2	0.12	360	Risk	No	0/2	
Benz(a)anthracene	56-55-3	2/2	1	0.01	Risk	Yes	2/2	
Benzo(a)pyrene	50-32-8	2/2	1.3	0.24	MCL	Yes	2/2	
Benzo(b)fluoranthene	205-99-2	2/2	1.2	0.035	Risk	Yes	2/2	
Benzo(ghi)perylene	191-24-2	1 / 2	0.75	0.35	Risk	Yes	1 / 2	
Benzo(k)fluoranthene	207-08-9	2/2	1.4	0.35	Risk	Yes	1 / 2	
Chrysene	218-01-9	2/2	1.1	1.1	Risk	No	0/2	
Dibenz (a,h) anthracene	53-70-3	1 / 2	0.24	0.011	Risk	Yes	1 / 2	
Fluoranthene	206-44-0	2/2	1.7	160	Risk	No	0/2	
Indeno(1,2,3-cd)pyrene	193-39-5	1 / 2	0.75	0.12	Risk	Yes	1 / 2	
Phenanthrene	85-01-8	1 / 2	0.32	120	Risk	No	0/2	
Pyrene	129-00-0	2/2	1.4	120	Risk	No	0/2	
		Inor	rganic Chemic	eals				
Aluminum	7429-90-5	18 / 18	18,000	55,000	Risk	No	0 / 18	
Barium	7440-39-3	18 / 18	180	82	MCL	Yes	4 / 18	
Beryllium	7440-41-7	18 / 18	1.9	3.2	MCL	No	0 / 18	
Cadmium	7440-43-9	4 / 18	0.59	0.38	MCL	Yes	1 / 18	
Copper	7440-50-8	18 / 18	27	46	MCL	No	0 / 18	
Lead	7439-92-1	18 / 18	29	14	MCL	Yes	16 / 18	
Manganese	7439-96-5	18 / 18	1,900	57	Risk	Yes	18 / 18	
Mercury	7439-97-6	18 / 18	0.08	0.1	MCL	No	0 /18	
Nickel	7440-02-0	18 / 18	37	48	Risk	No	0 /18	
Silver	7440-22-4	1 / 18	0.39	1.6	Risk	No	0 /18	
Zinc	7440-66-6	18 / 18	120	680	Risk	No	0 / 18	
	Polychlorinated Biphenyls							
PCB-1254	11097-69-1	2/2	0.23	0.0088	Risk	Yes	2/2	
			Miscellaneous					
Nitrocellulose	9004-70-0	1 / 1	2	24,000	Risk	No	0 / 1	

^aUSEPA SSL for protection of groundwater criteria from regional screening level tables (USEPA 2010)

Bold values indicate the SSL is less than the surface soil background concentration

CAS = Chemical Abstract Service

CMCOPC = Contaminant Migration Chemical of Potential Concern

PCB = Polychlorinated Biphenyl

SSL = Soil Screening Level

2

4 5

7

USEPA = United States Environmental Protection Agency

3.2.4.2 Migration Pathways and Receptors

The primary contaminant migration pathway at the AOC is surface water runoff. The steep topography and surface water flow patterns through the drainage swale facilitate contaminant migration from the AOC into nearby Sand Creek, which is located at distances ranging 30-170 ft. Infiltration of rainfall through remaining waste, with discharge into the drainage swale at the base of

^bMaximum contaminant level criteria were used when available; otherwise, risk-based criteria are shown

^cConstituent is considered a CMCOPC when one or more detected concentrations exceed the SSL

the slope may still occur; however, the soil cover and current dense vegetation maximize evapotranspiration rates (particularly during the growing season) and help minimize contaminant migration via this pathway. Surface water samples collected in 2003 immediately following the limited "RD/RA" indicated the presence of inorganic SRCs but did not contain explosives, VOCs, SVOCs, pesticides, or PCBs.

Groundwater may be a potential migration pathway; although, the occurrence of contaminants in groundwater is not documented by sampling because of a lack of monitoring wells. The January 2010 unconsolidated aquifer facility-wide potentiometric map (EQM 2010) indicates the potentiometric head in the vicinity of the Dump Along Paris-Windham Road likely falls between 950 and 975 ft above mean sea level (amsl). Sand Creek lies approximately 945 ft amsl to the west of the AOC, and the sample grid areas outlined in the limited "RD/RA" range from 950 to 960 ft amsl (Figure 3-1). Sand Creek, which lies approximately 30 ft to the north of the AOC on the northern end to about 170 ft west of the AOC on the southern end, is assumed to be an expression of the water table and the downgradient receptor for groundwater discharge. Therefore, available data indicate relatively short vertical (5-15 ft) and horizontal (less than 200 ft) flow paths exist for contaminant migration to the saturated zone and lateral transport to Sand Creek.

Results of the RVAAP facility-wide biological and water quality study sampling at station S9 at river mile (RM) 1.9 (Figure 3-1) were evaluated to determine possible surface water and groundwater contaminant migration to Sand Creek (USACE 2005a). This monitoring station is located at the southwest corner of the Paris-Windham Road bridge over Sand Creek and is immediately downstream of the AOC. As discussed in Section 4.4, results of chemical and biological samples collected during the facility-wide surface water study at this sampling station indicate any potential groundwater or surface water contamination due to past activities at the Dump Along Paris-Windham Road is not contributing to a decline in water quality in Sand Creek immediately downstream of the AOC.

3.2.4.3 <u>Uncertainties and Data Gaps</u>

Characterization of the nature and extent of contamination in soil (including dry sediment) and surface water at the Dump Along Paris-Windham Road is considered sufficient for risk assessment (Section 4.0) and development and analysis of remedial alternatives in the FFS (Sections 8.0 and 9.0). No residual data gaps are identified for these media.

- 36 As previously stated, groundwater in the vicinity of the AOC has not been characterized to date.
- 37 Therefore, uncertainty exists with respect to predicted impacts and the potential for contaminant
- 38 migration in groundwater. Groundwater will be assessed in a future report as part of the RVAAP
- 39 Facility-Wide Groundwater AOC (RVAAP- 66).

THIS PAGE INTENTIONALLY LEFT BLANK.




Figure 3-1. Limited "RD/RA" Location Map

THIS PAGE INTENTIONALLY LEFT BLANK.

4.0 HUMAN HEALTH RISK ASSESSMENT AND ECOLOGICAL RISK ASSESSMENT

Based on the confirmation sampling data, the limited "RD/RA" recommended that an evaluation of risk be performed for the AOC to determine if additional removal actions were required or if the limited "RD/RA" actions were sufficient to allow for regulatory AOC closure (MKM 2004). This portion of the SC presents the results of the recommended risk assessment. Potential exposure pathways and receptors, based on the CSM discussed in Section 3.0, are shown in the conceptual site exposure model (Section 4.1). Data evaluation for use in the risk assessments is described in Section 4.2. Methods and results are discussed in the HHRA (Section 4.3) and the ERA (Section 4.4).

4.1 CONCEPTUAL SITE EXPOSURE MODEL

The limited "RD/RA" for the Dump Along Paris-Windham Road consisted of removing all existing unconsolidated surface debris and some subsurface debris. Soil samples were collected from the excavated area following excavation and prior to AOC restoration. In addition, six co-located sediment/surface water samples were collected from a drainage swale at the base of the toe slope and within the neighboring floodplain to characterize impacts associated with runoff. The excavation area was restored to grade using a combination of clean, hard fill and Ohio EPA-approved soil backfill. The area was seeded and mulched. Reconnaissance data from an AOC walkover conducted by SAIC in August 2009 show extensive healthy re-vegetation of the area (Appendix A).

Figure 4-1 illustrates the conceptual site exposure model for the AOC. The primary source of contamination is the residual dump material located in the AOC. The potential migration pathway is surface water runoff with three secondary sources: surface water in the drainage swale, dry sediment/soil in the drainage swale, and subsurface soil and debris under the layers of clean hard fill. Human receptors evaluated for the Dump Along Paris-Windham Road are the National Guard Range Maintenance Soldier (representative receptor), Adult and Juvenile Trespassers, and Resident Farmer Adult and Child, as described in Section 4.3.3. The human receptor exposure pathways are based on the FWCUG Report (USACE 2010a). Potential ecological receptors are aquatic organisms (such as fish and stream macroinvertebrates) for surface water and terrestrial organisms (such as plants and wildlife) for soil.

In the swale to the west of the AOC, surface water from storm runoff, periods of snow melt, and occasional overflow conditions from nearby Sand Creek has been observed on a periodic basis. The presence of surface water in the drainage swale is most prevalent during seasonally wet periods of the year. Although field observations show surface water exists in the drainage swale on an intermittent basis, it is present at sufficient frequency and duration to be evaluated as a potential exposure pathway for human and ecological receptors. Sand Creek flows northward about 170 ft west of the south-central edge of the AOC. At the northern end of the AOC, the former dump limits are about 30 ft from Sand Creek near the bridge on Paris-Windham Road. Receptors also may be exposed to soil (inclusive of dry sediment for this evaluation). Exposure to sampled soil and residual waste in the

1 2	area that was excavated during the limited "RD/RA" is precluded by the presence of the clean, hard fill and a minimum of 2 ft of clean soil backfill placed on top of the excavated grids (MKM 2004).




Figure 4-1. Conceptual Site Exposure Model

4.2 DATA EVALUATION FOR HUMAN HEALTH AND ECOLOGICAL RISK ASSESSMENTS

The purpose of this data evaluation is to develop a set of chemical data suitable for use in the HHRA and ERA. Data were evaluated to establish data aggregates and identify a list of SRCs.

456

7

1 2 3

Data collected at the Dump Along Paris-Windham Road were aggregated by environmental media (e.g., surface soil and surface water), exposure unit (EU), and sample type (i.e., discrete or ISM). A description of the media to which human and ecological receptors are potentially exposed follows.

8 9

4.2.1 Data Aggregate – Soil

101112

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

Soil data at the Dump Along Paris-Windham Road include 10 discrete soil samples, 2 ISM soil samples, and 6 sediment samples collected in 2003. The dump area was divided into 10 equally sized grids (40 ft by 20 ft) to facilitate collection of discrete and ISM soil samples (Figure 3-1). One discrete soil sample was collected from each grid. One ISM sample was collected at Grid 9, and one ISM sample was collected to evaluate the extent of SVOC contamination over the entire AOC (i.e., across all 10 grids). All soil samples were collected from 0-1 ft bgs prior to AOC restoration. Six discrete sediment samples were collected from a drainage swale adjacent to the dump area. The 2003 limited "RD/RA" did not differentiate wet or dry sediment. Sampling logs indicate that sediment samples were collected from the 0 to 1 ft bgs interval. Surface water only occurs in the swale on an intermittent basis. For instance, during an August 2009 walkover, SAIC noted the sediment in the drainage swale had high moisture content, but no standing water was observed. However, during a November 2011 walkover following a rainfall event, water was observed. Based on the intermittent nature of surface water at the AOC and the sampling interval, the 2003 sediment samples are considered dry sediment. Evaluation of these samples as dry sediment/soil is a conservative approach because the representative receptor (Range Maintenance Soldier) is exposed to soil but is not exposed to wet sediment. Further discussion of characteristics and habitat within the drainage swale is presented in the ERA (Section 4.4).

28 29 30

31

Soil at the Dump Along Paris-Windham Road was evaluated as three EUs, based on the potential for exposure (i.e., exposed soil versus soil covered by fill) and sample coverage (i.e., AOC-wide ISM area). The three EUs are as follow:

323334

35

36

37

• **Fill Area EU** – The middle of the dump area (characterized by discrete sample locations SS-005, SS-006, SS-007, SS-008, and SS-009 and ISM sample location PWss-CONT1) was excavated and covered with at least 2 ft of clean fill. These samples were collected from 0-1 ft bgs prior to restoration. This sampled soil is currently under at least 2 ft of clean fill; therefore, it represents subsurface soil.

38 39 40

41

42

43

• **Surface Area EU** – The northern and southern ends of the dump area (characterized by discrete sample locations SS-001, SS-002, SS-003, SS-004, and SS-010) and the drainage swale (characterized by samples SD-001, SD-002, SD-003, SD-004, SD-005, and SD-006) lay outside the limited "RD/RA" excavation area (Figure 3-1). Limited, if any, backfill/cover soil was placed

3 4 5

> 6 7

1

2

AOC-Wide EU – One ISM sample was collected across the entire soil grid (i.e., all 10 grid areas). As with the discrete samples, this sample was collected following excavation and prior to restoration to grade. Portions of the sampled area were subsequently filled. Therefore, this sample (PWss-CONT2) represents a combination of surface and subsurface conditions at the AOC.

8 9 10

Samples included in the risk assessment datasets for soil are listed in Table 4-1.

11 12 13

Table 4-1. Risk Assessment Datasets for Soil

			Dept	h (ft bgs)
Station	Sample ID	Date	At Time of Sampling	Following AOC Restoration
	Fill Area EU: Discret	e Subsurface S	Soil Samples	
PWss-005	PWss-005-0001-SO	04/29/03	0-1	>2
PWss-006	PWss-006-0001-SO	04/29/03	0-1	>2
PWss-007	PWss-007-0001-SO	04/29/03	0-1	>2
PWss-008	PWss-008-0001-SO	04/29/03	0-1	>2
PWss-009	PWss-009-0001-SO	04/28/03	0-1	>2
	Fill Area EU: ISM	Subsurface So	oil Sample	
PWss-CONT1	PWss-CONT1-0001-SO	09/30/03	0-1	>2
	Surface Area EU: Dis	crete Surface :	Soil Samples	
PWsd-001	PWsd-001-0001-SD	04/29/03	0-1	0-1
PWsd-002	PWsd-002-0001-SD	04/29/03	0-1	0-1
PWsd-003	PWsd-003-0001-DUP	04/29/03	0-1	0-1
PWsd-003	PWsd-003-0001-SD	04/29/03	0-1	0-1
PWsd-004	PWsd-004-0001-SD	04/29/03	0-1	0-1
PWsd-005	PWsd-005-0001-SD	04/29/03	0-1	0-1
PWsd-006	PWsd-006-0001-SD	04/29/03	0-1	0-1
PWss-001	PWss-001-0001-SO	04/28/03	0-1	0-1
PWss-002	PWss-002-0001-SO	04/28/03	0-1	0-1
PWss-003	PWss-003-0001-SO	04/28/03	0-1	0-1
PWss-004	PWss-004-0001-DUP	04/28/03	0-1	0-1
PWss-004	PWss-004-0001-SO	04/28/03	0-1	0-1
PWss-010	PWss-010-0001-SO	04/28/03	0-1	0-1
AOC	C-Wide EU: ISM Combined	Surface and S	ubsurface Soil Se	ample
PWss-CONT2	PWss-CONT2-0001-SO	10/28/03	0-1	0 to >2

AOC = Area of Concern

bgs = below ground surface

EU = Exposure Unit

ISM = Incremental Sampling Method

14

4.2.2 Data Aggregate – Surface Water

15 16 17

18

Intermittent surface water at the Dump Along Paris-Windham Road is limited to a long, narrow drainage swale downslope of the excavated dump area. Clean backfill soil was not placed in the

drainage swale following the limited "RD/RA." The eastern edge of the drainage swale is estimated to be 15 ft wide by 400 ft long (approximately 0.14 acres). Sand Creek flows northward about 170 ft west of the south-central portion of the dump and flows as close as 30 ft at the northern end of the AOC. Six surface water samples were collected from the drainage swale at the base of the toe slope and within the neighboring floodplain in 2003. No data more recent than 2003 exists for surface water. The use of these samples in the HHRA and ERA is protective because the samples were collected prior to the placement of the soil and vegetation cover, and the potential for contaminant migration from the AOC and exposures was higher than following the completion of the limited "RD/RA."

Surface water at the Dump Along Paris-Windham Road was evaluated as a single EU (also referred to as the Surface Water EU in this SC/FFS). Samples included in the risk assessment dataset for surface water are listed in Table 4-2. Field duplicate sample PWsw-003-0001-F was not excluded from the dataset.

Table 4-2. Risk Assessment Dataset for Surface Water

Station	Sample ID	Date
PWsw-001	PWsw-001-0001-S	04/29/03
PWsw-002	PWsw-002-0001-S	04/29/03
PWsw-003	PWsw-003-0001-F	04/29/03
PWsw-003	PWsw-003-0001-S	04/29/03
PWsw-004	PWsw-004-0001-S	04/29/03
PWsw-005	PWsw-005-0001-S	04/29/03
PWsw-006	PWsw-006-0001-S	04/29/03

4.3 HUMAN HEALTH RISK ASSESSMENT

As described in the SC/FFS WP (USACE 2010c), the HHRA conducted for this SC/FFS consists of the following three steps:

• Evaluate representative AOC-specific receptors and exposure media.

• Identify COCs using appropriate RVAAP risk-based values, FWCUGs, and background concentrations.

• Identify the specific FWCUGs that are applicable for this SC/FFS and evaluate the nature and extent of COCs.

Recently, the RVAAP project team adopted a streamlined approach for performing risk-based decision making at RVAAP, taking advantage of the experience gained through previously completed work. To aid in streamlined decision making, the FWCUG Report (USACE 2010a) was developed to support environmental remediation of the remaining AOCs at RVAAP to complete final transfer of the land to OHARNG. The FWCUG Report contains calculated FWCUGs and guidance for their application to accelerate the decision-making process for the remaining AOCs, taking advantage of

the fact that many of the risk assessment inputs and decisions for the facility have already been agreed to by stakeholders through the application of the CERCLA process over the past 10 years. Most of

the agreed-to risk assessment methods have been documented in the FWHHRAM.

5 The streamlined approach to risk decision-making presented in the FWCUG Report (USACE 2010a) 6 is as follows.

8 1. Using the risk assessment process presented in the FWHHRAM (and appended by information in the Final White Paper provided in Appendix B of the FWCUG Report), develop FWCUGs for all COPCs identified from the facility-wide dataset at RVAAP.

12 2. Perform RI characterization sampling and analysis to establish the baseline chemical concentrationswithin an AOC.

Perform data analysis and mapping to identify COPCs, determine EUs, and calculate exposure point
concentrations (EPCs) for each COPC, following the requirements in the FWHHRAM and further
clarified in the position paper developed by USACE, Louisville District provided in Appendix B of
the FWCUG Report.

4. Compare EPCs to the FWCUGs to determine COCs.

5. Perform the FS, PP, and ROD to address any identified COCs.

Step 1 of this process (develop FWCUGs) has been completed in the FWCUG Report. The results of Step 2 (characterization sampling) and Step 3 (mapping and data analysis) are documented in the *Final Report for Remedial Design/Remedial Action Plan at Paris-Windham Road Dump* (MKM 2004) and summarized in Section 3.0 of this SC/FFS. The remainder of the process (Steps 4 and 5) is provided in the following subsections and follows the four steps for a streamlined risk assessment outlined in Figure 4-2: (1) identify media of concern (Section 4.3.1); (2) identify COPCs (Section 4.3.2); (3) determine the AOC land use and appropriate receptors (Section 4.3.3); and (4) compare to appropriate FWCUGs to identify COCs (Section 4.3.4). Section 4.3.5 identifies the specific FWCUGs that are applicable for this SC/FFS and provides an assessment of the nature and extent of

COCs.

4.3.1 Identify Media of Concern

Media of concern at the Dump Along Paris-Windham Road are surface soil, subsurface soil, and surface water, as described in Section 4.2. As described in Section 1.1, groundwater will be addressed by the U.S. Army under a future decision for the RVAAP Facility-Wide Groundwater AOC (RVAAP-66).

4.3.2 Identify Chemicals of Potential Concern

Section 3.2 presents the screening criteria used to identify SRCs for the Dump Along Paris-Windham Road. Details of the SRC screening for each exposure medium and sample type are provided in Appendix Tables B-1 (discrete soil samples at the Fill Area EU), B-2 (discrete soil samples at the Surface Area EU), B-3 (ISM soil samples), and B-4 (surface water). COPCs were identified as a subset of SRCs present at concentrations that indicate potential impacts to human receptors. The COPC screen follows the approach specified in the FWCUG Report (USACE 2010a) and summarized here.

To determine COPCs, the MDCs of all SRCs were screened against the chemical-specific FWCUGs at a target cancer risk of 1E-06 and a non-carcinogenic target hazard quotient (HQ) of 0.1 for the Resident Farmer Adult, Resident Farmer Child, and National Guard Trainee. These are the most conservative FWCUGs available and are used for all AOCs at RVAAP regardless of the current or future land use. If no FWCUGs were developed for an SRC, the USEPA residential RSL [at a target risk (TR) of 1E-06 or an HQ of 0.1] was used for this screen. As part of the conservative screening approach for identifying COPCs, the FWCUG for hexavalent chromium (the more toxic of the two chromium types evaluated) was used at this stage.

Details of the COPC screening for each exposure medium are provided in Appendix Tables B-1 (soil in the Fill Area EU), B-2 (soil in the Surface Area EU), B-3 (ISM sampled soil), and B-4 (surface water). These tables include all carcinogenic and non-carcinogenic risk-based FWCUG or RSL values for each chemical. SRCs were identified as COPCs if the MDC exceeded the most protective (i.e., lowest) FWCUG. The COPCs identified for the media of concern at the Dump Along Paris-Windham Road are summarized in Table 4-3 and following sections.




Figure 4-2. Risk Assessment Input to Support Remediation Decisions

4.3.2.1 Chemicals of Potential Concern in the Fill Area Exposure Unit for Soil

Thirty-one chemicals were detected in discrete soil samples collected in the excavated area of the former dump prior to filling and grading during restoration. These samples currently represent subsurface soil because at least 2 ft of clean fill was added to this area after these samples were collected. Nineteen of these chemicals were identified as SRCs. Risk-based screening identified six COPCs in this soil: five SVOCs [benz(a)anthracene, benzo(a)pyrene, benzo(b)fluoranthene, dibenz(a,h)anthracene, and indeno(1,2,3-cd)pyrene] and one PCB (PCB-1254).

8

1

2

4

5

6 7

4.3.2.2 Chemicals of Potential Concern in the Surface Area Exposure Unit for Soil

101112

13

14

Thirty-one chemicals were detected in discrete soil samples collected in the area of the former dump not covered by fill during restoration. Twenty-one of these chemicals were identified as SRCs. Risk-based screening identified five COPCs in this soil: two inorganic chemicals (aluminum and manganese) and three SVOCs [benz(a)anthracene, benzo(a)pyrene, and benzo(b)fluoranthene].

15 16 17

4.3.2.3 <u>Chemicals of Potential Concern in Incremental Sampling Method Soil Samples</u>

18 19

20

21

22

Two ISM soil samples were collected from the same areas from which discrete samples had previously been collected. These ISM samples were analyzed for SVOCs only. Twenty-three SVOCs were detected, and all were identified as SRCs. Risk-based screening identified five COPCs in these samples: benz(a)anthracene, benzo(a)pyrene, benzo(b)fluoranthene, dibenz(a,h)anthracene, and indeno(1,2,3-cd)pyrene.

2324

4.3.2.4 Chemicals of Potential Concern in Surface Water

252627

28

29

Sixteen chemicals were detected in surface water samples collected from the drainage swale at the base of the toe slope and within the neighboring floodplain; eight of these chemicals were identified as SRCs. Risk-based screening identified two inorganic chemicals as COPCs in surface water: arsenic and cobalt.

303132

Table 4-3. Summary of COPCs

Fill Area EU	Surface Area EU	ISM Samples	Coords a Walan
(Subsurface Soil)	(Surface Soil)	(Surface and Subsurface Soil)	Surface Water
<u>SVOCs</u>	Inorganic Chemicals	SVOCs	Inorganic Chemicals
Benz(a)anthracene	Aluminum	Benz(a)anthracene	Arsenic
Benzo(a)pyrene	Manganese	Benzo(a)pyrene	Cobalt
Benzo(b)fluoranthene		Benzo(b)fluoranthene	
Dibenz(a,h)anthracene	SVOCs	Dibenz(a,h)anthracene	
Indeno(1,2,3-cd)pyrene	Benz(a)anthracene	Indeno(1,2,3-cd)pyrene	
	Benzo(a)pyrene		
Pesticides/PCBs	Benzo(b)fluoranthene		
PCB-1254			

COPC = Chemical of Potential Concern

PCB = Polychlorinated Biphenyl

EU = Exposure Unit

SVOC = Semi-volatile Organic Compound

ISM = Incremental Sampling Method

4.3.3 Determine Area of Concern Land Use and Appropriate Receptors

The Dump Along Paris-Windham Road is located in the east-central portion of RVAAP, along a steep embankment on the west side of Paris-Windham Road between the bridge over Sand Creek and the intersection of Paris-Windham Road with Remalia Road. Sand Creek is located to the west and north at distances ranging from approximately 30 ft (north end of the AOC) to 170 ft (south-central portion of the AOC). The following information was considered when identifying representative receptors for evaluation in this SC/FFS:

• No specific development project is currently identified by the OHARNG for this AOC.

• Any proposed utilities would be located on the east side of Paris-Windham Road due to the presence of transite on the west side of the road in this area.

• The area is not fenced and does not have any additional security measures beyond those in place for the entire facility.

• The dump area is small and located on a steep slope. It begins at the berm to the west of Paris-Windham Road, and there is a floodplain at the bottom.

Based on these considerations, the RAFLU for the AOC is as follows:

• Dump Along Paris-Windham Road (RVAAP-51) Restricted Access OHARNG Military Use and Training Land Use.

• Representative Receptor – National Guard Range Maintenance Soldier

Dump Along Paris-Windham Road is described in more detail below.

RVAAP is a controlled-access facility that is fenced and patrolled by security personnel. Full-time OHARNG, BRAC, and contractor staff work at the facility. Military training and operations are conducted at the facility. The AOC is located in the eastern-central portion of the facility. The AOC is not currently used for military training activities but may receive periodic foot traffic. The OHARNG projected future land use for the AOC is Restricted Access due to residual asbestos at the AOC. The most representative receptor is the National Guard Range Maintenance Soldier. This anticipated future land use, in conjunction with the evaluation of agricultural-residential land uses and associated receptors, form the basis for identifying COCs in this RI. Residential land use, specifically the Resident (adult and child) Farmer scenario, is included to evaluate COCs for unrestricted land use at the AOC as required by the CERCLA process and as outlined in the FWHHRAM (USACE 2005); however, the topography of the area (i.e., steep slope and floodplain) precludes Residential Land Use. Because the AOC is located immediately adjacent to a primary road, trespassers may potentially visit the AOC; therefore, Adult and Juvenile Trespassers are also considered. The exposure assumptions for the Range Maintenance Soldier are also protective of the Adult and Child trespasser. Per guidelines in the FWCUG Report (USACE 2010a), the application of these receptor scenarios to the

• Range Maintenance Soldier – This receptor represents OHARNG personnel who may occasionally visit the AOC in connection with any adjacent range areas or for other routine or occasional monitoring of the area. This receptor is assumed to contact shallow surface soil, including dry sediment (0-1 ft bgs) and subsurface soil (>2 ft bgs). These two soil intervals represent the 0-4 ft deep surface soil interval as defined for the Range Maintenance Soldier in the FWCUG Report (USACE 2010a).

• Adult and Juvenile Trespassers – These receptors are assumed to contact shallow surface soil, including dry sediment (0-1 ft bgs) and surface water in the drainage swale at the base of the slope of the former dump. The Adult Trespasser is assumed to visit the AOC 75 days/year (USACE 2010a) and thus is also protective of "foot traffic" by National Guard Trainees.

• Resident Farmer Adult and Child – These receptors are generally assumed to contact shallow surface soil (0-1 ft bgs), subsurface soil, and surface water. This AOC is located on a steep embankment, is bordered by a floodplain and a road, and is not suitable for Residential Land Use (e.g., a house cannot be built directly on the AOC). However, for evaluation of Residential Land Use, a residence is assumed to be built across the road from the AOC with a yard that encompasses the road and hillside. Based on this scenario, the Resident Farmer is assumed to contact shallow surface soil, including dry sediment (0-1 ft bgs), and intermittent surface water in the drainage swale at the base of the toe slope of the former dump. Exposure to subsurface soil is not included because the foundation of a house would have to be located outside the AOC.

4.3.4 Compare to Appropriate Facility-Wide Cleanup Goals

The comparison to FWCUGs and determination of COCs follows guidance presented in Appendix B of the FWCUG Report (USACE 2010a). The screening process is as follows:

 • Select the FWCUGs for the planned National Guard end-use representative receptor (Range Maintenance Soldier) and Adult and Juvenile Trespassers at the Dump Along Paris-Windham Road. Also select the FWCUGs for the Resident Farmer Adult and Child receptors to evaluate an unrestricted land use scenario corresponding to a TR of 1E-05 and target HQ of 1.0.

 Report all carcinogenic- and non-carcinogenic-based FWCUGs for each COPC for all appropriate receptors (i.e., Range Maintenance Soldier, Adult and Juvenile Trespassers, and Resident Farmer Adult and Child).

Report critical effects and target organs for each of the non-carcinogenic-based FWCUGs.

- Complete a comparison of the selected FWCUG to the EPC, including a sum-of-ratios (SOR).
- o For non-carcinogens, compare the EPC to the target HQ FWCUG. Sum the ratios of the EPC/FWCUG for COPCs that affect similar target organs.
- o For carcinogens, compare the EPC to the TR FWCUG. Sum the ratios of the EPC/FWCUG for all carcinogens.

- The COPC is identified as a COC for a given receptor if
 - The EPC exceeds the more protective FWCUG for either the 1E-05 target cancer risk or the 1.0 target HQ; or
 - o The SOR for all carcinogens or all non-carcinogens that may affect the same organ is greater than 1.0; chemicals contributing at least 10% to the SOR also were considered COCs.

The selection of FWCUGs, calculation of EPCs for comparison to the FWCUGs, and results of the identification of Dump Along Paris-Windham Road COCs are detailed in the following sections.

7 8 9

1

2

3

4

5

6

4.3.4.1 <u>Selection of Appropriate Facility-Wide Cleanup Goals for the Dump Along Paris-</u> Windham Road

101112

13

14

15

16

The land use/receptor combination for selecting FWCUGs for the Dump Along Paris-Windham Road evaluation is a safety danger zone (SDZ) for a proposed future range complex (OHARNG 2008b), with the Range Maintenance Soldier as the representative receptor. The Adult and Juvenile Trespassers are also evaluated for this AOC. FWCUGs were identified for soil and surface water. In addition to this planned OHARNG land use, the Resident Farmer Adult and Child receptor FWCUGs were also used to evaluate a baseline scenario.

17 18 19

20

FWCUGs for these receptors from the FWCUG Report (USACE 2010a) are provided in Tables 4-4 and 4-5 (soil and surface water, respectively) for all COPCs. The critical effect or target organ associated with the toxicity values used to calculate the FWCUGs are also provided in these tables.

212223

Table 4-4. FWCUGs for COPCs in Soil

		Range Ma	aintenance				
		Sol	dier	Trespasser ^a		Resident Farmer ^b	
COPC	Target Organ	HQ = 1.0	TR = 1E-05	HQ = 1.0	TR = 1E-05	HQ = 1.0	TR = 1E-05
Aluminum	Reproductive ^c	1,000,000	NA	1,000,000	NA	76,800	NA
Manganese	CNS	204,672	NA	220,293	NA	2,927	NA
Benz(a)anthracene	NA	NA	26.2	NA	11.3	NA	2.21
Benzo(a)pyrene	NA	NA	2.62	NA	1.13	NA	0.221
Benzo(b)fluoranthene	NA	NA	26.2	NA	11.3	NA	2.21
Dibenz(a,h)anthracene	NA	NA	2.62	NA	1.13	NA	0.221
Indeno(1,2,3-cd)pyrene	NA	NA	26.2	NA	11.3	NA	2.21

^aTrespasser FWCUGs are the smaller of the adult and juvenile values for each COPC

CNS = Central Nervous System

COPC = Chemical of Potential Concern

 $FWCUG = Facility\text{-}Wide\ Cleanup\ Goal$

HQ = Hazard Quotient

NA = Not available

TR = Target Risk

^bResident Farmer FWCUGs are the smaller of the adult or child values for each COPC

^cNeurotoxicity in offspring

			FWCUG (mg/L)				
		Trespasser ^a		Trespasser ^a		Resident	: Farmer ^b
COPC	Target Organ	HQ = 1.0	TR = 1E-05	HQ = 1.0	TR = 1E-05		
Arsenic	Skin	0.705	0.0415	0.046	0.011		
Cobalt	Thyroid/Lung	NA	NA	NA	NA		

^aTrespasser FWCUGs are the smaller of the adult and juvenile values for each COPC

COPC = Chemical of Potential Concern

FWCUG = Facility-Wide Cleanup Goal

HQ = Hazard Quotient

NA = Not available

TR = Target Risk

4.3.4.2 <u>Exposure Point Concentrations for Comparison to Facility-Wide Cleanup Goals</u>

For discrete soil and surface water data, EPCs were calculated from the results of all of the discrete samples collected from each EU (listed in Tables 4-1 and 4-2), following the method and equations provided in the FWHHRAM (USACE 2005b). The EPC was either the 95% upper confidence limit (UCL $_{95}$) of the mean or the MDC, whichever value was lowest. If the UCL $_{95}$ could not be determined, the EPC was the MDC. For ISM soil data, the EPC was the detected concentration in each ISM sample.

11 12

2

4 5

6

7

8

9

10

4.3.4.3 <u>Identification of Dump Along Paris-Windham Road Chemicals of Concern: Range</u> <u>Maintenance Soldier Scenario</u>

131415

16 17

18

The Range Maintenance Soldier is assumed to contact shallow surface soil (0-1 ft bgs at the Surface Area EU) and subsurface soil (> 2 ft bgs at the Fill Area EU). This receptor is not expected to contact surface water. COC screening for the Range Maintenance Soldier receptor is detailed in Appendix Tables B-5, B-6, and B-7 (Surface Area EU, Fill Area EU, and ISM soil samples, respectively) and summarized below.

192021

No COCs were identified in the Surface Area EU, Fill Area EU, or in the ISM soil samples. The EPCs for all COPCs are below the FWCUGs for this receptor.

2324

22

No COCs were identified based on the SOR analysis, as summarized below.

2526

• None of the COPCs identified in soil have similar toxic endpoints; therefore, no non-cancer SOR was calculated.

272829

30 31

32

• Five COPCs [benz(a)anthracene, benzo(a)pyrene, benzo(b)fluoranthene, dibenz(a,h)anthracene, and indeno(1,2,3-cd)pyrene] in soil have FWCUGs for the cancer endpoint. An SOR was calculated for these potential carcinogens for the EPCs in the Surface Area EU and Fill Area EU, as well as for each of the ISM samples. All calculated SORs are ≤1; therefore, no additional COCs were identified.

^bResident Farmer FWCUGs are the smaller of the adult and child values for each COPC

4.3.4.4 <u>Identification of Dump Along Paris-Windham Road Chemicals of Concern: Trespasser</u> Scenario

Trespassers are assumed to contact shallow surface soil and surface water in the drainage conveyance at the base of the slope of the former dump. COC screening for the Trespasser scenario is detailed in Appendix Tables B-8 (Surface Area EU), B-9 (AOC-wide ISM soil sample), and B-12 (surface water) and summarized below. The most conservative (smallest) FWCUGs for the Adult and Juvenile Trespassers were used in the COC screening.

Soil COCs for the Trespasser scenario are summarized below and in Table 4-6.

• No COCs were identified for the Trespasser scenario in the Surface Area EU. All EPCs are less than FWCUGs for the Adult and Juvenile Trespassers.

• No COCs were identified for the Trespasser scenario in the AOC-wide ISM sample. Benzo(a)pyrene was detected at a concentration of 1.4 mg/kg in PWss-CONT2. The detected concentration slightly exceeds the FWCUG for the Adult Trespasser (1.13 mg/kg) and is below the FWCUG for the Juvenile Trespasser (4.5 mg/kg). Sample PWss-CONT2 was collected from across all 10 soil sample grids prior to filling and grading of the AOC. Approximately one-half the sampled area was covered with at least 2 ft of clean fill after this sample was collected; therefore, this sample does not entirely represent surface conditions. Because the Trespasser is not exposed to subsurface soil, and considering the information presented above, benzo(a)pyrene is not identified as a COC for this receptor.

No additional COCs were identified based on the SOR analysis as summarized below.

• None of the COPCs identified in soil have similar toxic endpoints; therefore, no non-cancer SOR was calculated.

• Five COPCs [benz(a)anthracene, benzo(a)pyrene, benzo(b)fluoranthene, dibenz(a,h)anthracene, and indeno(1,2,3-cd)pyrene] in soil have FWCUGs for the cancer endpoint. An SOR was calculated for these potential carcinogens for the EPCs in the Surface Area EU as well as for each ISM sample. All calculated SORs are <1; therefore, no additional COCs were identified.

No surface water COCs were identified for the Trespasser scenario. Two inorganic chemicals (arsenic and cobalt) were identified as COPCs for this medium. The EPC for arsenic (0.00685 mg/L) is less than the lowest FWCUG (0.0415 mg/L). No FWCUG is available for cobalt in surface water; however, the MDC (0.0015 mg/L) does not exceed the USEPA residential RSL (0.011 mg/L) for drinking water at an HQ of 1.0. Thus, cobalt is not a COC.

4.3.4.5 <u>Identification of Dump Along Paris-Windham Road Chemicals of Concern: Resident</u> Farmer Scenario

The Resident Farmer is assumed to contact shallow surface soil and surface water. Exposure to subsurface soil is not included because the foundation of a house would have to be located outside the AOC. COC screening for the Resident Farmer is detailed in Appendix Tables B-10 (Surface Area EU), B-11 (AOC-wide ISM soil sample), and B-13 (surface water) and summarized below.

Soil COCs for the Resident Farmer scenario are summarized below and in Table 4-6.

• Benzo(a)pyrene was identified as a COC for the Resident Farmer scenario in the Surface Area EU. The EPC (0.33) exceeds the FWCUG for the Resident Farmer Adult (0.221 mg/kg).

 • Benzo(a)pyrene and dibenz(a,h)anthracene were identified as COCs in ISM sample PWss-CONT2 collected from across all 10 soil sample grids. Approximately one-half of this area was covered with at least 2 ft of clean fill after this sample was collected. The detected concentrations of benzo(a)pyrene and dibenz(a,h)anthracene were 1.4 and 0.36 mg/kg, respectively. The FWCUG for the Resident Farmer Adult is 0.221 mg/kg for both of these chemicals.

No additional COCs were identified based on the SOR analysis as summarized below.

• None of the COPCs identified in soil have similar toxic endpoints; therefore, no non-cancer SOR was calculated.

• Five COPCs [benz(a)anthracene, benzo(a)pyrene, benzo(b)fluoranthene, dibenz(a,h)anthracene, and indeno(1,2,3-cd)pyrene] in soil have FWCUGs for the cancer endpoint. An SOR was calculated for these potential carcinogens for the EPCs in the Surface Area EU as well as for the AOC-wide ISM sample. All calculated SORs are ≤1; therefore, no additional COCs were identified.

No surface water COCs were identified for the Resident Farmer. Two inorganic chemicals (arsenic and cobalt) were identified as COPCs for this medium. The EPC for arsenic (0.00685 mg/L) is less than the lowest FWCUG (0.011 mg/L). No FWCUG is available for cobalt in surface water; however, the MDC (0.0015 mg/L) is less than the USEPA residential RSL (0.011 mg/L) for drinking water at an HQ of 1.0. Thus, cobalt is not a COC.

4.3.5 Uncertainty Assessment

The sources of uncertainty, as well as the potential bias they impart to the risk assessment (i.e., whether conservatism is increased or decreased), are briefly discussed below.

4.3.5.1 Uncertainty in Estimating Potential Exposure

Sources of uncertainty in estimating potential human exposure include limitations of the sampling and analysis, comparison to background concentrations to identify SRCs, and estimation of EPCs.

Sampling Limitations – Uncertainties arise from limits on the media sampled, the total number and specific locations that can be sampled, and the parameters chosen for analysis to characterize the AOC. Sampling at the Dump Along Paris-Windham Road was targeted primarily at inorganic chemicals and asbestos. A subset of the total samples collected was analyzed for VOCs, SVOCs, pesticides, PCBs, explosives, and propellants. Soil has been characterized using both discrete and ISM sampling biased toward areas anticipated to have the highest level of potential contamination. Uncertainty is associated with exactly what sampled areas are currently covered by fill. A conservative estimate was made of the extent of excavation and fill. Some fill may extend onto areas included in the Surface Area EU, but its depth is assumed to be much less than that applied to the Fill Area EU.

Analytical Limitations – Uncertainty is associated with the contaminant concentrations detected and reported by the analytical laboratory. The quality of the analytical data used in the risk assessment was maximized and uncertainty minimized by implementing quality assurance/quality control (QA/QC) procedures that specify how samples are selected and handled; however, sampling errors, laboratory analysis errors, and data analysis errors can occur. Beyond the potential for errors, there is normal variability in analytical results. Some current analytical methods are limited in their ability to achieve detection limits at or below risk-based screening levels. Under these circumstances, it is uncertain whether the true concentration is above or below the screening levels that are protective of human health. When analytes have a mixture of detected and non-detected concentrations, EPC calculations may be affected by these detection limits. Risks may be overestimated as a result of some sample concentrations being reported as non-detected at the method detection limit (MDL) when the actual concentration may be much smaller than the MDL. Risks also may be underestimated if some analytes that were not detected in any sample were removed from the COPC list. If the concentrations of these analytes are below the MDL but above the screening level, the risk from these analytes would not be included in the risk assessment results.

Identification of SRCs – Uncertainty is associated with screening against background results from statistical limitations and natural variation in background concentrations. Because of this variation, metal concentrations below the background screening value are likely representative of background concentrations. Metal concentrations above the background screening level may be above background concentrations or may reflect natural variation. This is especially true for measured concentrations close to the background screening value.

EPCs – Soil was characterized using both discrete and ISM sampling techniques. ISM samples provide a physical average concentration across an exposure area. Use of ISM sampling reduces the uncertainty associated with estimating a statistical average exposure. Generally, the upper confidence limit on the arithmetic mean was adopted as the EPC for discrete sample results and was considered

to represent a conservative estimate of the average concentration. This imparts a small but intentional conservative bias to the risk assessment, provided the sampling captured the most highly contaminated areas. Representative EPCs for the EU were calculated from discrete data or measured with ISM data based on the assumption that samples collected from the EU were truly random samples. This assumption is not true for the Dump Along Paris-Windham Road. Sample locations were biased to identify the areas of highest contaminant concentrations. ISM sample PWss-CONT2 was collected from across the entire soil sampling grid. After this sample was collected, approximately one-half of the area sampled was covered with at least 2 ft of clean fill. Therefore, this sample is not representative of current surface soil conditions, but the area is also not completely covered by fill.

4.3.5.2 Uncertainty in Use of Facility-Wide Cleanup Goals

Sources of uncertainty in the FWCUGs used to identify COCs include the selection of appropriate receptor scenarios and exposure parameters, exposure models, and toxicity values used in the calculation of FWCUGs.

Selection of Representative Receptors – The OHARNG will control future use of the property and implement any LUCs that may be required as a component of RAs. As discussed in Section 4.3.3, the RAFLU for the AOC is Restricted Access, with the Range Maintenance Soldier as the representative receptor. The AOC location and topographic conditions preclude Residential Land Use. A low degree of uncertainty exists with respect to the future OHARNG-controlled land use and the assumption that RVAAP will not be released for Residential Land Use; however, a Resident Farmer receptor is included to provide a baseline assessment. Because this area is located immediately adjacent to a primary road, trespassers may visit the AOC; therefore, Adult and Juvenile Trespassers were also evaluated. The Dump Along Paris-Windham Road is located in the eastern-central area of Camp Ravenna (well distant from the property boundary), and it is unlikely a trespasser will visit this small area 75 days/year for 30 years. Therefore, some uncertainty exists as to the exact number of hours or days a trespasser may be present. The exposure assumptions for the Range Maintenance Soldier are also protective of the Adult and Child trespasser.

Exposure Parameters and Exposure Models – For each primary exposure pathway included in the FWCUGs, assumptions are made concerning the exposure parameters (e.g., amount of contaminated media a receptor can be exposed to and intake rates for different routes of exposure) and the routes of exposure. Most exposure parameters have been selected so that errors occur on the side of conservatism. When several of these upper-bound values are combined in estimating exposure for any one pathway, the resulting risks can be in excess of the 99th percentile and outside of the range that may be reasonably expected. Therefore, the consistent conservatism employed in the estimation of these parameters generally leads to overestimation of the potential risks.

Toxicity Values – The toxicity of chemicals is under constant study, and values change from time to time. The toxicity values used in the calculation of the FWCUGs were the most recent values

available at the time of those calculations (September 2008). These values are designed to be conservative and provide an upper-bound estimate of risk.

4.3.5.3 Uncertainty in the Identification of Chemicals of Concern

One of the two COPCs identified in surface water (cobalt) does not have FWCUGs. The MDC of cobalt (0.0015 mg/L) is less than the USEPA residential RSL (0.0011 mg/L) for drinking water at an HQ of 0.1. Thus, cobalt is not expected to contribute significantly to uncertainty in the results of the risk assessment.

4.3.6 Summary of Human Health Risk Assessment

This HHRA documents the COCs that may pose potential health risks to human receptors resulting from exposure to contamination at the Dump Along Paris-Windham Road. This HHRA was conducted as part of this SC/FFS and was based on the streamlined approach described in the FWCUG Report (USACE 2010a). The components of the risk assessment (receptors, exposure media, EPCs, and results) are summarized below.

Receptors – RVAAP is a controlled-access facility that is fenced and patrolled by security personnel. Full-time OHARNG, BRAC, and contractor staff work at the facility. Military training and operations are conducted at the facility. The AOC is not currently used for military training activities but may receive periodic foot traffic. The OHARNG projected future land use for the AOC is Restricted Access due to residual asbestos at the AOC. The representative receptor at the AOC is the Range Maintenance Soldier. This RAFLU (Restricted Access – Range Maintenance Soldier) forms the basis for identifying COCs. Because the AOC is located immediately adjacent to a primary road, trespassers may visit the AOC; therefore, Adult and Juvenile Trespassers were also evaluated. Topography (e.g., steep slopes and floodplain) and the presence of the covered dumpsite precludes unrestricted or Residential Land Use on the AOC. However, unrestricted or Residential Land Use could potentially occur adjacent to the AOC east of Paris-Windham Road. Therefore, an unrestricted scenario was evaluated in the HHRA as a comparative baseline, in accordance with CERCLA.

Exposure Media – Media of concern at the Dump Along Paris-Windham Road are surface soil, subsurface soil, and surface water. All soil samples were collected from 0-1 ft bgs. Some of these samples were subsequently covered with at least 2 ft of clean fill and now represent subsurface conditions.

Estimation of EPCs – For discrete soil and surface water, data EPCs were calculated from the results of all the discrete samples collected from each EU (listed in Tables 4-1 and 4-2). The EPC was either the UCL $_{95}$ or the MDC, whichever value was lowest. If the UCL $_{95}$ could not be determined, the EPC was the MDC. For ISM soil data, the EPC was the detected concentration in each ISM sample.

Results of HHRA – No COCs were identified in surface water for any receptor scenario. No COCs were identified in soil for the Range Maintenance Soldier or Adult and Juvenile Trespassers. Two

PAHs were identified as COCs in soil for the Resident Farmer. COCs and FWCUGs are summarized in Table 4-6.

3

Table 4-6. Summary of COCs and FWCUGs

		Chemicals	of Concern (FWCUG)			
Exposure Unit	Range Maintenance Soldier	Trespasser ^a	Resident Farmer ^b			
Soil						
Surface Area - Discrete Samples	None	None	Benzo(a)pyrene (0.221 mg/kg)			
Fill Area - Discrete Samples	None	NA	NA			
Fill Area ISM Sample (PWss-CONT1)	None	NA	NA			
AOC-Wide ISM Sample (PWss-CONT2)	None	None	Benzo(a)pyrene (0.221 mg/kg) Dibenz(a,h)anthracene (0.221 mg/kg)			
Surface Water						
Surface Water - Discrete Samples	None	None	None			

^aBoth Adult and Juvenile Trespasser scenarios were evaluated

AOC = Area of Concern

COC = Chemical of Concern

 $FWCUG = Facility\text{-}Wide\ Cleanup\ Goal$

ISM = Incremental Sampling Method

NA = Exposure medium not applicable to this receptor

5

4.4 ECOLOGICAL RISK ASSESSMENT

7 8

4.4.1 Introduction

9 10

11

12

13

14

15

16 17

18

19

20

The ERA presented in this SC/FFS follows a unified approach of methods integrating U.S. Army, Ohio EPA, and USEPA guidance. This ERA approach is consistent with the general approach by these agencies and primarily follows the Level I Scoping Level ERA, Level II Screening ERA, and Level III Baseline ERA outlined in the *Guidance for Conducting Ecological Risk Assessments* (Ohio EPA 2008), with specific application of components from the *RVAAP Facility-Wide Ecological Risk Work Plan* (USACE 2003c) (herein referred to as the FWERWP), *Risk Assessment Handbook Volume II: Environmental Evaluation* (USACE 2010b), and *Ecological Risk Assessment Guidance for Superfund: Process for Designing and Conducting Ecological Risk Assessments* (USEPA 1997). The process implemented in this SC/FFS combines these guidance documents to meet requirements of the Ohio EPA and U.S. Army, while following previously accepted methods established for RVAAP. This unified approach resulted from coordination between USACE and Ohio EPA during the summer of 2011.

212223

4.4.2 Scope and Objective

2425

2627

28

The Dump Along Paris-Windham Road contains habitat that supports ecological receptors. The habitat has known chemical contamination (USACE 2003c). Habitat types and an assessment of the ecological resources found at the Dump Along Paris-Windham Road are presented in subsequent sections. Additionally, the limited "RD/RA" confirmatory sample results (MKM 2004) are provided

^bBoth Resident Farmer Adult and Child scenarios were evaluated

to determine whether a qualitative ERA (Level I) is sufficient, based on the quality of the habitat and 2 the presence of contamination, or whether a more rigorous ERA (Level II or Level III) should be 3 conducted.

4 5

1

4.4.3 Level I: Scoping Level Ecological Risk Assessment

6 7

8

The ERA method for Level I follows guidance documents listed in Section 4.4.1. Level I is intended to evaluate if the AOC had past releases or the potential for current contamination, and if there are important ecological resources on or near the AOC.

9 10 11

The following two questions should be answered at the completion of the Level I ERA:

12

13 1. Are current or past releases suspected at the AOC? Current or past releases are determined by 14 evidence that chemical contaminants or chemicals of potential ecological concern (COPECs) are 15 present.

16 17

2. Are important ecological resources present at or in the locality of the AOC? Important ecological resources are defined in the Guidance for Conducting Ecological Risk Assessments (Ohio EPA 2008) and Technical Document for Ecological Risk Assessment: Process for Developing Management Goals (BTAG 2005).

20 21

18

19

22 If an AOC has contaminants but lacks important ecological resources, the ERA process can stop at 23 Level I. Contamination and important ecological resources must both be present to proceed to a 24 Level II Screening Level ERA.

25 26

4.4.3.1 **AOC Description and Land Use**

27 28

29

30

31

The Dump Along Paris-Windham Road is approximately 30 ft wide by 400 ft long or about 0.25 acres in size. There are two small wetlands at the AOC. The primary habitat is forest and is not large enough to completely support cover and food for small birds and mammals that typically require approximately 1 acre (USEPA 1993). The habitat area at the Dump Along Paris-Windham Road represents 0.001% of the 21,683 acres at RVAAP.

- 34 Activity on the AOC will consist of occasional foot traffic associated with minor maintenance 35 activities (e.g., mowing and vegetation control) and road maintenance (e.g., mowing along the road 36 berm and road surface repairs/patching). Activities could also include foot traffic by range control 37 (because the AOC is in the SDZ) and wildlife and natural resource management activities. The
- 38 representative receptor for the Dump Along Paris-Windham Road is the Range Maintenance Soldier.
- 39 The Adult and Juvenile Trespassers and Adult and Child Resident are also evaluated in the HHRA.
- 40 U.S. Army natural and ecological resource management activities may apply if habitat disturbance
- 41 occurs.

4.4.3.2 Evidence of Chemical Contamination

Previous investigative activities at the Dump Along Paris-Windham Road include an RRSE in 1998, environmental sampling conducted by USACE, Louisville District in 2001, confirmatory sampling performed during the April 2003 limited "RD/RA," and biological and water quality sampling conducted in June 2003 (USACE 2005a).

The RRSE summarized in the *Relative Risk Site Evaluation for Newly Added Sites* (USACHPPM 1998) identified potential ecological receptors for exposure to surface soil and sediment contamination and assumed complete exposure pathways due to the AOC's proximity to Sand Creek. As a result, the RRSE score for this AOC was "High." The "High" score was prior to the limited "RD/RA" in April 2003.

In 2003, USACE, Louisville District prepared a *Decision Document for a Removal Action at Paris-Windham Road Dumpsite (RVAAP-51)* (USACE 2003a). According to the DD, USACE, Louisville District collected soil samples in 2001 to further characterize the AOC. The DD reported the principal contaminants detected during the 2001 sampling with potential impact to ecological receptors were cadmium, PCBs, benz(a)anthracene, benzo(a)pyrene, benzo(b)fluoranthene, chrysene, and indeno(1,2,3-cd)pyrene.

The confirmatory sampling performed during the April 2003 limited "RD/RA" is summarized in Section 3.0 and the *Final Report for Remedial Design/Remedial Action Plan at Paris-Windham Road Dump* (MKM 2004). The limited "RD/RA" was conducted in accordance with CERCLA to mitigate risks related to potential contact with exposed waste material. The limited "RD/RA" removal activities consisted of removing all existing surface debris and limited removal of subsurface debris. Approximately 300 tons of surface and subsurface debris were removed from the AOC (see Section 3.1.3).

Ohio EPA and USACE, Louisville District investigated several streams at RVAAP using a network of various biological/water quality sampling stations (USACE 2005a). The purpose of this investigation was to document ecological effects of AOCs on stream or pond biota and conditions. The biological/water quality sampling was conducted between June and September 2003 after the limited "RD/RA" and the associated confirmatory sampling but prior to any site restoration conducted in November 2003. The site restoration consisted of returning the excavation area to grade using a combination of clean hard fill and soil backfill from an Ohio EPA-approved source. The area was then seeded and mulched (see Section 3.1.3).

The goal of this ERA is to identify COPECs for the Dump Along Paris-Windham Road using available analytical data from the 2003 limited "RD/RA" confirmatory sampling. The screening level approach to evaluate limited "RD/RA" confirmatory sample results followed instructions presented in the *Guidance for Conducting Ecological Risk Assessments* (Ohio EPA 2003) and consisted of the first two of six steps listed in Figure III of the FWERWP (USACE 2003c). These two steps identify the evaluation procedures, which were used to determine AOC-related COPECs. Section 3.2 of this

SC/FFS details chemical concentration data. The limited "RD/RA" confirmatory sampling included collection of discrete surface soil (0-1 ft bgs) samples and ISM surface soil (0-1 ft bgs) samples. Discrete soil samples are used in the COPEC screening. In addition, one ISM soil sample from Grid 9 that was analyzed for SVOCs was used in the COPEC screening. These samples were collected from 0-1 ft bgs after the removal action and prior to the placement of the soil and vegetation cover.

5 6 7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

1

2

3

4

The 2003 limited "RD/RA" did not differentiate wet or dry sediment. Therefore, as part of this SC/FFS, SAIC scientists conducted a field survey to determine the sediment type (wet or dry per RVAAP guidelines as explained in Section 1.1). Surface water occurs in the drainage swale on only an intermittent basis. During an August 2009 walkover/assessment, SAIC scientists noted the sediment in the drainage swale had high moisture content, but no standing water was observed. By contrast, SAIC scientists did observe standing water in the drainage swale in November 2011, following a rainfall event (see photographs in Appendix A). Based on the conditions of the AOC, sediment in the drainage swale is considered dry sediment because of the intermittent surface water. It is not considered permanent habitat for aquatic organisms. Therefore, dry sediment (0-1 ft bgs) is addressed as surface soil in terms of contaminant nature and extent, fate and transport, and risk exposure models. This approach is consistent with the FWCUG Report. For surface water, discrete samples collected during the limited "RD/RA" were used to evaluate the drainage swale (i.e., former stream channel) located along the western border of the AOC. Duplicate samples were used in this assessment. This ERA uses updated ecological screening values (ESVs) that follow the revised Guidance for Conducting Ecological Risk Assessments (Ohio EPA 2008), as provided in Appendix Tables C-1 and C-2.

222324

25

2627

28

29

30

31

32

33

34

In the first step of the COPEC screening process, the MDC of each chemical is compared to its respective facility-wide background concentration. Chemicals are not considered site-related if the MDC is below the background concentration. For all chemicals detected above background concentrations, the MDC is compared to the chemical-specific ESV. The hierarchy of screening values was based on the guidance included in the FWERWP and *Guidance for Conducting Ecological Risk Assessments* (Ohio EPA 2008). In addition to the ESV comparison, it was determined if the chemical is a persistent, bioaccumulative, and toxic (PBT) compound. Chemicals are retained as COPECs if they exceed background concentrations and the ESV, if the chemical exceeds background concentrations and had no toxicity information, or if the chemical is considered a PBT compound. Ratios of MDC to ESV are used to determine the COPECs that result from the limited "RD/RA" dataset. A ratio greater than 1 suggests a possible environmental consequence. Any chemicals with ratios greater than 1 are identified as COPECs.

353637

Groundwater was not included in the ERA. As explained in Section 3.2.2 of the FWERWP, groundwater is not considered an exposure medium to ecological receptors.

38 39

The ERA tables for soil and surface water are included in Appendix Tables C-3, C-4, and C-5 and contain the following:

42 43

- Frequency of detection;
- MDC;

- 1 Average results;
- Background concentrations;
- SRC determination;
- ESVs used for COPEC determinations;
- 5 Ratio of MDC to ESV:
- PBT compound identification;
- 7 COPEC determination; and
- 8 COPEC rationale.

9 10

11 12

13 14

15

16

17

18

19

20

As discussed in Section 4.2.1, the soil within the geographic area of the Dump Along Paris-Windham Road was subdivided into three spatial aggregates: the Fill Area EU, the Surface Area EU (Figure 4-3), and the AOC-Wide EU. The Fill Area EU is located in the middle of the dump area [characterized by discrete sample locations PWss-005, PWss-006, PWss-007, PWss-008, and PWss-009 and ISM sample location PWss-CONT1 (associated with Grid 9)] and was excavated and covered with approximately 2 ft of clean fill. The Surface Area EU is in the northern and southern ends of the dump area [characterized by discrete sample locations PWss-001, PWss-002, PWss-003, PWss-004 (including a duplicate sample), and PWss-010] and the drainage swale [characterized by samples PWsd-001, PWsd-002, PWsd-003 (including a duplicate sample), PWsd-004, PWsd-005, and PWsd-006] and is located outside the excavation area. The AOC-Wide EU was not evaluated in the ERA because it consisted of only a single ISM sample. Rather, the ERA focused on the other two soil EUs that had discrete samples.

212223

24

25

26

Intermittent surface water at the Dump Along Paris-Windham Road is limited to a long, narrow drainage swale downslope of the excavated dump area. The Surface Water EU includes all samples collected in the drainage swale (i.e., former stream channel) located along the western border of the AOC. These surface water samples were co-located with sediment samples that were later classified as soil and included in the Surface Area EU.




Figure 4-3. Exposure Units at the Dump Along Paris-Windham Road

COPECs for Soil at the Fill Area EU. Thirty-two chemicals were detected in surface soil at the Fill Area EU. Five chemicals (calcium, iron, magnesium, potassium, and sodium) were essential nutrients and were excluded as SRCs, as described in Section 3.2.1. Five inorganic chemicals and 14 organic chemicals were determined to be SRCs because they exceeded background concentrations, or they did not have an associated background concentration for comparison. Of the 19 SRCs, two inorganic chemicals (mercury and zinc) exceeded the ESVs and are identified as COPECs (Table 4-7). Mercury is also a PBT compound. Though it did not exceed its ESV, PCB-1254 was also identified as a COPEC because it is a PBT compound. The calculated ratio of MDC to ESV is shown in Table 4-7 for each COPEC. Appendix Table C-3 presents the details of the ESV comparisons for surface soil at the Fill Area EU.

10 11 12

1

2

3

4

5

7

8

9

Table 4-7. Summary of COPECs for Surface Soil at the Fill Area EU

COPEC	MDC (mg/kg)	ESV (mg/kg)	Ratio of MDC to ESV	Comments
Mercury	0.048	0.00051	94.1	Highest ratio at 94x; PBT compound
Zinc	100	46	2.2	None
PCB-1254	0.23	0.371	0.62	PBT compound

Table excludes nutrients

-- = not applicable, no ESV is available for comparison

15 COPEC = Chemical of Potential Ecological Concern 16

ESV = Ecological Screening Value

EU = Exposure Unit

MDC = Maximum Detected Concentration

19 PBT = Persistent, Bioaccumulative, and Toxic 20

PCB = Polychlorinated Biphenyl

x = multiplier

21 22 23

24

25

26

27

28

29

30

31

32

33

13

14

17

18

COPECs for Soil at the Surface Area EU. Thirty chemicals were detected in surface soil at the Surface Area EU. Five chemicals (calcium, iron, magnesium, potassium, and sodium) were essential nutrients and were excluded as SRCs, as described in Section 3.2.1. Eleven inorganic chemicals and 10 organic chemicals were determined to be SRCs because they either exceeded background concentrations or they did not have an associated background concentration for comparison. Of the 21 SRCs, six inorganic chemicals (aluminum, cadmium, lead, manganese, mercury, and zinc) exceeded the ESVs and are identified as COPECs (Table 4-8). Mercury is also a PBT compound. Though it did not exceed its ESV, PCB-1254 was also identified as a COPEC because it is a PBT compound. One organic chemical (nitrocellulose) was selected as a COPEC because it did not have an ESV. The calculated ratio of MDC to ESV is shown in Table 4-8 for each COPEC. Appendix Table C-4 presents the details of the ESV comparisons for surface soil at the Surface Area EU.

COPEC	MDC (mg/kg)	ESV (mg/kg)	Ratio of MDC to ESV	Comments
Aluminum	18,000	50	360	Highest ratio at 360x
Cadmium	0.59	0.36	1.6	None
Lead	29	11	2.6	None
Manganese	1,900	220	8.6	None
Mercury	0.08	0.00051	157	Second highest ratio at 160x; PBT compound
Zinc	120	46	2.6	None
Nitrocellulose	2	No ESV		None
PCB-1254	0.09	0.371	0.23	PBT compound

Table excludes nutrients

-- = not applicable, no ESV is available for comparison

COPEC = Chemical of Potential Ecological Concern

ESV = Ecological Screening Value

EU = Exposure Unit

MDC = Maximum Detected Concentration

PBT = Persistent, Bioaccumulative, and Toxic

PCB = Polychlorinated Biphenyl

x = multiplier

10 11 12

23456789

COPECs for Sediment. Sediment in the drainage swale is considered dry sediment because of the intermittent nature of the surface water. It is not considered permanent habitat for aquatic organisms. Therefore, dry sediment (0-1 ft bgs) is addressed as surface soil in the Surface Water EU.

141516

17

18

19

20

21

22

23

24

13

COPECs for Surface Water. Seventeen chemicals were detected in surface water. Five chemicals (calcium, iron, magnesium, potassium, and sodium) were essential nutrients and were excluded as SRCs, as described in Section 3.2.1. Seven detected inorganic chemicals, and one organic chemical were determined to be SRCs because they either exceeded background concentrations or they did not have an associated background concentration for comparison. Of the nine SRCs, one inorganic chemical (manganese) exceeded its ESV and is identified as a COPEC (Table 4-9). In addition, nitrocellulose was selected as a COPEC because it does not have an ESV for comparison. Mercury did not exceed its ESV in surface water but is retained as a COPEC because it is a PBT compound. The calculated ratio of MDC to ESV is shown in Table 4-9 for each COPEC. Appendix Table C-5 presents the details of the ESV comparisons for surface water.

252627

28

29

30

31

32

Summary of ERA. The ERA was performed using the limited "RD/RA" confirmatory sampling results to determine COPECs at the Dump Along Paris-Windham Road in surface soil and surface water. There are three surface soil COPECs identified in the ERA for the Fill Area EU: mercury, zinc, and PCB-1254. There are eight surface soil COPECs identified in the ERA for the Surface Area EU: aluminum, cadmium, lead, manganese, mercury, zinc, nitrocellulose, and PCB-1254. There are three surface water COPECs identified in the ERA: manganese, mercury, and nitrocellulose. Based on the identification of COPECs, ecological risk in surface soil and surface water was predicted.

Table 4-9. Summary of COPECs for Surface Water

COPEC	MDC (mg/kg)	ESV (mg/kg)	Ratio of MDC to ESV	Comments
Manganese	0.56	0.12	4.7	Highest ratio at 5x
Mercury	0.0001	0.0017	0.06	PBT compound
Nitrocellulose	0.094	No ESV		None

Table excludes nutrients

 ${ ext{--}}=$ not applicable, no ESV is available for comparison

COPEC = Chemical of Potential Ecological Concern

ESV = Ecological Screening Value

MDC = Maximum Detected Concentration

PBT = Persistent, Bioaccumulative, and Toxic

x = multiplier

4.4.3.3 <u>Ecological Significance</u>

4 5

Sources of data and information about the ecological resources at the Dump Along Paris-Windham Road include the *Integrated Natural Resource Management Plan* (INRMP) (OHARNG 2008a), *Facility-Wide Biological and Water Quality Study* (USACE 2005a), previous characterization work (USACHPPM 1998, USACE 2003a, and MKM 2004), and visits to the AOC conducted for the SC/FFS.

One of the two key questions to answer in the Level I Scoping Level ERA is whether there are ecologically important and especially ecologically significant resources at the Dump Along Paris-Windham Road. Ecological importance is defined as a place or resource that exhibits unique, special, or other attributes that makes it of great value. Ecological significance is defined as an important resource found at an AOC or in its vicinity that is subject to contaminant exposure. The underlying basis for this distinction can be found in *Ecological Significance and Selection of Candidate Assessment Endpoints* (USEPA 1996), stated as follows:

"A critical element in the ERA process requires distinguishing important environmental responses to chemical releases from those that are inconsequential to the ecosystem in which the site resides: in other words, determining the ecological significance of past, current, or projected site-related effects."

Important places and resources identified by the U.S. Army and Ohio EPA (Appendix Table C-6) include wetlands, terrestrial areas used for breeding by large or dense aggregations of animals, habitat known to be used by threatened or endangered species, state land designated for wildlife or game management, locally important ecological places, and state parks. Both the U.S. Army and Ohio EPA recognize 17 important places and resources. The U.S. Army recognizes an additional 16 important places (BTAG 2005), and the Ohio EPA recognizes another 6 important places (Ohio EPA 2008). In total, there are 39 important places. Presence or absence of an ecologically important place can be determined by comparing environmental facts and characteristics of the Dump Along Paris-Windham Road with each of the important places and resources listed in Appendix Table C-6.

Ecological significance is defined as an important resource found at an AOC or in its vicinity that is subject to contaminant exposure. Thus, important places and resources listed in Appendix Table C-6 are elevated to ecologically significant when present on the AOC and there is exposure to contaminants. For all 39 important places and resources, it is relatively clear the ecological place or resource is present or absent on the AOC; therefore, the decision process is objective. If no important or significant resource is present at an AOC, the evaluation will not proceed to Level II regardless of the presence of contamination. Instead, the Level I Scoping Level ERA would acknowledge there are important ecological places but that those resources are not ecologically significant, and no further evaluation is required.

Management Goals for the AOC. Regardless of whether the evaluation is concluded at Level I or continues to Level II, there is another level of environmental protection for the Dump Along Paris-Windham Road through the natural resource management goals expressed in the INRMP (OHARNG 2008a). The U.S. Army is required to monitor ecological conditions to maintain or enhance the facility's natural resources and ecosystem. While the monitoring focuses on the potential adverse effects from training activities, degradation from contamination would be noticed as well.

Some Natural Resources Management Goals of OHARNG (listed in Appendix Table C-7) benefit the Dump Along Paris-Windham Road. For example, Goal 1 states natural resources need to be managed in a compatible way with the military mission, and Goal 5 requires the U.S. Army to sustain usable training lands and native natural resources by implementing a natural resource management plan which incorporates invasive species management and by utilizing native species mixes for revegetation after ground disturbance activities. These management goals help detect degradation (whether from training activities or historical contamination). While the applicability of the remaining nine management goals to the Dump Along Paris-Windham Road varies, all of the management goals are intended to monitor, maintain, or enhance the RVAAP natural resources and ecosystem. While these goals are for the management of all types of resources at and near the Dump Along Paris-Windham Road, they do not affect the decisions concerning the presence or absence of important or significant ecological places or resources there.

Important Places and Resources. Ecological importance means a place or resource that exhibits a unique, special, or other attribute that makes it of great value. Examples of important places and resources include wetlands, terrestrial areas used for breeding by large or dense aggregations of animals, and habitat of state-listed or federally-listed species. An important resource becomes significant when found on an AOC and there is contaminant exposure.

As noted in Appendix Table C-6, a small portion (0.04 acres) of wetlands is within the AOC. The wetland is an important ecological resource at the AOC. The wetlands are discussed in greater detail later in this section.

Terrestrial Resources. The Dump Along Paris-Windham Road is dominated by terrestrial resources.

Habitat Descriptions and Species. The INRMP and AOC visits by SAIC scientists indicated the habitat in the immediate vicinity of the Dump Along Paris-Windham Road contains two types of vegetation (Figure 4-4). The dominant vegetation is a temporarily flooded forest alliance of green ash (Fraxinus pennsylvanica), American elm (Ulmus americana), and hackberry (Celtis occidentalis and laevigata) (Photographs 4-1 and 4-2). One other forest alliance consisting of American beech (Fagus grandifolia), oak (Quercus Spp.), and maple (Acer spp.) is found along the southeast border of the AOC. This characterization was originally established by a vegetation study using aerial photography and field verification (USACE 1999) and was later used in the INRMP (OHARNG 2008a).

During a field survey conducted at the Dump Along Paris-Windham Road in August 2009, SAIC scientists confirmed the main habitat type: green ash (*Fraxinus pennsylvanica*)/American elm (*Ulmus americana*)/hackberry (*Celtis occidentalis* and *laevigata*) temporarily flooded forest alliance.

The green ash (*Fraxinus pennsylvanica*)/American elm (*Ulmus americana*)/hackberry (*Celtis occidentalis* and *laevigata*) temporarily flooded forest alliance and the American beech (*Fagus grandifolia*)/oak (*Quercus* Spp.)/maple (*Acer* spp.) forest alliance includes small open areas and understory that results in multi-story vegetation (Photographs 4-1 and 4-2), providing layers of vegetation for various foraging height preferences of birds, mammals, insects, and other organisms.

Based on August 2009 and November 2011 observations (Photographs 4-1 and 4-2 and Appendix A), SAIC scientists assessed the habitat at the Dump Along Paris-Windham Road to be healthy and functioning. Functional habitat was determined by noting the absence of large bare spots and dead vegetation or other obvious visual signs of an unhealthy ecosystem. Some vegetation was removed during the limited "RD/RA;" however, the AOC walkover conducted by SAIC scientists in August 2009 and November 2011 showed vegetative recovery has occurred since the limited "RD/RA" in 2003. Appendix A provides photographs of current conditions at the AOC and the state of vegetative recovery observed during the AOC walkovers.

Threatened and Endangered and Other State-listed or Federally-listed Species. There are currently no federally-listed species or critical habitat on Camp Ravenna. The Dump Along Paris-Windham Road has not been previously surveyed for rare species; however, there have been no documented sightings of state-listed, federally-listed, threatened, or endangered species at the AOC (OHARNG 2008a).

Other Terrestrial Resources. While there are no other known important terrestrial places and resources (Appendix Table C-6), there are other resources at or near the Dump Along Paris-Windham Road (e.g., vegetation, animals) that interact in their ecosystems and support nutrient cycling and energy flow. For example, wildlife such as wild turkey (*Meleagris gallopavo*) and white-tailed deer (*Odocoileus virginianus*) could use the area. Also, it is possible that burrowing animals could be exposed to soil at depths greater than 1 ft. The INRMP provides information about species and habitat surveys at RVAAP (e.g., timber and ecological succession) (OHARNG 2008a). There are no other reported surveys of habitats and wildlife at the Dump Along Paris-Windham Road beyond those summarized in the INRMP (OHARNG 2008a).




Figure 4-4. Natural Resources (OHARNG 2008) Inside and Near Habitat Area at the Dump Along Paris-Windham Road



Photograph 4-1. Looking South along Paris-Windham Road; Green Ash, American Elm, and Hackberry Temporary Flooded Forest Alliance in Background (August 2009)



Photograph 4-2. Drainage Swale with no Standing Water Along the Northwest Portion of the AOC; Green Ash, American Elm, and Hackberry Temporary Flooded Forest Alliance in Background (August 2009)

Aquatic Resources. The Dump Along Paris-Windham Road has one type of aquatic resource, as described below.

Habitat Descriptions and Species. Wetlands exist in the small area in the northern portion of the AOC and within the drainage swale immediately adjacent to the southern portion of the AOC (Figure 4-4).

Wetlands. Wetlands are important habitats with water-saturated soil or sediment whose plant life can survive saturation. Wetlands are home to many different species and are also chemical sinks that can serve as detoxifiers and natural water purifiers. It is expected that the wetlands/drainage swale at the Dump Along Paris-Windham Road (Figure 4-4) perform these and other related functions.

In November 2011, an SAIC Professional Wetland Scientist used the Ohio Rapid Assessment Method (ORAM) (Ohio EPA 2001) to assess the condition and ecological importance of the wetlands. The wetlands are located on the western side of the AOC, with one in the northern portion of the AOC and the other in the southern portion of the AOC. The wetlands at the Dump Along Paris-Windham Road consist of two small wetlands (designated northern and southern wetlands) and were evaluated together because they are within 140 ft of each other and part of the same landform. Using the ORAM, wetlands are classified into three categories:

• Category 1 wetlands are described as "limited quality waters." They are considered to be a resource that has been degraded, has limited potential for restoration, or is of such low functionality that lower standards for avoidance, minimization, and mitigation can be applied. Scores range from 1 to 29.

• Category 2 includes wetlands of moderate quality and wetlands that are degraded but exhibit reasonable potential for restoration. Scores range from 30 to 59.

• Category 3 includes wetlands of very high quality and wetlands of concern regionally and/or statewide, such as wetlands that provide habitat for species listed as threatened or endangered. Scores range from 60 to 100.

The field sheet detailing the ORAM is presented in Appendix Figure C-1. Figure 4-4 shows the location of the evaluated wetlands with jurisdictional and planning level survey wetlands [i.e., based on desktop surveys conducted for the OHARNG of wetlands data and resources (i.e., NWI maps, aerials)] within the vicinity. Based on the ORAM, the wetlands at the Dump Along Paris-Windham Road are classified as Category 2 (with a score of 37), indicating a moderate wetland quality with some degradation of wetland functions and conditions (Appendix Figure C-1).

Dominant vegetation near the wetlands is forest habitat that has developed since the modifications to Sand Creek and the dumping at the AOC. The northern wetland covers 0.03 acres and lies almost entirely within the AOC. The southern wetland covers 0.18 acres, with 0.01 acres of the wetland inside the AOC. The combined area of the northern and southern wetlands is 0.21 acres. The two

small wetlands associated with the Dump Along Paris-Windham Road appear to be relic floodplain features. The wetlands may be former overflow channels on the Sand Creek floodplain, or they may represent an original channel of Sand Creek prior to dredging and channelization by the U.S. Army. Both wetlands have been mostly disconnected from Sand Creek by a large berm on the right bank that was apparently created during channelization of the creek. Wetland hydrology is largely dependent on precipitation, with secondary inputs from high flows in Sand Creek.

Because there is contamination within the Dump Along Paris-Windham Road, further contaminant analysis was conducted to determine if the contamination is at a level of concern to ecological receptors in the wetlands.

Eight COPECs [six inorganic chemicals (aluminum, cadmium, lead, manganese, mercury, and zinc) and two organic chemicals (nitrocellulose and PCB-1254)] were identified at the Surface Area EU (Table 4-8), which included dry sediment samples in the wetland/drainage swale area (Section 4.4.3.2). Three COPECs (manganese, mercury, and nitrocellulose) were identified in surface water (Table 4-9). To determine if the dry sediment and surface water COPECs were impacting the wetlands, the concentrations of COPECs in the seven dry sediment and surface water samples (PWsd/sw-001, PWsd/sw-002, PWsd/sw-003, PWsd/sw-003(duplicate), PWsd/sw-004, PWsd/sw-005, and PWsd/sw-006) were assessed (Table 4-10).

The results are as follows:

• Manganese was detected below its ESV in the seven dry sediment samples. Although concentrations of manganese exceeded its ESV in all seven surface water samples collected in or around the wetlands, the ESV was below the RVAAP background concentration. Therefore, this ESV is judged to be conservative. When the average concentration of manganese (0.38 mg/L) from the seven surface water samples is compared to the RVAAP background concentration (0.391 mg/L), the average concentration of manganese is less than its background concentration. This suggests manganese is not present in dry sediment and surface water at concentrations of concern for ecological receptors in the wetlands.

• Although the maximum concentrations of aluminum (9,900 mg/kg) and lead (25 mg/kg) in dry sediment exceeded their ESVs (50 mg/kg and 11 mg/kg, respectively), aluminum and lead were not detected above their background concentrations (17,700 mg/kg and 26.1 mg/kg, respectively) in any of the dry sediment samples. As a result, these inorganic chemicals are not present at concentrations of concern for ecological receptors in the wetlands.

• Although cadmium (0.59 mg/kg) exceeded its ESV (0.36 mg/kg) in one of the seven dry sediment samples (PWsd-002), this concentration is slightly above its respective ESV. Cadmium was not detected in the remaining six dry sediment samples. In surface soil and surface water samples, cadmium is not detected above its ESV. As a result, cadmium is not present at concentrations of concern for ecological receptors in the wetlands.

- Although concentrations of mercury and zinc exceeded their ESVs in all seven dry sediment samples, the ESVs were below the RVAAP background concentrations. Therefore, these ESVs are judged to be conservative. When the average concentrations of mercury (0.066 mg/kg) and zinc (92 mg/kg) from the seven dry sediment samples are compared to the RVAAP background concentrations (0.036 mg/kg and 61.8 mg/kg, respectively), the average concentrations of these samples are similar to their background concentrations. This suggests these inorganic chemicals are not present at concentrations of concern for ecological receptors in the wetlands. Mercury is a COPEC for surface water because it is a PBT compound; however, is not likely a concern for ecological receptors in the wetlands because concentrations do not exceed the ESV, and bioaccumulation in higher trophic levels is assumed to be considered in development of the ESV per Ohio Administrative Code (OAC) 3745-1-37.
- Nitrocellulose in dry sediment and surface water is essentially non-toxic to wildlife (USEPA 1987) and is not a concern for ecological receptors.
 - PCB-1254 was identified as a COPEC because it is a PBT compound. PCB-1254 was analyzed in only one dry sediment sample (PWsd-004) in the southern wetland and was detected at a concentration (0.086 mg/kg) below the ESV for total PCBs (0.371 mg/kg). PCB-1254 was analyzed in one surface soil sample (PWss-009) and was also detected at a concentration (0.23 mg/kg) below the ESV for total PCBs (0.371 mg/kg). While PCB-1254 is a PBT compound, it is not likely a concern for ecological receptors in the wetlands because concentrations do not exceed the ESV and bioaccumulation in higher trophic levels is considered in development of the ESV (DOE 1997).

In summary, although contamination is present in dry sediment and surface water samples, review of the data suggests that any migration of contamination from the AOC to the wetlands/drainage swale along the western boundary has not resulted in concentrations of concern to ecological receptors. As a result, although the wetlands are an important place, they are not ecologically significant with respect to the contamination at the Surface Area EU.

Table 4-10. Summary of COPEC Concentrations for Dry Sediment and Surface Water at and in the Vicinity of the Dump Along Paris-Windham Road

					Sampling Stations					
COPEC	Units	Background Concentration	ESV	PWsd/sw-001	PWsd/sw-002	PWsd/sw-003	PWsd/sw-003 (Duplicate)	PWsd/sw-004	PWsd/sw-005	PWsd/sw-006
					Dry Sediment	t				
Aluminum	mg/kg	17,700	50	8,000	9,000	7,100	7,400	8,400	9,900	7,600
Cadmium	mg/kg	0	0.36	ND	0.59	ND	ND	ND	ND	ND
Lead	mg/kg	26.1	11	19	25	19	18	18	16	20
Manganese	mg/kg	1,450	220	99	150	97	95	120	100	150
Mercury	mg/kg	0.036	0.00051	0.059	0.08	0.058	0.069	0.073	0.077	0.05
Zinc	mg/kg	61.8	46	81	120	75	73	88	99	90
Nitrocellulose	mg/kg	No BKG	No ESV	NR	NR	NR	NR	2	NR	NR
PCB-1254	mg/kg	No BKG	No ESV	NR	NR	NR	NR	0.086	NR	NR
	•	•		-	Surface Water	r	•			
Manganese	mg/L	0.391	0.12	0.32	0.27	0.26	0.26	0.51	0.47	0.56
Mercury	mg/L	No BKG	0.0017	0.0007	0.00009	0.00009	0.000091	ND	0.0001	0.00008
Nitrocellulose	mg/L	No BKG	No ESV	NR	NR	NR	NR	0.094	NR	NR

Background concentrations for surface soil (0-1 ft bgs) and surface water is from the final facility-wide background concentrations for RVAAP, published in the *Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio* (USACE 2001b)

COPEC = Chemical of Potential Ecological Concern

ESV = Ecological Screening Value

ND = Not Detected

No BKG = A background concentration does not exist for the specified chemical

NR = Not reported

PCB = Polychlorinated Biphenyl

Bold = Concentration exceeds the background concentration and the ESV

1 Threatened and Endangered and Other State-listed or Federally-listed Species. There are currently no federally listed species or critical habitat on Camp Ravenna. The Dump Along Paris-Windham

Road has not been previously surveyed for rare species; however, there have been no documented

4 sightings of rare, threatened or endangered species at the AOC (OHARNG 2008a).

Other Aquatic Resources. There are no other known aquatic resources (Appendix Table C-6) at or near the Dump Along Paris-Windham Road (e.g., vegetation, animals). There are no other reported surveys of habitats and wildlife at the Dump Along Paris-Windham Road beyond those summarized in the INRMP (OHARNG 2008a). There are two nearby biological and water quality stations. The following subsections provide a summary of the biological and water quality stations in the vicinity of the Dump Along Paris-Windham Road.

Biological/Water Quality Sampling Stations. Ohio EPA and USACE, Louisville District investigated several streams at RVAAP in a network of various biological/water quality sampling stations (USACE 2005a). The purpose of this investigation was to document ecological effects of AOCs on stream or pond biota and conditions. Two sampling stations were located in the vicinity of the Dump Along Paris-Windham Road. Station S-7 was located upstream of the AOC, and station S-9 was located downstream of the AOC. The upstream biological/water quality station (S-7) provides information regarding potential contamination from upstream AOCs and if upstream AOCs may be contributing to adverse biological, chemical, and physical measurements in the vicinity of the Dump Along Paris-Windham Road. The downstream sampling station (S-9) provides information about potential contamination from the Dump Along Paris-Windham Road and upstream AOCs. If the downstream sampling station has a positive rating (e.g., good, excellent, full attainment, and other positive terms reported in the study), it means that the Dump Along Paris-Windham Road and other upstream AOCs are not adversely impacting the quality of Sand Creek.

According to the *Facility-Wide Biological and Water Quality Study* (USACE 2005a), each sampling location included sediment sampling/assessment, surface water sampling/assessment, fish and macroinvertebrate community assessment, and habitat assessment. The sampling reach for stream sampling stations ranged 120-210 meters.

Sediment evaluations were conducted in June 2003 using guidelines established in *Development and Evaluation of Consensus-based Sediment Quality Guidelines for Freshwater Ecosystems* (MacDonald et al. 2000), sediment reference values (SRVs) for inorganic chemicals (Ohio EPA 2003), and USEPA Region 5 Ecological Screening Levels (ESLs) (USEPA 2003). Sediment samples were analyzed for SVOCs, pesticides, PCBs, TAL metals, explosives, percent solids, cyanide, ammonia, nitrate, and phosphorus. Surface water grab samples collected in June and September 2003 were evaluated using Ohio Water Quality Standards (WQS) criteria, reference conditions, or the *Facility-wide Biological and Water Quality Study*. Surface water samples were analyzed for TAL metals, pesticides, PCBs, explosives, SVOCs, and several nutrients.

Fish and macroinvertebrate sampling and assessments occurred in August and September 2003. Fish were sampled using electrofishing methods. Macroinvertebrate communities were assessed using

artificial substrates (quantitative sampling), supplemented with a composite natural substrate (qualitative sampling). Both the fish and macroinvertebrate community assessments followed the methods in the *Biological Criteria for the Protection of Aquatic Life: Volume III, Standardized Biological Field Sampling and Laboratory Methods for Assessing Fish and Macroinvertebrate Communities* (Ohio EPA 1989).

The physical habitat assessment was conducted in June 2003 and used the Qualitative Habitat Evaluation Index (QHEI) developed by the Ohio EPA (Rankin 1989, 1995). The types(s) and quality of substrates, amount and quality of instream cover, channel morphology, extent and quality of riparian vegetation, pool, run, riffle development and quality, and gradient are some of the habitat characteristics used to determine the QHEI score.

Sampling Station Locations. Station S-7 (RM 2.4) is located on Sand Creek approximately 1,900 ft upstream from the Dump Along Paris-Windham Road and provides the closest upstream reference point for the AOC. Station S-9 (RM 1.9) is located on Sand Creek at Paris-Windham Road. The station is immediately downstream from the Dump Along Paris-Windham Road and provides the closest downstream sampling point to the AOC.

Summary of Sampling/Assessment Results. Table 4-11 shows the ratings of the attributes for sampling stations S-7 and S-9. Review of the Facility-Wide Biological and Water Quality Study (USACE 2005a) data from the two stations showed many positive attribute ratings (e.g., good, excellent, full attainment) and little to no sign of aquatic impairment. Each station was rated at Full Use Attainment Status, which indicated all indices met the Ohio EPA biological criteria.

At S-7, all inorganic chemicals tested in sediment were below Ohio SRVs and threshold effects concentration (TEC) levels. All tested explosives, pesticides, and PCBs were not detected in sediment samples collected from S-7. The few SVOCs were measured at low levels, with all concentrations below TEC or ESL guidelines. Ammonia and total phosphorus levels were measured below screening guidelines. None of the surface water chemical concentrations at S-7 exceeded Ohio WQS aquatic life maximum or average water quality criteria, and none of the chemicals measured exceeded criteria protective of the Warm Water Habitat (WWH) aquatic life use (USACE 2005a). Overall, the sediment quality and water quality at S-7 was rated "excellent." The fish community at S-7 was rated "marginally good." The index of biotic integrity (IBI) score was 36, and 15 species were reported. The macroinvertebrate community at S-7 was rated "exceptional." Based on the fish and macroinvertebrate community assessment, no biological impairment associated with chemical contaminants was observed at S-7. The physical habitat was also evaluated at S-7, and the QHEI score was 70, indicating "good" stream habitat capable of supporting WWH biological communities.

At S-9, all inorganic chemicals tested in sediment were below Ohio SRVs and TEC levels. All tested explosives, pesticides, and PCBs were not detected in sediment samples collected from S-9. The few SVOCs were measured at low levels, with all concentrations below TEC or ESL guidelines. Ammonia and total phosphorus levels were measured below screening guidelines. None of the surface water chemical concentrations at S-9 exceeded Ohio WQS aquatic life maximum or average

water quality criteria, and none of the chemicals measured exceeded criteria protective of the WWH aquatic life use (USACE 2005a). Overall, the sediment quality and water quality at S-9 was rated "excellent." The fish community at S-9 was rated "good." The IBI score was 43, and 19 species were reported. The macroinvertebrate community at S-9 was rated "exceptional." Based on the assessment attributes, low body burdens to fish and macroinvertebrates would be expected, given the positive conditions in the fish and macroinvertebrate communities. High fecundity and other measures of reproductive success would also be expected. The physical habitat was also evaluated at S-9, and the QHEI score was 71.5, indicating "good" stream habitat capable of supporting WWH biological communities. These favorable sediment/water quality findings at S-9 support the observation that the Dump Along Paris-Windham Road is not contributing contamination to Sand Creek.

11 12 13

14

1

2

3

4

5

6 7

8

9

10

Table 4-11. Comparison of Five Assessment Techniques at Sampling Stations Near the Dump Along Paris-Windham Road

Attributes	S-7 (RM 2.4) (upstream)	S-9 (RM 1.9) (downstream)	Comments
Sediment quality	Excellent	Excellent	Downstream station rating is equivalent to upstream station, suggesting no negative impacts from the AOC.
Water quality	Excellent	Excellent	Downstream station rating is equivalent to upstream station, suggesting no negative impacts from the AOC.
Fish community (IBI) ^a	Marginally Good	Good	Downstream station rating is slightly better to upstream station, suggesting no negative impacts from the AOC.
Macroinvertebrate community (ICI) ^b	Exceptional	Exceptional	Downstream station rating is equivalent to upstream station, suggesting no negative impacts from the AOC.
Habitat (QHEI) ^c	Good	Good	Downstream station rating is equivalent to upstream station, suggesting no negative impacts from the AOC.
Use Attainment Status ^d	Full	Full	Downstream station rating is equivalent to upstream station, suggesting no negative impacts from the AOC.

^aFish communities range from 0-60, with <18 being "very poor," 18-27 being "poor," 28-35 being "fair," 36-39 being "marginally good," 40-45 being "good," 46-49 being "very good," and 50-60 being "excellent" (Ohio EPA 2009).

AOC = Area of Concern

IBI = Index of Biotic Integrity

ICI = Invertebrate Community Index

Ohio EPA = Ohio Environmental Protection Agency

QHEI = Qualitative Habitat Evaluation Index

RM = River Mile

15 16

17

Ecosystem and Landscape Roles and Relationships. There are four spatial areas evaluated to assess the ecosystem and landscape roles and relationships: the AOC, the vicinity of the AOC, the

^bMacroinvertebrate communities range from 0-60 with <2 being "very poor," 2-12 being "poor," 14-32 being "fair," 34-46 being "good," and 48-60 being "exceptional" (Ohio EPA 1988).

^{&#}x27;Habitat ranges from 30 to <100 with <30 being "very poor," 30-44 being "poor," 45-59 being "fair,"

⁶⁰⁻⁷⁴ being "good," and 75-100 being "excellent" (Ohio EPA 2009).

^dFull-attainment means all of the applicable indices meet the Ohio EPA biocriteria (USACE 2005a).

entire RVAAP, and the northeastern or ecoregion of Ohio. Information about the first spatial area (the AOC) was provided in the subsections above on terrestrial and aquatic resources.

Vicinity of the AOC. Two vegetation communities border the Dump Along Paris-Windham Road (Figure 4-4): the green ash (Fraxinus pennsylvanica)/American elm (Ulmus americana)/hackberry (Celtis occidentalis and laevigata) temporary flooded forest alliance and the American beech (Fagus grandifolia)/oak (Quercus spp.)/and maple (Acer spp.) forest alliance. The AOCs bordering vegetation communities are similar to those observed at Dump Along Paris-Windham Road; there are no apparent differences in habitat quality of these plant communities inside or outside of the AOC. The types and qualities of habitat are not unique and can be found at many other areas at RVAAP. Figure 4-4 shows there are two wetlands along the western border of the AOC. Other wetlands are located to the west along the drainage swale (i.e., former stream channel), west along Sand Creek, and east of Paris-Windham Road. No perennial surface water features exist in the AOC boundary; however, Sand Creek is located 100-170 ft west of the AOC (Figure 4-4).

The closest recorded state-listed or federally-listed species [butternut (*Juglans cinerea*)] was located approximately 5 ft west of the northwestern border of the AOC (Table 4-12) (OHARNG 2008a); it is a state potentially threatened plant. The next closest recorded state-listed or federally-listed species [yellow-bellied sapsucker (*Sphyrapicus varius*)] was previously sighted about 500 ft southwest of the AOC; it is a state endangered species.

No beaver dams are in or near the AOC. There is a 100-year floodplain to Sand Creek located approximately 40 ft west of the AOC boundary, and there is a biological and water quality station (stream sampling station) within 60 ft of the AOC.

Table 4-12. Survey of Proximity to the AOC of Various Ecological Resources

Natural Resource	Inside Habitat Area	Near the AOC	Distances to Nearest Resource ^a and Comments
Wetlands (Planning Level Survey and Jurisdictional)	Two small jurisdictional wetlands	West along the drainage swale, west along Sand Creek, and east of Paris- Windham Road	Others in vicinity (Figure 4-4)
State-listed or Federally-listed Species	No known sightings	Along western border	5 ft west 500 ft southwest See text for species names
Beaver dams	None	None	1,600 ft north 1,700 ft north
100-year floodplain	None	Sand Creek floodplain located 40 ft to the west	100-year floodplain to Sand Creek located 40 ft west of the AOC
Stream sampling ^b	None	Sampling location (S-9) is located 60 ft north	An additional stream sampling location (S-7) is located approximately 1,900 ft upstream of the AOC
Pond sampling ^b	None	None	Nearest pond station at Cobbs Ponds about 2,000 ft south

^a Measurements of distance and direction are taken from the nearest boundary of the AOC to the resource being measured

AOC = Area of Concern

^bStream and pond sampling refers to Facility-Wide Biological and Water Quality Study 2003 (USACE 2005a)

The Entire RVAAP. The Dump Along Paris-Windham Road is approximately 0.25 acres in size, which represents 0.001% of the total area of RVAAP (21,683 acres). There are approximately 2,310 acres of forest type FL1 [temporarily flooded cold-deciduous forest alliance (e.g., green ash and American elm)] at RVAAP, based on the INRMP map (OHARNG 2008a); this represents 10.7% of the habitat at RVAAP. There are approximately 2,290 acres of forest type FU2 (American beech, oak, maple) (OHARNG 2008a), representing 10.6% of the habitat at RVAAP. There are approximately 1,990 acres of wetlands (jurisdictional and planning level survey) as defined in the INRMP (OHARNG 2008a), representing 9% of the habitat at RVAAP. These types of resources are abundant and are not unique to the Dump Along Paris-Windham Road at RVAAP.

Ecoregion. In the area surrounding RVAAP, forests occupy a high percentage of the terrain. Ohio's forests cover approximately 8,000,000 acres or 30% of the state (USDA 2009). The Erie/Ontario Drift and Lake Plain ecoregion (USEPA 2011) is located in the northeastern part of Ohio, and both contain the communities of temporarily flooded, cold-deciduous forest alliance (e.g., green ash and American elm) and American beech/oak/maple forest alliance. The Erie/Ontario Drift and Lake Plain ecoregion exhibits rolling to level terrain formed by lacustrine and low lime drift deposits. Lakes, wetlands, and swampy streams occur where stream networks converge or where the land is flat and clayey (USEPA 2011). The United States Forest Service (USFS) has a Forest Inventory Data Online tool that was queried for the forest types in the surrounding counties in or near RVAAP (USFS 2011). In 2009, approximately 93,900 acres of forest type FL1 and 621,100 acres of forest type FU2 were found throughout northwestern Ohio in Cuyahoga, Geauga, Mahoning, Portage, Stark, Summit, and Trumbull counties that surround RVAAP (USFS 2011). Wetlands across the ecoregion make up 207,800 acres (USEPA 1999). The vegetation communities and wetlands at the Dump Along Paris-Windham Road are also found in the surrounding counties in the ecoregion of northeastern Ohio.

In summary, the current vegetation types of temporarily flooded, cold-deciduous forest alliance (e.g., green ash and American elm); the American beech/oak/maple forest alliance; and wetlands are found in the vicinity of the Dump Along Paris-Windham Road. The two forest types and wetlands are in abundance at RVAAP and the larger surrounding local ecoregion. There is no known unique resource at the Dump Along Paris-Windham Road that cannot be found in the immediate vicinity of the AOC, RVAAP, and in the large part of the ecoregion of northeastern Ohio.

4.4.3.4 Evaluation of Chemical Contamination and Ecological Significance

There are three surface soil COPECs identified in the ERA for the Fill Area EU: mercury, zinc, and PCB-1254. There are eight surface soil COPECs identified in the ERA for the Surface Area EU: aluminum, cadmium, lead, manganese, mercury, zinc, nitrocellulose, and PCB-1254. There are three surface water COPECs identified in the ERA: manganese, mercury, and nitrocellulose (Section 4.4.3.2).

Section 4.4.3.3 provides information about presence of important ecological resources and the lack of significant ecological resources at the AOC. Approximately 0.25 acres of forest habitat exists within

43 the boundaries of the Dump Along Paris-Windham Road. The current forest community consists

primarily of green ash, American elm, and hackberry. Small wetlands are found at the western boundary of the AOC along a drainage swale. The entire extent of the wetlands is 0.21 acres; however, only 0.04 acres of the wetlands lie within the AOC boundary. Although the wetlands are an important resource, they are not a significant resource, as dry sediment and surface water sampling results in and around the wetlands (discussed in Section 4.4.3.3) do not indicate exposure to elevated concentrations of contaminants would occur within the wetlands/drainage swale (i.e., former stream channel). As a result, there are no significant ecological resources. Also, the downstream biological and water quality sampling station shows no impairment, indicating contaminants are not migrating from the landfill to Sand Creek.

4.4.3.5 Summary and Recommendations of Scoping Level Ecological Risk Assessment

While a removal action occurred in the Fill Area EU, the limited "RD/RA" confirmatory sample results indicate there are three surface soil COPECs for the Fill Area EU, eight surface soil COPECs for the Surface Area EU, and three surface water COPECs identified in the Surface Water EU at the Dump Along Paris-Windham Road. These COPECs consist of inorganic chemicals, PCBs, and propellants. There are no sediment COPECs at the AOC.

The information in Section 4.4.3.3 regarding ecological resources at the AOC was compared to the list of important ecological places and resources (Appendix Table C-6). One of the 39 important places (wetlands) was present. Although the wetlands are an important resource, the wetlands are not a significant resource, as dry sediment and surface water sampling results (Section 4.4.3.2) do not indicate chemicals are present at concentrations of concern for ecological receptors in the wetlands/drainage swale. Environmental management goals and objectives of OHARNG are applicable to the AOC, as presented in Appendix Table C-6. Some of the management goals benefit the AOC, including Goal 1 that requires management of natural resources to be compatible with the military mission, and Goal 5 that requires the U.S. Army to sustain usable training lands and native natural resources by implementing a natural resource management plan which incorporates invasive species management and by utilizing native species mixes for revegetation after ground disturbance activities.

The Dump Along Paris-Windham Road is approximately 0.25 acres and is vegetated with: (1) green ash/American elm/hackberry temporary flooded forest alliance; (2) American beech/oak/maple forest alliance; and (3) small wetlands. These same types of habitats are found adjacent to the AOC and elsewhere at RVAAP (OHARNG 2008a). The habitats are also found in the larger, local ecoregion that surrounds RVAAP (USFS 2011). There is no known unique resource at the AOC.

Although there is contamination at the AOC and an important ecological resource is present, the AOC has no known significant ecological places or resources. Also, the downstream biological and water quality sampling station shows no impairment, suggesting contaminants have not migrated from the landfill to Sand Creek. Consequently, the ERA for the Dump Along Paris-Windham Road can conclude with a Level I Scoping Level ERA and the recommendation of NFA from the ecological risk perspective.

4.4.4 Conclusions

There is chemical contamination present at the Dump Along Paris-Windham Road. While a removal action occurred in the Fill Area EU, the limited "RD/RA" confirmatory sample results indicate there are three surface soil COPECs at the Fill Area EU, eight surface soil COPECs at the Surface Area EU, and four surface water COPECs at the Surface Water EU. Although the wetlands are an important resource, they are not a significant resource because dry sediment and surface water sampling results do not indicate chemicals are present at concentrations of concern for ecological receptors in the wetlands/drainage swale. Thus, there are no significant ecological resources at the AOC. Also, the downstream biological and water quality sampling station shows no impairment, suggesting contaminants are not migrating from the landfill to Sand Creek. Further, the vegetation types are found elsewhere near the AOC, at RVAAP, and in the ecoregion. Based on the results of the ERA, there is sufficient justification to recommend NFA for the Dump Along Paris-Windham Road from the ecological perspective.

THIS PAGE INTENTIONALLY LEFT BLANK.

5.1 REMEDIAL ACTION OBJECTIVE

RAOs specify the requirements remedial alternatives must fulfill to protect human health and the environment from COCs at the Dump Along Paris-Windham Road. Media-specific objectives that identify major contaminants and associated media-specific cleanup goals (CUGs) are developed to provide this protection. These objectives specify COCs, exposure routes and receptors, and acceptable constituent concentrations for long-term protection of receptors.

In accordance with CERCLA, a residential receptor (Resident Farmer Adult and Child) was addressed in the risk assessment (see Section 4.3) as a comparative baseline. However, a remedial alternative based on Residential Land Use is not evaluated in this FFS. Similarly, a remedial alternative based on unrestricted land use is not evaluated in this FFS. The representative receptor for the Dump Along Paris-Windham Road is the Range Maintenance Soldier. The Adult and Juvenile Trespassers was also evaluated in the HHRA. No COCs are identified in soil and surface water for these receptors; however, COCs were identified in shallow surface soil for the Resident Farmer receptor. LUCs and awareness training is necessary as part of the final remedy due to future Camp Ravenna training missions, AOC characteristics, the presence of shallow surface soil COCs for a Resident Farmer receptor, and the presence of residual transite, all of which make unrestricted use of the AOC impractical. The HHRA identified no surface water COCs for any receptor; therefore, RAs are not required for surface water. The ERA recommended NFA for protection of ecological resources. As noted earlier, RAs for groundwater are not included in the alternatives evaluated in this FFS; groundwater will be addressed under a future decision by the U.S. Army.

Based upon the SC results, the RAO at the Dump Along Paris-Windham Road is to prevent exposure of the Resident Farmer to shallow surface soil (0-1 ft bgs) with COC levels exceeding the TR of 1E-05 and an HQ of 1.0.

5.2 REASONABLE AND ANTICIPATED FUTURE LAND USE

Under the Camp Ravenna training mission, OHARNG has established that the Dump Along Paris-Windham Road will be a part of the SDZ for a proposed future range complex (OHARNG 2008b). No range construction activities are proposed to be conducted within the AOC. The presence of the former dump and residual transite preclude placement of utilities along the west side of Paris-Windham Road in the vicinity of this AOC. The Range Maintenance Soldier is the representative receptor for this land use.

 Activities could also include foot traffic by range control (due to the fact the AOC is in the SDZ) and wildlife and natural resource management activities. Because this area does not have elevated security measures, trespassers may visit the AOC; therefore, Adult and Juvenile Trespassers were also evaluated. The National Guard Trainee is not considered an appropriate receptor for this area because

the AOC is a small area on a steep road berm and is not suitable for training use. However, the exposure assumptions for the Adult Trespasser are protective of foot traffic by the National Guard Trainee. Characteristics of the AOC (e.g., proximity to the road, steep slope, and floodplain at the bottom) preclude Residential Land Use. These considerations determined the selection of the representative receptors denoted in Section 4.3.3 for the most likely foreseeable land use.

5 6 7

1

2

3

5.3 FACILITY-WIDE CLEANUP GOALS

8

FWCUGs have been established in the FWCUG Report (USACE 2010a). These FWCUGs are the remediation levels for the designated user for any COCs at the Dump Along Paris-Windham Road, unless there are additive effects to be considered.

111213

14

10

The COCs identified at the Dump Along Paris-Windham Road are listed in Table 5-1. No COCs were identified in soil or surface water for the Range Maintenance Soldier or Adult and Juvenile Trespassers.

15 16 17

Table 5-1. Chemicals of Concern and Cleanup Goals by Media and Receptor

		FWCUG					
Media	COC	(mg/kg)					
Range M	Range Maintenance Solder						
Surface Soil (0-4 ft bgs)	None	NA					
Wet Sediment	None ^a	NA					
Surface Water	None ^b	NA					
Adult and	l Juvenile Trespasser						
Shallow Surface Soil (0-1 ft bgs)	None	NA					
Subsurface Soil (1-13 ft bgs)	None ^b	NA					
Wet Sediment	None ^a	NA					
Surface Water	None	NA					
Re	sident Farmer						
Shallow Surface Soil (0-1 ft bgs)	Benzo(a)pyrene	0.211 mg/kg					
Shahow Surface Son (0-1 it bgs)	Dibenz(a,h)anthracene	0.211 mg/kg					
Subsurface Soil (1-13 ft bgs)	None	NA					
Wet Sediment	None ^a	NA					
Surface Water	None	NA					

^aWet sediment does not exist within the boundaries of the area of concern. Dry sediment is addressed the same as surface soil in terms of contaminant nature and extent, fate and transport, and risk exposure models and is consistent with the FWCUG Report (USACE 2010a).

bgs = Below ground surface

COC = Chemical of Concern

FWCUG = Facility-Wide Cleanup Goal

NA = Not applicable

^bA complete exposure pathway does not exist for the specified receptor and media

6.0 APPLICABLE OR RELEVANT AND APPROPRIATE REQUIREMENTS

Agencies responsible for RAs under CERCLA must ensure selected remedies meet ARARs. This section describes the proposed ARARs for the Dump Along Paris-Windham Road.

6.1 Introduction

CERCLA Sections 121(d)(1) and (2) provide that RAs selected for an AOC must attain a degree of cleanup of hazardous substances, pollutants, and contaminants that: (1) assures protection of human health and the environment; and (2) complies with ARARs. ARARs are developed in accordance with the statutory and regulatory provisions set forth in CERCLA and the National Oil and Hazardous Substance Pollution Contingency Plan (NCP).

An RA will comply with ARARs if the RA attains the standard established in the ARAR for a particular hazardous substance. When a hazardous substance, pollutant, or contaminant will remain on-site at the completion of an RA, that substance must meet any limit or standard set forth in any legal ARAR, criteria, or limitation under a federal environmental law. These standards apply unless such standard, requirement, criteria, or limitation is waived in accordance with CERCLA Section 121(d)(4). Any promulgated standard, requirement, criteria, or limitation under a state environmental or facility citing law that is more stringent than any federal standard, requirement, criteria, or limitation, and that has been identified by the state in a timely manner, can be an ARAR as well.

Regulatory language interpreting and implementing the statutory directive is found in the NCP. One provision, 40 *Code of Federal Regulations (CFR)* Section 300.400(g), provides that the lead agency (U.S. Army) and support agency (Ohio EPA) shall identify applicable requirements based on an objective determination of whether the requirement specifically addresses a hazardous substance, pollutant, contaminant, RA, location, or other circumstance found at a CERCLA site. Under 40 *CFR* Section 300.430(e), the lead agency has the ultimate authority to decide what requirements are ARARs for the potential remedial activities.

Identifying ARARs involves determining whether a requirement is legally applicable, and (if it is not legally applicable) whether a requirement is relevant and appropriate. Individual ARARs for each AOC must be identified on a site-specific basis. Applicable requirements are those cleanup standards, standards of control, and other substantive environmental protection requirements, criteria, or limitations promulgated under federal or state environmental or facility siting laws that specifically address a hazardous substance, pollutant, contaminant, RA, location, or other circumstance found at a CERCLA site (40 *CFR* Section 300.5).

 If a requirement is determined to not be legally applicable to a specific release, the requirement may still be relevant and appropriate to the circumstances of the release. Determining whether a rule is relevant and appropriate is a two-step process that involves determining whether the rule is relevant, and appropriate. A requirement is relevant if it addresses problems or situations sufficiently similar to the circumstances of the RA contemplated. It is appropriate if its use is well suited to the AOC. In addition to ARARs, the lead and support agencies may identify other advisories, criteria, or guidance to be considered for a particular release. The "to be considered" category consists of advisories, criteria, or guidance that were developed by USEPA, other federal agencies, or states that may be useful in developing CERCLA remedies. "To be considered" will be regarded as guidance or justification for a standard used in the remediation if no other standard is available for a situation to help determine the necessary level of cleanup for protection of health or the environment.

While on-site actions must comply with both applicable and relevant and appropriate requirements, off-site actions taken outside of the CERCLA site must fully comply with the regulations in their entirety, including any administrative requirements. Also, a determination of relevance and appropriateness may be applied to specific portions of a requirement so that only parts of a requirement need be complied with; whereas, a determination of applicability is made for the requirement as a whole so that the entire requirement must be complied with.

CERCLA provides for a permit waiver for RAs that are conducted on-site and in accordance with NCP. Although the administrative requirement of permits has been waived by the statute, substantive requirements of rules that would otherwise be enforced through permits are still applicable. The Ohio EPA Division of Emergency and Remedial Response (DERR) has addressed this issue in two policies, one in final form and one in draft form. The policy in final form, Final Policy Number DERR-00-RR-001, "ARARs," July 30, 1998, states "cleanup projects will not be subject to the administrative requirements of permits, including permit applications, public notice, etc." particularly when the cleanup project is governed by an enforcement order. The policy in draft form, Draft Policy Number DERR-00-RR-034, "Use of ARARs in the Ohio EPA Remedial Response Program," September 2, 2003, states "it has been DERR's policy to require responsible parties to acquire and comply with all necessary permits, including all substantive and administrative requirements." Permit waivers are specifically addressed in Section VII, General Provisions (Paragraph No. 12e) of the DFFO:

"It is Ohio EPA's position that if state law related to a remedial or removal action requires a permit, then a permit must be acquired in accordance with CERCLA Section 120(a)(4). It is the Respondent's position that these Orders implement a CERCLA-based remediation program and that a permit is not required in accordance with CERCLA Section 121(e). The Parties agree that the remedial or removal actions anticipated at RVAAP are not of the type that routinely requires a permit under state law. If Ohio EPA determines that a permit is required for a particular remedial or removal action at RVAAP, the Parties will meet and attempt in good faith to resolve to [sic] this issue."

Any remedial response action at RVAAP must be conducted in accordance with the DFFOs, which provide that, irrespective of ARARs, "all activities undertaken...pursuant to these Orders shall be

performed in accordance with the requirements of CERCLA, the NCP, and all other applicable federal and state laws and regulations."

6.2 POTENTIAL APPLICABLE OR RELEVANT AND APPROPRIATE REQUIREMENTS

USEPA classifies ARARs as chemical-specific, action-specific, and location-specific to provide guidance for identifying and complying with ARARs (USEPA 1988).

• Chemical-specific ARARs are health- or risk-based numerical values or methodologies which, when applied to site-specific conditions, allow numerical values to be established. These values establish the acceptable amount or concentration of a chemical that may be found in, or discharged to, the ambient environment.

• Action-specific ARARs are rules, such as performance or design or other activity-based rules, that place requirements or limitations on actions.

• Location-specific ARARs are rules that place restrictions on the concentration of hazardous substances or the conduct of activities solely because they occur in special locations.

As explained in the following paragraph, rules from each of these categories are ARARs only to the extent they relate to the degree of cleanup.

CERCLA Section 121 governs cleanup standards at CERCLA sites. ARARs originate in the subsection of CERCLA that specifies the degree of cleanup at each AOC, CERCLA Section 121(d). In Section 121(d)(2), CERCLA expressly directs that ARARs are to address specific COCs at each AOC, specifying the level of protection to be attained by any chemicals remaining at the AOC. CERCLA Section 121(d)(2) provides that, with respect to hazardous substances, pollutants, or contaminants remaining on-site at the completion of an RA, an ARAR is:

"any standard, requirement, criteria, or limitation under any Federal environmental law...or any promulgated standard, requirement, criteria, or limitation under a State environmental or facility siting law that is more stringent than any Federal standard, requirement, criteria, or limitation."

CERCLA Section 121(d)(2) further provides that the RA attain a level of control established in rules determined to be ARARs.

- In some cases, most ARARs will be chemical-specific, depending on the identified COCs and (in some cases) the media that have been contaminated from the release of these contaminants (e.g.,
- 40 MCLs for groundwater contamination). Action- or location-specific requirements will be ARARs to
- 41 the extent they establish standards addressing COCs that will remain at the AOC. In addition,
- 42 CERCLA Section 121(d)(1) directs that RAs taken to achieve a degree of cleanup that is protective of
- 43 human health and the environment are to be relevant and appropriate under the circumstances

presented by the release. Accordingly, any chemical-, action-, or location-specific requirements will be ARARs to the extent that they ensure that the degree of cleanup will be protective of human health and the environment under the circumstances presented by the release.

In summary, chemical-, action-, or location-specific requirements will be ARARs to the extent: (1) they establish standards protective of human health and the environment for chemicals that will remain on-site after the RA; and (2) to the extent they ensure a degree of cleanup that is protective of human health and the environment under the circumstances presented by the release.

6.3 POTENTIAL CHEMICAL-SPECIFIC APPLICABLE OR RELEVANT AND APPROPRIATE REOUIREMENTS

The actions evaluated within this FFS address the potential risk from contaminated soil at the Dump Along Paris-Windham Road. A review of regulations governing the remediation of soil did identify the requirements of 40 *CFR* 761 governing PCBs as a potential ARAR. However, these standards were found to not be applicable or relevant and appropriate, as the highest concentration of PCBs found within the soil was 0.23 mg/kg, and the guidance documents from USEPA concerning the Toxic Substances Control Act (TSCA) requirements state that triggering of these requirements is based on a finding of unacceptable risk during the risk assessment. As no such finding was made, the TSCA requirements for PCBs at 40 *CFR* Part 761 were deemed not to be an ARAR (chemical- or action-specific). Due to the nature of the AOC, the identified COCs, and the media of concern, no chemical-specific ARARs were identified.

6.4 POTENTIAL ACTION-SPECIFIC APPLICABLE OR RELEVANT AND APPROPRIATE REQUIREMENTS

The proposed actions at the Dump Along Paris-Windham Road include No Action and LUC alternatives. Neither of these activities would result in the excavation or generation of contaminated soil. By leaving the soil in place, no potential waste treatment or disposal requirement is triggered; therefore, the Resource Conservation and Recovery Act (RCRA) and TSCA requirements would not be considered ARARs for either of these actions.

Although passive in nature, the potential presence of transite (an ACM) would trigger certain provisions within the OAC for inactive asbestos disposal sites. Even though sampling has not indicated that asbestos was present, past operating knowledge indicates such material is in the landfill that would result in these rules being considered (at minimum) relevant and appropriate. The requirements of OAC 3745-20-07 are considered ARARs for both the No Action and the LUC alternatives (see Table 6-1). These regulations require that a former asbestos waste disposal site must be covered and posted in accordance with the specific requirements. Because all visible surface debris was removed and the subsurface transite excavation areas were backfilled and covered with clean soil and vegetated (Section 3.1.3), the cover requirements have been achieved in compliance with this ARAR. In addition to the cover requirements, these rules specify the AOC must be posted as a former asbestos disposal site. The No Action alternative would not comply with this

requirement. However, such posting is anticipated to be conducted under the LUC alternative and would comply with the posting requirements of this ARAR.

As previously discussed, the Dump Along Paris-Windham Road historically was used as a waste disposal site. Under the evaluated alternatives, the disposed waste will remain in place. Based on the majority of the material observed within the AOC and the accompanying analytical data, the dumpsite is believed to have been used primarily for the disposal of debris from C&D activities during its operation. Because the exact historical dates of operation are unknown but thought to have been as late as the 1970s, the solid waste landfill closure requirements (OAC 3745-27-11) have not been identified as ARARs for this facility. Although the closure requirements have not been identified as ARARs, the requirements of OAC 3745-27-14 (landfill post-closure for existing facilities) are considered an applicable requirement and an ARAR (see Table 6-1). The No Action alternative would not include provisions such as quarterly inspections [OAC 3745-27-14(c)(4)]. Based on available information and observations noted above, the material disposed of consisted of debris that would not result in the generation of methane or leachate; therefore, this subparagraph is the only provision within this rule identified as an ARAR.

Table 6-1. Potential Action ARARs for Disposal of RCRA Hazardous Waste

Media and Citation	Description of Requirement	Potential ARAR Status	Standard
Standard for	These rules require that	If ACM is present	An inactive asbestos disposal site
Inactive Asbestos	inactive asbestos disposal	within the AOC,	must be covered by 6 inches of
Waste Disposal	sites be covered and posted	these rules are	compacted soil with a vegetated
Sites	to ensure access to ACM is	potentially	cover or 2 ft of compacted soil. In
OAC 3745-20-07	controlled. In addition,	applicable	addition, the AOC must be posted as
	these rules require that no		having ACM present and must have
	visible emissions be		access control to ensure exposure to
	allowed from the AOC		asbestos does not occur
Post-Closure	These rules specify the	Because material	Required inspection and
Care for Sanitary	required post-closure care	that would be	maintenance of the cover.
Landfill Facilities	activities required for solid	considered solid	Additional provisions are not
OAC 3745-27-14	waste facilities, including	waste is disposed at	considered ARARs, as the debris
	existing facilities	the AOC, these	disposed at the AOC does not
		requirements are	generate methane gas or leachate
		considered relevant	
		and appropriate	

ACM = Asbestos-containing Material

AOC = Area of Concern

ARAR = Applicable or Relevant and Appropriate Requirement

OAC = Ohio Administrative Code

RCRA = Resource Conservation and Recovery Act

6.5 POTENTIAL LOCATION-SPECIFIC APPLICABLE OR RELEVANT AND APPROPRIATE REQUIREMENTS

Location requirements include those established for potential remedial activities conducted within wetlands or a floodplain area or with respect to threatened and endangered (T&E) species. Generally, for wetlands and floodplains, rules require alternatives to remedial activity within the sensitive area

be pursued. If that is not feasible, then adverse effects from any actions taken within the sensitive area must be mitigated to the extent possible. These requirements do not relate to specific chemicals nor do they further the degree of cleanup in the sense of protecting human health or the environment from the effects of harmful substances. Rather, their purpose is to protect the sensitive areas to the extent possible. Under CERCLA Section 121(d), relevance and appropriateness are related to the circumstances presented by the release of a hazardous substance, with the goal of attaining a degree of cleanup and control of further releases that ensures protection of human health and the environment.

Rules ensuring protection of sensitive resources do not represent requirements that are relevant and appropriate to circumstances presented by the release of a hazardous substance, with a goal of attaining a degree of cleanup and control of further releases that ensure protection of human health and the environment. Location requirements for wetlands and floodplains do not relate to the degree of cleanup as much as they relate to the protection of these sensitive areas from the effects of remedial activities. This purpose does not address problems or situations sufficiently similar to those encountered at the CERCLA site that their use is well suited to the particular AOC as an ARAR; that is, the rule requirements are not sufficiently relevant and appropriate under CERCLA Section 121(d) as related to the circumstances of the release, degree of cleanup, or protectiveness of RA, to include these requirements as ARARs.

The Endangered Species Act (ESA) exists to protect the habitat or body of flora and fauna that are T&E. Once again, these rules do not relate to specific chemicals nor do they further the degree of cleanup in the sense of protecting human health or the environment from the effects of harmful substances. The purpose of these rules is to protect sensitive areas and plant and animal life to the degree possible. This purpose does not address problems or situations sufficiently similar to those encountered at the CERCLA site that their use is well suited to the particular AOC as an ARAR; that is, the rule requirements are not sufficiently relevant and appropriate under CERCLA Section 121(d) as related to the circumstances of the release, degree of cleanup, or protectiveness of RA to include these requirements as ARARs.

Having determined these requirements are not ARARs, any action taken by the Federal Government must be conducted in accordance with requirements established under the National Environmental Policy Act, National Historic Preservation Act, ESA, and federal and state wetlands and floodplains construction and placement of material considerations, even though these laws and rules do not establish standards, requirements, limitations, or criteria relating to the degree of cleanup for chemicals remaining on-site at the close of the response action. As the No Action or LUC alternatives would not result in impacts upon endangered species or their habitats, these ARARs would be complied with.

7.0 TECHNOLOGY TYPES AND PROCESS OPTIONS

This section describes the general response actions (GRAs) and remedial technologies that are potentially applicable at the Dump Along Paris-Windham Road. GRAs are actions that will satisfy the RAO (Section 5.1) for shallow surface soil. Given no COCs were identified for the representative receptor (Range Maintenance Soldier) or the Adult and Juvenile Trespassers, and NFA has been recommended for ecological receptors, the appropriate GRAs for this FFS are No Action (as required by the NCP) and LUCs (to prevent exposure to human receptors). The residential receptor (Resident Farmer) has two PAHs identified as COCs, thus preventing unrestricted land use.

7.1 NO ACTION

In this GRA, no action would be undertaken to reduce any hazard to human health or the environment. Any current actions, restrictions, or monitoring would be discontinued. This action complies with the CERCLA requirement to provide an appropriate option (or component of a remedial alternative if no unacceptable risks are present) and to provide a baseline against which other alternatives can be compared.

7.2 LAND USE CONTROLS

Generally, LUCs reduce the potential for exposure to contaminants but do not reduce contaminant volume or toxicity. These controls are utilized to supplement and affect the engineering component(s) of a remedy (e.g., treatment and removal) during short- and long-term implementation. The primary goal of LUCs is to restrict the use of, or limit access to, real property using physical, legal, and/or administrative mechanisms to ensure protectiveness of the remedy. Particular LUCs under consideration at the Dump Along Paris-Windham Road include measures that will restrict land use changes over the long term, such as governmental controls and enforcement tools. Governmental controls could include a Property Management Plan (PMP) and facility-specific regulations to manage property and enforce management strategies, while enforcement tools may involve administrative orders or consent decrees.

THIS PAGE INTENTIONALLY LEFT BLANK.

8.0 DEVELOPMENT OF REMEDIAL ALTERNATIVES

1 This section describes the remedial alternatives assembled for impacted shallow surface soil at the

- 2 Dump Along Paris-Windham Road. Remedial alternatives should assure adequate protection of
- 3 human health and the environment; achieve the RAO; meet ARARs; and permanently and
- 4 significantly reduce the volume, toxicity, and/or mobility of COCs. The remedial alternatives are
- 5 listed below.

6 7

- Alternative 1: No Action; and
- 8 Alternative 2: LUCs.

9

Alternative 1 is the No Action response required under the NCP. Alternative 2 relies on LUCs. No source control or removal actions are implemented under Alternative 2.

12

8.1 NO ACTION

13 14 15

16

17

18

19

20

Under Alternative 1, no actions regarding access or LUCs would be implemented. Alternative 1 provides no additional protection to human health and the environment. This remedial alternative is required under the NCP as a no action baseline against which other remedial alternatives can be compared. Any current legal and administrative LUC mechanisms at the AOC would be discontinued. No future legal, administrative, or physical LUC mechanisms would be employed at the AOC. Environmental monitoring would not be performed. In addition, no restrictions on land use would be pursued.

212223

8.2 LAND USE CONTROLS

2425

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

For Alternative 2, LUCs would be implemented for the Dump Along Paris-Windham Road. This alternative relies on LUCs to limit access to the AOC and prevent exposure by possible receptors (e.g., Resident Farmer) to COCs in shallow surface soil. Unrestricted land use of the AOC is hindered by concentrations of benzo(a)pyrene and dibenz(a,h)anthracene in shallow surface soil, which exceed FWCUGs for the Resident Farmer. However, no COCs were identified for the Range Maintenance Soldier (the representative receptor at the AOC as determined by the RAFLU) or the possible Adult and Juvenile Trespassers. Alternative 2 would leave impacted media in place and implement no active remedial measures. Instead, long-term management to ensure land use remains protective of potential receptors would be implemented. Awareness training and signs (posted every 300 ft or less along the AOC perimeter) would be employed to alert persons having a need to access the AOC that the location was formerly used to dispose of ACM. Controls on digging within the AOC would be incorporated due to the potential presence of ACM and to maintain integrity of restored sections of the dump. Because: (1) surface debris was removed; (2) subsurface transite was excavated to the extent possible without undermining and compromising the integrity of Paris-Windham Road; (3) soil confirmation samples did not indicate the presence of asbestos in soil, dry sediment, or surface water; and (4) the AOC is heavily vegetated, potential exposures to asbestos are

currently controlled, and physical access controls other than warning signs (e.g., fencing/gates) are not proposed as part of Alternative 2. Prior to implementation of Alternative 2, an RD detailing the 5-year review requirements and any supplemental access restrictions to address chemical contamination of soil would be developed.

An RD would be developed to address specific maintenance activities, monitoring requirements (i.e., 5-year reviews), and LUCs. The RD would incorporate existing access restrictions. A more detailed discussion of the LUCs would be developed as part of the RD, including notification requirements for changes in land use. The RVAAP PMP would capture all LUCs prescribed by the approved RD and serve as a formal tool to help manage and set forth procedures for the established LUCs. Coordination with any planned OHARNG AOC improvement and environmental monitoring activities would be necessary to ensure consistency with the Dump Along Paris-Windham Road's designated land use and RAO. Pursuant to CERCLA, a review would be conducted every 5 years, as COCs would remain on-site above unrestricted (i.e., residential) land use CUGs. Five-year reviews permit evaluation of all remedy components, including LUCs, to assess the presence and behavior of the remaining COCs. Continued surveillance would ensure any land use changes or disturbances of impacted areas are identified.

9.0 ANALYSIS OF REMEDIAL ALTERNATIVES

1	9.1 Introduction	
2		
3	This section presents a detailed analysis of the two remedial alternatives for the Dump Along Pa	
4 5	Windham Road. Under the CERCLA remedy selection process, the preferred remedial alternative will be suggested in the PP and set forth in final form in the ROD. A detailed evaluation of each of the preferred remedial alternative will be suggested in the PP and set forth in final form in the ROD.	
6	alternative is performed in this section to provide the basis and rationale for identifying a prefe	
7	remedy and preparing the PP.	1100
8		
9	To ensure the analysis of alternatives provides information of sufficient quality and quantity to just	stify
10	the selection of a remedy, it is helpful to understand the requirements of the remedy selection proc	•
11	This process is driven by the requirements set forth in CERCLA Section 121. In accordance	
12	these requirements (USEPA 1988), RAs must:	
13		
14	• Be protective of human health and the environment;	
15		
16	• Attain ARARs;	
17		
18	• Be cost effective;	
19		
20	• Use permanent solutions and alternative treatment technologies to the maximum ex	tent
21	practicable; and	
22		
23	• Satisfy the preference for treatment that, as a principle element, reduces volume, toxicity	, or
24	mobility.	
25		
26	CERCLA emphasizes long-term effectiveness and related considerations for each reme	dial
27	alternative. These statutory considerations are as follows:	
28		
29	 Long-term uncertainties associated with land disposal; 	
30		
31	• The goals, objectives, and requirements of the Solid Waste Disposal Act;	
32		
33	• The persistence, toxicity, and mobility of hazardous substances and their propensity	/ to
34	bio-accumulate;	
35		
36	• Short- and long-term potential for adverse health effects from human exposure;	
37	· ·	
38	• Long-term maintenance costs;	
39		

40

The potential for future RA costs if the remedial alternative in question was to fail; and

• The potential threat to human health and the environment associated with excavation, transportation, re-disposal, or containment.

2 3 4

5

6 7

1

These statutory requirements are implemented through the use of nine evaluation criteria presented in the NCP. These nine criteria are grouped into threshold criteria, balancing criteria, and modifying criteria, as described below. A detailed analysis of each alternative against the evaluation criteria is presented in the following sections. The detailed analysis provides further definition of each alternative (if necessary), compares the alternatives against one another, and presents considerations common to alternatives.

9 10 11

8

9.1.1 Threshold Criteria

12

- 13 Two of the NCP evaluation criteria relate directly to statutory findings that must be made in the ROD.
- 14 These criteria are considered to be threshold criteria that must be met by any remedy selected. The

15 criteria are:

16 17

- 1. Overall protection of human health and the environment; and
- 2. Compliance with ARARs.

18 19

Each alternative must be evaluated to determine how it achieves and maintains protection of human health and the environment. Similarly, each remedial alternative must be assessed to determine how it complies with ARARs or, if a waiver is required, an explanation of why a waiver is justified. An alternative is considered to be protective of human health and the environment if it complies with CUGs.

2526

9.1.2 Balancing Criteria

2728

The five balancing criteria represent the primary criteria upon which the detailed analysis of alternatives and the comparison of alternatives are based. They are:

29 30

- 31 1. Long-term effectiveness and permanence;
- 32 2. Reduction of toxicity, mobility, or volume through treatment;
- 33 3. Short-term effectiveness;
- 34 4. Implementability; and
- 35 5. Cost.

- 37 Long-term effectiveness and permanence is an evaluation of the magnitude of residual risk (risk
- 38 remaining after implementation of the alternative) and the adequacy and reliability of controls used to
- 39 manage the remaining waste (untreated waste and treatment residuals) over the long term.
- 40 Alternatives that provide the highest degree of long-term effectiveness and permanence leave little or
- 41 no untreated waste at the AOC, make long-term maintenance and monitoring unnecessary, and
- 42 minimize the need for LUCs.

Reduction of toxicity, mobility, or volume through treatment is an evaluation of the ability of the alternative to reduce the toxicity, mobility, or volume of the waste. The irreversibility of the treatment process and the type and quantity of residuals remaining after treatment are also assessed.

Short-term effectiveness addresses the protection of workers and the community during the RA, the environmental effects of implementing the action, and the time required to achieve CUGs.

Implementability addresses the technical and administrative feasibility of implementing an alternative and the availability of various services and material required during implementation. Technical feasibility assesses the ability to construct and operate a technology, the reliability of the technology, the ease in undertaking additional RAs, and the ability to monitor the effectiveness of the alternative. Administrative feasibility is addressed in terms of the ability to obtain approval from federal, state, and local agencies.

Cost analyses provide an estimate of the dollar cost of each alternative. The cost estimates in this report are based on estimating reference manuals, historical costs, vendor quotes, and engineering estimates. Costs are reported in base year 2010 dollars. The present value analysis is a method to evaluate expenditures, either capital or operation and maintenance (O&M), which occur over different time periods. Present value calculations allow for cost comparisons of different remedial alternatives on the basis of a single cost figure. The cost estimates are for guidance in project evaluation and implementation and are believed to be accurate within a range of -30 to +50%, in accordance with USEPA guidance (USEPA 1988). Actual costs could be higher than estimated due to unexpected conditions or potential delays. Details and assumptions used in developing cost estimates for Alternative 2 are provided in Appendix D.

9.1.3 Modifying Criteria

The two modifying criteria below will be evaluated as part of the ROD after the public has had an opportunity to comment on the PP. They are:

31 1. State acceptance; and

2. Community acceptance.

State acceptance considers comments received from agencies of the state of Ohio. The primary state agency supporting this investigation is Ohio EPA. Comments will be obtained from state agencies on the SC/FFS and the preferred remedy presented in the PP. This criterion will be addressed in the responsiveness summary of the ROD.

- Community acceptance considers comments made by the community, including stakeholders, on the alternatives being considered. Input has been encouraged during the ongoing investigation process to ensure the remedy ultimately selected for the Dump Along Paris-Windham Road is acceptable to the
- 42 public. Comments will be accepted from the community on the preferred remedy presented in the PP.
- 43 This criterion will be addressed in the responsiveness summary of the ROD. Because the actions

above have not yet taken place, the detailed analysis of alternatives presented below cannot account for these criteria at this time. Therefore, the detailed analysis is carried out only for the first seven of the nine criteria.

9.2 INDIVIDUAL ANALYSIS OF ALTERNATIVES

Detailed analyses of the retained remedial alternatives for the Dump Along Paris-Windham Road are presented below. Each relevant alternative is described and evaluated against the criteria outlined in Section 9.1.

9.2.1 Alternative 1: No Action

Under this alternative, contaminated shallow surface soil would remain in place. Existing access restrictions (e.g., the RVAAP perimeter fence) would not be continued. No restrictions on land use would be pursued.

9.2.1.1 Overall Protection of Human Health and the Environment

Under Alternative 1, the representative receptor (Range Maintenance Soldier) and possible Adult and Juvenile Trespassers would not be exposed to unacceptable risk due to contaminants in shallow surface and subsurface soil at the AOC. However, the AOC has COC concentrations above CUGs for the Resident Farmer. Consequently, a No Action alternative would not be protective, as LUCs would be required to prevent Residential Land Use of the AOC while the COC concentrations exceed Resident Farmer COCs. Alternative 1 is not considered protective for human health.

The ERA concluded there is chemical contamination and possible risk but no significant ecological resources at the AOC, and the recommendation is NFA for protection of ecological receptors. Under Alternative 1, current risk is not reduced and the ecological resources at the AOC remain unchanged. Current land use and RAFLU allows for sustainability of terrestrial habitat for ecological receptors.

9.2.1.2 Compliance with ARARs

Potential ARARs for the final remedy of shallow surface soil at the Dump Along Paris-Windham Road are presented in Section 6.0. There are no identified chemical- or location-specific ARARs for Alternative 1.

OAC 3745-20-07 requires that a former asbestos waste disposal site must be covered and posted in accordance with the specific requirements. Because all visible surface debris was removed and the excavation areas covered with clean soil and vegetated, the cover requirements have been achieved in compliance with this ARAR. However, in addition to the cover requirements, these rules specify the AOC must be posted as a former asbestos disposal site. The No Action alternative would not comply with this requirement, as no signs would be posted at the AOC.

9.2.1.3 Long-Term Effectiveness and Permanence

Alternative 1 includes no long-term management measures to prevent exposures to, or the spread of, contamination. This alternative does not have controls in place outside the existing cover over portions excavated during the limited "RD/RA" and does not provide any additional new controls in the future.

9.2.1.4 Reduction of Toxicity, Mobility, or Volume through Treatment

Alternative 1 does not reduce contaminant toxicity, mobility, or volume because no treatment process is proposed.

9.2.1.5 Short-Term Effectiveness

There are no significant short-term human health risks associated with Alternative 1. No additional short-term health risks to the community would occur because no RAs would be implemented. There would be no transportation risks nor would workers be exposed to any additional health risks. Alternative 1 would not directly cause adverse impacts to soil, air quality, water resources, or biotic

19 resources.

9.2.1.6 Implementability

No RAs would be implemented under this alternative.

9.2.1.7 <u>Cost</u>

The present value cost to complete Alternative 1 is \$0. No capital costs are associated with this alternative.

9.2.2 Alternative 2: Land Use Controls

Alternative 2 maintains the current status of the Dump Along Paris-Windham Road and includes LUCs and annual inspections to identify potential exposures and/or changes in the nature or extent of AOC contamination. LUCs would be implemented in accordance with an approved RD and PMP. In addition, signs would be posted at the AOC stating that the area was a former ACM disposal location.

Pursuant to CERCLA, a review would be conducted every 5 years, as contaminants remain on-site above unlimited use and unrestricted exposure FWCUGs. These 5-year reviews will evaluate the effectiveness of LUCs and ensure any land use changes are identified.

9.2.2.1 Overall Protection of Human Health and the Environment

Under Alternative 2, the representative receptor (Range Maintenance Soldier) and possible Adult and Juvenile Trespassers are not exposed to unacceptable risk due to contaminants in shallow surface or subsurface soil at the AOC. Implementation of LUCs prevents exposure to the Resident Farmer. Alternative 2 is considered protective for human receptors.

The ERA concluded there is chemical contamination and possible risk but no significant ecological resources at the Dump Along Paris-Windham Road, and the recommendation is NFA for protection of ecological receptors. Under Alternative 2, current risk is not reduced and the ecological resources at the AOC remain unchanged. Current land use and RAFLU allow for sustainability of terrestrial habitat for ecological receptors.

9.2.2.2 Compliance with ARARs

Potential ARARs for the final remedy of shallow surface soil at the Dump Along Paris-Windham Road are presented in Section 6.0. These enforceable standards would be protective of representative receptors under the Range Maintenance Soldier and Trespasser scenario. There are no identified chemical- or location-specific ARARs for Alternative 2.

OAC 3745-20-07 requires that a former asbestos waste disposal site must be covered and posted in accordance with the specific requirements. Because all visible surface debris was removed and the excavation areas covered with clean soil and vegetated, the cover requirements have been achieved in compliance with this ARAR. In addition to the cover requirements, these rules specify that the AOC must be posted as a former asbestos disposal site. Alternative 2 would comply with this posting requirement.

9.2.2.3 Long-Term Effectiveness and Permanence

Alternative 2 is protective in the long term. It relies on LUCs to eliminate or reduce exposures to contaminants. The effectiveness of this approach is related to the adequacy and reliability of the LUCs. However, with appropriate documentation and procedures, LUCs can reasonably be expected to be effective in protecting human health and the environment while preserving the RAFLU anticipated for the Dump Along Paris-Windham Road.

Because contaminants would remain on-site above Resident Farmer CUGs, reviews would need to be conducted every 5 years, pursuant to CERCLA requirements. The purpose of these reviews is to ensure that land use and engineering controls are retaining effectiveness.

9.2.2.4 Reduction of Toxicity, Mobility, or Volume through Treatment

Alternative 2 does not involve reduction in contaminant toxicity, mobility, or volume because no treatment is proposed.

Short-Term Effectiveness 9.2.2.5

1 2 3

4

5

6

There are no significant short-term human health risks associated with Alternative 2. No additional short-term health risks to the community would occur because no RAs would be implemented. Alternative 2 would not directly cause adverse impacts on soil, air quality, water resources, or biotic resources. The alternative's remedial measures would require less than 1 year to complete and would include an O&M period (30 years assumed for cost-estimating purposes).

7 8 9

9.2.2.6 **Implementability**

10 11

12

13

LUCs are technically implementable. No technical difficulties are anticipated in establishing or maintaining monitoring programs, signs, or access restrictions. There are currently access restrictions implemented facility-wide at RVAAP. Implementing proposed LUCs would supplement and support restrictions already in place at the Dump Along Paris-Windham Road.

14 15 16

9.2.2.7 <u>Cost</u>

17 18

19 20

The present value (discounted) cost to complete Alternative 2 is approximately \$93,384 (in base year 2010 dollars). O&M and monitoring costs are estimated for a 30-year period. The development of a RD, including LUCs and CERCLA 5-year reviews, is included in this cost. A detailed description of Alternative 2 costs is contained in Appendix D.

21 22

9.3 **COMPARATIVE ANALYSIS**

23 24 25

A comparison of the two alternatives for the Dump Along Paris-Windham Road is presented in Table 9-1.

26 27 28

Table 9-1. Comparison of Alternatives by Evaluation Criteria

NCP Evaluation Criteria	Alternative 1:No Action	Alternative 2: LUCs
Overall Protectiveness for Human Health and the Environment	Somewhat protective	Protective
2. Compliance with ARARs	Not compliant	Compliant
3. Long-Term Effectiveness and Permanence	Low	High
4. Reduction of Toxicity, Mobility, or Volume through Treatment	Low	Low
5. Short-Term Effectiveness	High	Medium
6. Implementability	High	Medium
7. Cost	High	Medium

[&]quot;High" = highly favorable

²⁹ 30 "Medium" = moderately favorable 31

[&]quot;Low" = not favorable

³² ARAR = Applicable or relevant and appropriate requirement

³³ LUC = Land Use Control 34

NCP = National Oil and Hazardous Substances Pollution Contingency Plan

THIS PAGE INTENTIONALLY LEFT BLANK.

10.0 AGENCY COORDINATION AND PUBLIC INVOLVEMENT

- The U.S. Army is the lead agency under the Defense Environmental Restoration Program responsible for achieving remedy of the Dump Along Paris-Windham Road. This section reviews actions that have been conducted and that are planned in the future to ensure regulatory agencies and the public have been provided with appropriate opportunities to stay informed of progress of the AOC's
- 5 remediation and to provide meaningful input on the planning effort and final selection of a remedy.

As described in Section 9.0, two of the nine NCP evaluation criteria are known as "modifying criteria." These are state acceptance and community acceptance. These criteria provide a framework for obtaining the necessary agency coordination and public involvement in the remedy selection process.

10.1 STATE ACCEPTANCE

State acceptance considers comments received from agencies of the state of Ohio on the remedial alternatives being considered. For the process supporting remedy of the Dump Along Paris-Windham Road, Ohio EPA is the lead regulatory agency, and this SC/FFS has been prepared in consultation with Ohio EPA. Ohio EPA has provided input during the ongoing investigation and report development process to ensure the remedy selected for this AOC meets the needs of the state of Ohio and fulfills the requirements of the DFFO (Ohio EPA 2004). Comments will be solicited from Ohio EPA on this SC/FFS and on the PP. The U.S. Army will obtain Ohio EPA concurrence prior to selecting the final remedy for the Dump Along Paris-Windham Road.

10.2 COMMUNITY ACCEPTANCE

Community acceptance considers comments provided by the community on the remedial alternatives being considered. CERCLA 42 U.S. Code 9617(a) emphasizes early, constant, and responsive community relations. The U.S. Army has prepared a *Community Relations Plan* (USACE 2003b) for RVAAP to ensure the public has convenient access to information regarding project progress. The community relations program interacts with the public through news releases; public meetings; and Restoration Advisory Board meetings with local officials, interest groups, and the general public.

The public also is provided the opportunity to comment on draft documents submitted to the Administrative Record that support remedy of the Dump Along Paris-Windham Road.

CERCLA 42 U.S. Code 9617(a) requires that an Administrative Record be established "at or near the facility at issue." Relevant documents regarding RVAAP have been made available to the public for review and comment. The Administrative Record for this project is available at the following location:

1	Ravenna Army Ammunition Plant
2	Building 1037 Conference Room
3	8451 State Route 5
4	Ravenna, Ohio 44266-9297
5	
6	Access to RVAAP is restricted but can be obtained by contacting facility management at (330) 358
7	7311. In addition, an Information Repository of current information and final documents is available
8	to any interested reader at the following libraries:
9	
10	Reed Memorial Library
11	167 East Main Street
12	Ravenna, Ohio 44266
13	
14	Newton Falls Public Library
15	204 South Canal Street
16	Newton Falls, Ohio 44444-1694
17	
18	Also, RVAAP has an online resource for restoration news and information. This website is available
19	at: <www.rvaap.org>.</www.rvaap.org>
20	
21	Similar to state agencies, comments will be received from the community upon issuance of this FFS
22	and the PP. The U.S. Army will request public comments on the PP for the Dump Along Paris-
23	Windham Road, as required by the CERCLA regulatory process and the RVAAP Community
24	Relations Plan. These comments will be considered in the final selection of a remedy for the Dump
25	Along Paris-Windham Road. Responses to these comments will be addressed in the responsiveness
26	summary of the ROD.

11.0 CONCLUSIONS

11.1 CONCLUSIONS

The primary purpose of this SC/FFS is to: (1) evaluate the nature and extent of contamination at the Dump Along Paris-Windham Road following the limited "RD/RA" using data collected during previous investigations; (2) determine the potential risk to appropriate human and ecological receptors; and (3) develop, screen, and evaluate remedial alternatives in compliance with the CERCLA process. This SC/FFS examined the history of the Dump Along Paris-Windham Road, summarized previous investigations, outlined CUGs and RAO for the AOC, and identified alternatives potentially applicable for meeting these CUGs.

The RAFLU for the Dump Along Paris-Windham Road is Restricted Access, which includes continued access restrictions associated with the covered dumpsite. Chemical-specific CUGs were identified for the representative receptor (Range Maintenance Soldier), Adult and Juvenile Trespassers, and Resident Farmer. CUGs were identified for a Resident Farmer to provide a baseline for evaluating whether this AOC may be eligible for unrestricted land use.

This SC/FFS establishes the RAO for the Dump Along Paris-Windham Road and evaluates RAs to reduce risks to the environment to obtain a final remedy with respect to shallow surface soil. The RAO analysis identified COCs in impacted shallow surface soil at the Dump Along Paris-Windham Road that require further evaluation of potential remedial alternatives for Residential Land Use. The RAO analysis indicates the representative receptor (Range Maintenance Soldier) and Adult and Juvenile Trespassers do not have COCs in media at the AOC, and the RAFLU is protective with respect to impacted shallow surface soil. NFA is recommended for the protection of ecological resources within the AOC. However, COCs were identified for the Resident Farmer; therefore, the following potential remedial alternatives were developed:

- Alternative 1: No Action; and
- Alternative 2: LUCs.

These alternatives were assessed and compared against one another to provide information of sufficient quality and quantity to justify the selection of a remedy.

11.2 RECOMMENDED ALTERNATIVE

The recommended alternative for the final remedy of the Dump Along Paris-Windham Road is Alternative 2: LUCs. COCs do not exist for the representative receptor for the RAFLU (Range Maintenance Soldier) and Adult and Juvenile Trespassers. However, COCs exist within shallow surface soil for the Resident Farmer; therefore, LUCs are required to ensure protection of this receptor. ACM is also known to be present within the subsurface. Alternative 2 fully complies with ARARs by including signs alerting persons of the presence of ACM and offers long-term

effectiveness and permanence when implemented and maintained. Alternative 2 is easily implementable in a relatively short time frame and is expected to have a discounted cost of approximately \$93,384.

The next step in the CERCLA process is to prepare a PP to solicit public input regarding the remedial alternatives. The PP will present alternatives evaluated in the FFS together with the preferred alternative for the Dump Along Paris-Windham Road.

The ROD will document the remedy for the Dump Along Paris-Windham Road. Comments on the PP received from state and federal agencies and the public will be considered in drafting the ROD for the AOC. The ROD will provide a brief summary of the history, characteristics, risks, and selected remedy. The ROD also will include a responsiveness summary, which addresses comments received on the PP.

12.0 REFERENCES

1	United States Army Biological Technical Assistance Group (BTAG) 2005. Technical Document for
2 3	Ecological Risk Assessment: Process for Developing Management Goals. August 2005.
4	United States Department of Energy (DOE) 1997. Preliminary Remediation Goals for Ecological
5	Endpoints. Oak Ridge National Laboratory, Oak Ridge Tennessee. August 1997.
6	
7	Environmental Quality Management (EQM) 2010. Facility-Wide Groundwater Monitoring Program
8	Report on the January 2010 Sampling Event, Ravenna Army Ammunition Plant, Ravenna,
9	Ohio. July 2010.
10	
11 12	MacDonald, D.D., C.G., Ingersoll, and T.A., Berger. 2000. Development and Evaluation of Consensus-based Sediment Quality Guidelines for Freshwater Ecosystems, Arch. Environ.
13	Contam. Toxicol. 39, 20-31. 2000.
14	
15	MKM Engineers, Inc. (MKM) 2004. Final Report for Remedial Design/Remedial Action Plan at
16 17	Paris-Windham Road Dump. March 2004.
18	Ohio Army National Guard (OHARNG) 2008a. Integrated Natural Resources Management Plan
19	and Environmental Assessment for the Ravenna Training and Logistics Site, Portage and
20	Trumbull Counties, Ohio. March 2008.
21	
22	OHARNG 2008b. Land Use Summary Table, personal communication from Katie Elgin OHARNG.
23	August 14, 2008.
24	
25	Ohio Environmental Protection Agency (Ohio EPA) 1988. Biological Criteria for the Protection of
26	Aquatic Life: Vol. II: User's Manual for Biological Field Assessment of Ohio Surface
27	Waters. Table 8-3. February 1988.
28	
29	Ohio EPA. 1989. Biological Criteria for the Protection of Aquatic Life: Vol. III: Standardized
30	Biological Field Sampling and Laboratory Methods for Assessing Fish and
31	Macroinvertebrate Communities. September 1989.
32	
33	Ohio EPA 2001. Ohio Rapid Assessment Method for Wetlands v. 5.0, User's Manual and Scoring
34	Forms. Division of Surface Water, 401/Wetland Ecology Unit. February 2001.
35 36	Ohio EPA 2003. Guidance for Conducting Ecological Risk Assessments (Ohio EPA). Division of
37	Emergency and Remedial Response. February 2003.
38	Emergency and Remedian Response. Teorum y 2005.
39	Ohio EPA 2004. Director's Final Findings and Orders for the Ravenna Army Ammunition Plant.
40	June 2004.
41	

1 2	Ohio EPA 2008. <i>Guidance for Conducting Ecological Risk Assessments (Ohio EPA)</i> . Division of Emergency and Remedial Response. April 2008.
3	Emergency and remedial response. April 2000.
4 5	Ohio EPA 2009. <i>Total Maximum Daily Loads for the White Oak Creek Watershed.</i> Division of Surface Water. December 2009.
6	
7	Rankin, E.T. 1989. The Qualitative Habitat Evaluation Index (QHEI): Rationale, Methods, and
8 9	Application. Div. Water Qual. Plan. & Assess. Ecol. Assess. Sect. Columbus, Ohio. 1989.
10	
11	Rankin, E. T. 1995. The use of habitat assessments in water resource management programs, pp.
12	181-208. in W. Davis and T. Simon (eds.). Biological Assessment and Criteria: Tools for
13	Water Resource Planning and Decision Making. Lewis Publishers, Boca Raton, FL. 1995.
14	
15	United States Army Corps of Engineers (USACE) 1999. Plant Community Survey for the Ravenna
16	Army Ammunition Plant Summary Report. Prepared for Ohio Army National Guard,
17	Adjutant General's Department, Columbus, Ohio. August 1999.
18	
19	USACE 2001a. Facility-Wide Sampling and Analysis Plan for Environmental Investigations at the
20	Ravenna Army Ammunition Plant, Ravenna, Ohio. March 2001.
21	
22	USACE 2001b. Phase II Remedial Investigation Report for the Winklepeck Burning Grounds at the
23	Ravenna Army Ammunition Plant, Ravenna, Ohio. April 2001.
24	
25	USACE 2003a. Decision Document for a Removal Action at Paris-Windham Road Dumpsite
26	(RVAAP-51). USACE, Louisville District, KY. 2003.
27	
28	USACE 2003b. Ravenna Army Ammunition Plant, Ravenna, Ohio Community Relations Plan.
29	September 2003.
30	
31	USACE 2003c. RVAAP Facility-wide Ecological Risk Work Plan. April 2003.
32	
33	USACE 2005a. Facility-Wide Biological and Water Quality Study 2003 Ravenna Army Ammunition
34	Plant, Ravenna, Ohio, Part I-Streams, Part II-Ponds. November 2005.
35	
36	USACE 2005b. RVAAP Facility-Wide Human Health Risk Assessor Manual, Amendment 1.
37	December 2005.
38	
39	USACE 2008. Wet and Dry Sediment Definition, email communication from Thomas M. Chanda,
40	USACE, Louisville District. May 27, 2008.
41	
42	USACE 2010a. Facility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition
43	Plant, Ravenna, Ohio. March 2010.

44

USACE 2010b. Risk Assessment Handbook Volume II: Environmental Evaluation. December 2010.
USACE 2010c. Site Characterization and Focused Feasibility Study Work Plan for the RVAAP-51
Dump Along Paris-Windham Road. August 2010.
U.S. Army Center for Health Promotion and Preventive Medicine (USACHPPM) 1998. Relative Risk
Site Evaluation for Newly Added Sites at the Ravenna Army Ammunition Plant, Ravenna,
Ohio. Hazardous and Medical Waste Study No. 37-EF-5360-99. October 1998.
omo. Hazardous and Medicar Waste Study 110. 37 Er 3300 77. October 1770.
United States Department of Agriculture (USDA) 2009. Forest Service. Resource Bulletin NRS-36.
http://www.nrs.fs.fed.us/. 2009.
r.
United States Environmental Protection Agency (USEPA) 1987. Health Advisory for Nitrocellulose.
Office of Drinking Water. September 1987.
USEPA 1988. Guidance for Conducting Remedial Investigations/Feasibility Studies Under
CERCLA. October 1988.
USEPA 1989. Risk Assessment Guidance for Superfund, Volume 1: Human Health Evaluation
Manual (Part A). December 1989.
USEPA 1993. Wildlife Exposure Factors Handbook. EPA/600/R-93/187a. Office of Research and
Development, Washington, D.C. Volume 1 of 2. December 1993.
USEPA 1996. Ecological Significance and Selection of Candidate Assessment Endpoints. ECO
Update. Volume 3, Number 1. EPA 540/F-95/037. January 1996.
USEPA 1997. Ecological Risk Assessment Guidance for Superfund: Process for Designing and
Conducting Ecological Risk Assessments. Office of Solid Waste and Emergency Response,
Washington, D.C. June 1997.
USEPA 1999. Ohio Wetland Restoration and Mitigation Strategy Blueprint. Wetland Grant Program
Federal Grant No. CD985853-01-0. Prepared by Ohio Department of Natural Resources and
Ohio Environmental Protection Agency. September 1999.
Omo Environmental Protection rigency. September 1999.
USEPA 2003. Region 5 RCRA Ecological Screening Levels. http://www.epa.gov/reg5rcra/ca/ESL.pdf.
28 August 2003.
USEPA 2010. EPA Regional Screening Level (RSL). Website:
http://www.epa.gov/reg3hwmd/risk/human/rb-concentration_table/index.htm. November
2010.

1	USEPA 2011. Ecoregions of EPA Region	n 3: Delaware, Maryla	nd, Pennsylvania, V	Virginia, and West
2	Virginia.	Western	Ecology	Division.
3	http://www.epa.gov/wed/pages/ed	coregions/reg3_eco.htm	. October 2011.	
4	United States Forest Service (USFS) 201	1. Forest Inventory Do	uta Online (FIDO).	Forest Inventory
5	and Analysis National Program.	. http://www.fia.fs.fed	l.us/tools-data/defa	ult.asp. February
6	2011.			

APPENDIX A

Photographs

SAIC Site Photographs August 2009



Photograph 1. View from Paris-Windham Road Bridge over Sand Creek (sample location S-9 on left side of creek)



Photograph 2. Looking South Along Paris-Windham Road (AOC on right)



Photograph 3. Dense Growth at North End of AOC



Photograph 4. Dense Growth Downgradient of Former Dump Site



Photograph 5. Drainage Swale Facing East with No Standing Water (sheen from high moisture content of sediment)



Photograph 6. Drainage Swale Facing Southeast with No Standing Water (sheen from high moisture content of sediment)



Photograph 7. View of Floodplain Located West of Drainage Swale



Photograph 8. Toe of Slope in Vicinity of Grid 5

SAIC Site Photographs November 2011



Photograph 9. View to West of Northern Wetland from Paris-Windham Road (Grid 2)



Photograph 10. View to West of Sand Creek Floodplain from Paris-Windham Road (Grid 4)



Photograph 11. View to Southwest of Southern Wetland from Paris-Windham Road (Grid 6)



Photograph 12. View to Southwest of Southern Wetland from Paris-Windham Road (excavation and fill area in Grids 8-9 in foreground; Grid 10 in background left)



Photograph 13. View to Northeast of Northern Wetland from Sand Creek Floodplain (Grids 1-3 and Paris-Windham Road in background)



Photograph 14. View to North of Sand Creek Floodplain between Northern and Southern Wetlands (Grids 1-5 on right)



Photograph 15. View to East of Grid 4 from Sand Creek Floodplain (Paris-Windham Road in background)



Photograph 16. View to Southeast of Southern Wetland from Sand Creek Floodplain (excavation and fill area in Grids 6-8 in background)



Photograph 17. View to Southeast of Southern Wetland from Sand Creek Floodplain (Grids 9 and 10 in background)



Photograph 18. View to Southeast of Southern Wetland from Sand Creek Floodplain (Grids 7-10 on left)

APPENDIX B

Human Health Risk Assessment Supporting Data

List of Tables

SRC and COPC	Screening
---------------------	-----------

Table B-1.	SRC and COPC Screening for Subsurface Soil (> 2 ft bgs Discrete Samples) at Paris-Windham Dump AOC: Fill Area EU
Table B-2.	SRC and COPC Screening for Surface Soil (0-1 ft bgs Discrete Samples) at
Table D 2	Paris-Windham Dump AOC: Surface Area EU
Table B-3. Table B-4.	SRC and COPC Screening for Soil (ISM Samples) at Paris-Windham Dump AOC
COC Scree	ning: Soil – Range Maintenance Soldier
Table B-5.	COC Screening for Surface Soil (0-1 ft bgs Discrete Samples) at Paris-Windham Dump AOC: Surface Area EU
	Representative Receptor: Range Maintenance Soldier
Table B-6.	COC Screening for Subsurface Soil (>2 ft bgs Discrete Samples) at Paris-Windham Dump AOC: Fill Area EU
Table B-7.	Representative Receptor: Range Maintenance Soldier
14010 2 71	Representative Receptor: Range Maintenance Soldier
COC Scree	ning: Soil – Trespasser
Table B-8.	COC Screening for Surface Soil (0-1 ft bgs Discrete Samples) at Paris-Windham Dump AOC: Surface Area EU
Table B-9.	Representative Receptor: Trespasser
COC Scree	ning: Soil – Resident Farmer
Table B-10.	COC Screening for Surface Soil (0-1 ft bgs Discrete Samples) at Paris-Windham Dump AOC: Surface Area EU
	Baseline Receptor: Resident Farmer
Table B-11.	COC Screening for Soil at Paris-Windham Dump AOC: ISM Samples Baseline Receptor: Resident Farmer
COC Scree	ning: Surface Water – Trespasser
Table B-12.	COC Screening for Surface Water at Paris-Windham Dump AOC
	Representative Receptor: Trespasser
COC Scree	ning: Surface Water – Resident Farmer
Table B-13.	COC Screening for Surface Water at Paris-Windham Dump AOC
	Baseline Receptor: Resident Farmer 13

Table B-1. SRC and COPC Screening for Subsurface Soil (> 2 ft bgs Discrete Samples) at Paris-Windham Dump AOC: Fill Area EU

	CAS	_				Background	SRC?		(HQ= 0			Risk Screening	Screening Level	COPC?		Station at	Date Collected at
Analyte (mg/kg)	Number	Detect	Detect	Detect	Result	Criteria ^a	(yes/no)	SRC Justification	RFA	RFC	NGT	Level	Source	(yes/no)	COPC Justification	Max Detect	Max Detect
Inorganic Chemi									#2 00			l vom			P	1/20/2002	
Aluminum	7429-90-5	5/ 5	6500	11000	8240	17700	No	Below background	52923	7380	3496	3496	NGT	No	Below background	PWss-005	4/29/2003
Arsenic	7440-38-2	5/ 5	9.2	13	11.2	15.4	No	Below background	0.425	0.524	2.78	0.425	RFA	No	Below background	PWss-008	4/29/2003
Barium	7440-39-3	5/ 5	47	180	76.6	88.4	Yes	Exceeds background	8966	1413	351	351	NGT	No	Below risk screening criteria	PWss-005	4/29/2003
Beryllium	7440-41-7	5/ 5	0.34	1.2	0.576	0.88	Yes	Exceeds background				16	RSL	No	Below risk screening criteria	PWss-005	4/29/2003
Calcium	7440-70-2	5/ 5	1500	39000	9860	15800	No	Essential Nutrient				1000000	RDA	No	Essential Nutrient	PWss-005	4/29/2003
Chromium ^d	7440-47-3	5/ 5	8.3	11	9.96	17.4	No	Below background	90.4	19.9	1.64	1.64	NGT	No	Below background	PWss-008	4/29/2003
Cobalt	7440-48-4	5/ 5	4.3	7.1	5.98	10.4	No	Below background	803	131	7.03	7.03	NGT	No	Below background	PWss-009	4/28/2003
Copper	7440-50-8	5/ 5	9.3	19	14	17.7	Yes	Exceeds background	2714	311	25368	311	RFC	No	Below risk screening criteria	PWss-005	4/29/2003
Iron	7439-89-6	5/ 5	14000	22000	18000	23100	No	Essential Nutrient	19010	2313	184370	180000	RDA	No	Essential Nutrient	PWss-005	4/29/2003
Lead	7439-92-1	5, 5	14	19	16.2	26.1	No	Below background				400	RSL	No	Below background	PWss-005	4/29/2003
Magnesium	7439-95-4	5, 5	1500	6100	2580	3030	No	Essential Nutrient				1000000	RDA	No	Essential Nutrient	PWss-005	4/29/2003
Manganese	7439-96-5	<i>z</i> , 5	390	880	540	1450	No	Below background	1482	293	35.1	35.1	NGT	No	Below background	PWss-005	4/29/2003
Mercury	7439-97-6	5, 5	0.025	0.048	0.036	0.036	Yes	Exceeds background	16.5	2.27	172	2.27	RFC	No	Below risk screening criteria	PWss-006	4/29/2003
Nickel	7440-02-0	5, 5	10	21	14.2	21.1	No	Below background	1346	155	12639	155	RFC	No	Below background	PWss-009	4/28/2003
Potassium	7440-09-7	<i>z</i> , 5	740	1100	892	927	No	Essential Nutrient				1000000	RDA	No	Essential Nutrient	PWss-005	4/29/2003
Sodium	7440-23-5	5, 5	130	380	202	123	No	Essential Nutrient				1000000	RDA	No	Essential Nutrient	PWss-005	4/29/2003
Vanadium	7440-62-2	5, 5	10	15	12.2	31.1	No	Below background	156	44.9	2304	44.9	RFC	No	Below background	PWss-008	4/29/2003
Zinc	7440-66-6	5, 5	50	100	66.8	61.8	Yes	Exceeds background	19659	2321	187269	2321	RFC	No	Below risk screening criteria	PWss-005	4/29/2003
	-	3/						Semi-volatile Organic (Сотроип	ds					-		
Acenaphthylene	208-96-8	1/ 1	0.13	0.13	0.13		Yes	Detected organic				340	RSL	No	Below risk screening criteria	PWss-009	4/28/2003
Anthracene	120-12-7	1/ 1	0.12	0.12	0.12		Yes	Detected organic				1700	RSL	No	Below risk screening criteria	PWss-009	4/28/2003
Benz(a)anthracene	56-55-3	1, 1	1	1	1		Yes	Detected organic	0.221	0.65	4.77	0.221	RFA	Yes	Exceeds screening level	PWss-009	4/28/2003
Benzo(a)pyrene	50-32-8	1/ 1	1.3	1.3	1.3		Yes	Detected organic	0.022	0.065	0.477	0.022	RFA	Yes	Exceeds screening level	PWss-009	4/28/2003
Benzo(b)fluoranthene	205-99-2	1/ 1	1.2	1.2	1.2		Yes	Detected organic	0.221	0.65	4.77	0.221	RFA	Yes	Exceeds screening level	PWss-009	4/28/2003
Benzo(ghi)perylene	191-24-2	1/ 1	0.75	0.75	0.75		Yes	Detected organic				1.5	RSL	No	Below risk screening criteria	PWss-009	4/28/2003
Benzo(k)fluoranthene	207-08-9	1/ 1	1.4	1.4	1.4		Yes	Detected organic	2.21	6.5	47.7	2.21	RFA	No	Below risk screening criteria	PWss-009	4/28/2003
Chrysene	218-01-9	1/ 1	1.1	1.1	1.1		Yes	Detected organic	22.1	65	477	22.1	RFA	No	Below risk screening criteria	PWss-009	4/28/2003
Dibenz(a,h)anthracene	53-70-3	1/ 1	0.24	0.24	0.24		Yes	Detected organic	0.022	0.065	0.477	0.022	RFA	Yes	Exceeds screening level	PWss-009	4/28/2003
Fluoranthene	206-44-0	1/ 1	1.7	1.7	1.7		Yes	Detected organic	276	163	5087	163	RFC	No	Below risk screening criteria	PWss-009	4/28/2003
Indeno(1,2,3-cd)pyrene	193-39-5	1/ 1	0.75	0.75	0.75		Yes	Detected organic	0.221	0.65	4.77	0.221	RFA	Yes	Exceeds screening level	PWss-009	4/28/2003
Phenanthrene	85-01-8	1/ 1	0.32	0.32	0.32		Yes	Detected organic				170	RSL	No	Below risk screening criteria	PWss-009	4/28/2003
Pyrene	129-00-0	1/ 1	1.4	1.4	1.4		Yes	Detected organic	207	122	3815	122	RFC	No	Below risk screening criteria	PWss-009	4/28/2003
		-1/						Pesticides/PCI				•			6		
PCB-1254	11097-69-1	1, 1	0.23	0.23	0.23		Yes	Detected organic	0.203	0.12	3.46	0.12	RFC	Yes	Exceeds screening level	PWss-009	4/28/2003
		' 1/ -															

^aBackground criteria for soil >1 ft bgs from final facility-wide background values for RVAAP, published in th*Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio* (USACE 2001).

 $RDA = Concentration \ associated \ with \ recommended \ daily \ allowance \ of \ essential \ nutrient$

RFA = FWCUG for Resident Farmer Adult

RFC = FWCUG for Resident Farmer Child

RSL = United States Environmental Protection Agency Residential Regional Screening Level

^dFWCUG is the most conservative (smallest) of the FWCUGs for hexavalent and trivalent chromium.

AOC = Area of Concern bgs = Below ground surface CAS = Chemical Abstract Service COPC = Chemical of Potential Concern

EU = Exposure Unit

HQ = Hazard Quotient

PCB = Polychlorinated Biphenyl SRC = Site-related Contaminant

-- = no value available

Bold = chemical is a COPC

^bFacility-Wide Cleanup Goals (FWCUGs) for Resident Farmer Adult (RFA), Resident Farmer Child (RFC), and National Guard Trainee (NGT) from acility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition Plant (USACE 2010).

^cScreening Level Source:

Table B-2. SRC and COPC Screening for Surface Soil (0-1 ft bgs Discrete Samples) at Paris-Windham Dump AOC: Surface Area EU

Numinum		CAS		Minimum			Background	SRC?		$(\mathbf{HQ} = 0)$	ning FW	k=1E-6)	Risk Screening	Screening Level	COPC?		Station at	Date Collected at
Aluminum	Analyte (mg/kg)	Number	Detect	Detect	Detect	Result	Criteria"	(yes/no)			KFC	NGI	Level	Source	(yes/no)	COPC Justification	Max Detect	Max Detect
Antimory	Aluminum	7420 00 5	12/ 12	5200	19000	9250	17700		7290	2406	2406	NCT	Voc	Evenede comoning level	DW/99 002	4/28/2003		
Arsenic 7440-38-2 17 13 2.6 13 7.53 15.4 No. Below background 0.425 0.524 2.78 0.425 RFA No. Below background PWs-010 Berjillium 7440-43-9 17 13 0.33 19 0.556 0.88 Ves Exceeds background FWs-02 2.73 13 15 No. Below risk screening criteria PWs-002 Cadmium 7440-43-9 13 0.1 0.59 0.204 0 Ves Exceeds background 16 RSI. No. Below risk screening criteria PWs-002 Cadmium 7440-43-9 13 1700 55000 6550 15800 No. Essential Nutrient 1000000 RDA No. Essential Nutrient PWs-002 Chromium 7440-43-3 17 13 1700 55000 6550 15800 No. Essential Nutrient 1000000 RDA No. Essential Nutrient PWs-002 Choalt 7440-43-3 17 13 1700 745000 14800 23100 No. Below background 904 199 1.64 1.64 NGT No. Below background PWs-010 Copper 7440-59-8 13 13 12000 18000 14800 23100 No. Essential Nutrient PWs-010 RDA No. RESential Nutrient PWs-010 RDA No. RDA			15/													Ü		4/28/2003
Barium									ŭ							Ü		4/28/2003
Beryllium					_				9							Ü		4/28/2003
Cadinium									,							- U		4/28/2003
Calcium			15/						ŭ							č		4/29/2003
Chromium			47						Ü							Ü		4/28/2003
Cobat			15/															
Copper									ŭ							- U		4/29/2003
Iron																Ü		4/28/2003
Ecad																č		4/29/2003
Magnesium																		4/29/2003
Manganese									ŭ									4/28/2003
Mercury 7439-97-6 73 73 0.045 0.08 0.0631 0.036 Yes Exceeds background 16.5 2.27 172 2.27 RFC No Below risk screening criteria PWsd-002 PWsd-005 PW	- U																	4/28/2003
Nickel 7440-02-0 13/13 9.9 37 19.1 21.1 Yes Exceeds background 1346 155 12639 155 RFC No Below risk screening criteria PWsd-005			10/						ū							U		4/28/2003
Potassium					0.00		0.000					_						4/29/2003
Silver 7440-22-4									ŭ							Ŭ		4/29/2003
Sodium			10															4/29/2003
Vanadium										324	38.6					Ŭ		4/28/2003
Zinc 7440-66-6 13/ 13 51 120 81.7 61.8 Yes Exceeds background 19659 2321 187269 2321 RFC No Below risk screening criteria PWsd-002 Explosives			11/															4/28/2003
Nitrocellulose 9004-70-0 1/2 1 2 2 2 2 2 2 2 2									ŭ							Ü		4/29/2003
Nitrocellulose 9004-70-0 1 1 2 2 2 2 Yes Detected organic 23000000 RSL No Below risk screening criteria PWsd-004	Zinc	7440-66-6	13/ 13	51	120	81.7	61.8	Yes	8		2321	187269	2321	RFC	No	Below risk screening criteria	PWsd-002	4/29/2003
Semi-volatile Organic Compounds Semi-volatile Organic Compounds Semi-volatile Organic Compounds										es						<u> </u>		
Benz(a)anthracene 56-55-3	Nitrocellulose	9004-70-0	1/ 1	2	2	2		Yes					23000000	RSL	No	Below risk screening criteria	PWsd-004	4/29/2003
Benzo(a)pyrene 50-32-8									Semi-volatile Organi	с Сотроі	ınds							
Benzolb/fluoranthene 205-99-2	Benz(a)anthracene		1/ 1			0.25		Yes								Exceeds screening level		4/29/2003
Benzo(k)fluoranthene 207-08-9	Benzo(a)pyrene		1/ 1		0.33	0.33			Detected organic	0.022	0.065					Exceeds screening level	PWsd-004	4/29/2003
Chrysene 218-01-9 1 1 0.33 0.34 0.35	Benzo(b)fluoranthene	205-99-2	1/ 1	0.39	0.39	0.39		Yes	Detected organic	0.221	0.65	4.77	0.221	RFA	Yes	Exceeds screening level	PWsd-004	4/29/2003
Fluoranthene 206-44-0 1/1 0.44 0.44 0.44 Yes Detected organic 276 163 5087 163 RFC No Below risk screening criteria PWsd-004 Pyrene 129-00-0 1/1 0.44 0.44 0.44 Yes Detected organic 207 122 3815 122 RFC No Below risk screening criteria PWsd-004 PWsd-004 PCB-1254 11097-69-1 1/1 0.086 0.086 0.086 0.086 Yes Detected organic 0.203 0.12 3.46 0.12 RFC No Below risk screening criteria PWsd-004	Benzo(k)fluoranthene	207-08-9	1/ 1	0.33	0.33	0.33		Yes	Detected organic	2.21	6.5	47.7	2.21	RFA	No	Below risk screening criteria	PWsd-004	4/29/2003
Pyrene 129-00-0 1/1 0.44 0.44 0.44 Yes Detected organic 207 122 3815 122 RFC No Below risk screening criteria PWsd-004	Chrysene	218-01-9	1/ 1			0.33			Detected organic		65				No	Below risk screening criteria	PWsd-004	4/29/2003
Pesticides/PCBs PCB-1254 11097-69-1 1 0.086 0.086 Yes Detected organic 0.203 0.12 RFC No Below risk screening criteria PWsd-004	Fluoranthene	206-44-0	1/ 1	0.44	0.44	0.44		Yes	Detected organic	276	163	5087	163	RFC	No	Below risk screening criteria	PWsd-004	4/29/2003
PCB-1254 11097-69-1 1/2 0.086 0.086 0.086 0.086 Yes Detected organic 0.203 0.12 3.46 0.12 RFC No Below risk screening criteria PWsd-004	Pyrene	129-00-0	1/ 1	0.44	0.44	0.44		Yes			122	3815	122	RFC	No	Below risk screening criteria	PWsd-004	4/29/2003
			1/						Pesticides/F	PCBs						<u> </u>		
Volatile Organic Compounds	PCB-1254	11097-69-1	1/ 1	0.086	0.086	0.086		Yes	Detected organic	0.203	0.12	3.46	0.12	RFC	No	Below risk screening criteria	PWsd-004	4/29/2003
voulite Organic Compounts			17						Volatile Organic C	Compound	ls							·
Acetone 67-64-1 1 0.041 0.041 0.041 - Yes Detected organic 6100 RSL No Below risk screening criteria PWsd-004	Acetone	67-64-1	1/ 1	0.041	0.041	0.041		Yes	Detected organic				6100	RSL	No	Below risk screening criteria	PWsd-004	4/29/2003

^aBackground criteria for soil 0-1 ft bgs from final facility-wide background values for RVAAP, published in the internal Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio (USACE 2001).

RDA = Concentration associated with recommended daily allowance of essential nutrient

RFA = FWCUG for Resident Farmer Adult

RFC = FWCUG for Resident Farmer Child

RSL = United States Environmental Protection Agency Residential Regional Screening Level

^dFWCUG is the most conservative (smallest) of the FWCUGs for hexavalent and trivalent chromium.

AOC = Area of Concern

HQ = Hazard Quotient

bgs = Below ground surface CAS = Chemical Abstract Service PCB = Polychlorinated Biphenyl SRC = Site-related Contaminant

COPC = Chemical of Potential Concern

-- = no value available

EU = Exposure Unit

Bold = chemical is a COPC

^bFacility-Wide Cleanup Goals (FWCUGs) for Resident Farmer Adult (RFA), Resident Farmer Child (RFC), and National Guard Trainee (NGT) from actility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition Plant (USACE 2010). ^cScreening Level Source:

Table B-3. SRC and COPC Screening for Soil (ISM Samples) at Paris-Windham Dump AOC

Analyte (mg/kg)	CAS Number	Freq of Detect	Minimum Detect	Maximum Detect	Average Result	Background Criteria ^a	SRC? (yes/no)	SRC Justification		ning FW .1 or Ris	k=1E-6)	Risk Screening Level	Screening Level Source ^c	COPC? (yes/no)	COPC Justification	Station at Max Detect		Sample ID at Max Detect
								Semi	-volatile	Organic (Сотроин	ds						
1,4-Dichlorobenzene	106-46-7	1/ 2	0.23	0.23	0.16		Yes	Detected organic				2.4	RSL	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
2-Methylnaphthalene	91-57-6	2/ 2	0.0055	0.064	0.0348		Yes	Detected organic	238	30.6	2384	30.6	RFC	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Acenaphthene	83-32-9	1/ 2	0.12	0.12	0.0688		Yes	Detected organic				340	RSL	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Acenaphthylene	208-96-8	2/ 2	0.056	0.12	0.088		Yes	Detected organic	-			340	RSL	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Anthracene	120-12-7	2/ 2	0.041	0.22	0.131		Yes	Detected organic	-			1700	RSL	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Benz(a)anthracene	56-55-3	2/ 2	0.36	1	0.68		Yes	Detected organic	0.221	0.65	4.77	0.221	RFA	Yes	Exceeds screening level	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Benzo(a)pyrene	50-32-8	2/ 2	0.46	1.4	0.93		Yes	Detected organic	0.022	0.065	0.477	0.022	RFA	Yes	Exceeds screening level	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Benzo(b)fluoranthene	205-99-2	2/ 2	0.5	1.4	0.95		Yes	Detected organic	0.221	0.65	4.77	0.221	RFA	Yes	Exceeds screening level	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Benzo(ghi)perylene	191-24-2	2/ 2	0.3	0.79	0.545		Yes	Detected organic	-			1.5	RSL	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Benzo(k)fluoranthene	207-08-9	2/ 2	0.45	1.4	0.925		Yes	Detected organic	2.21	6.5	47.7	2.21	RFA	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Bis(2-ethylhexyl)phthalate	117-81-7	1/ 2	0.025	0.025	0.0575		Yes	Detected organic				35	RSL	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Carbazole	86-74-8	1/ 2	0.19	0.19	0.14		Yes	Detected organic	69.4	44.6	835	44.6	RFC	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Chrysene	218-01-9	2/ 2	0.41	1.2	0.805		Yes	Detected organic	22.1	65	477	22.1	RFA	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Dibenz(a,h)anthracene	53-70-3	2/ 2	0.14	0.36	0.25		Yes	Detected organic	0.022	0.065	0.477	0.022	RFA	Yes	Exceeds screening level	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Dibenzofuran	132-64-9	2/ 2	0.0064	0.051	0.0287		Yes	Detected organic	119	15.3	1192	15.3	RFC	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Diethyl phthalate	84-66-2	1/ 2	0.0093	0.0093	0.0214		Yes	Detected organic	-			4900	RSL	No	Below risk screening criteria	PWss-CONT1	9/30/2003	PWss-CONT1-0001-SO
Di-n-butyl phthalate	84-74-2	1/ 2	0.041	0.041	0.0655		Yes	Detected organic				610	RSL	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Fluoranthene	206-44-0	2/ 2	0.67	2.9	1.79		Yes	Detected organic	276	163	5087	163	RFC	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Fluorene	86-73-7	2/ 2	0.011	0.1	0.0555		Yes	Detected organic	737	243	11458	243	RFC	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Indeno(1,2,3-cd)pyrene	193-39-5	2/ 2	0.31	0.7	0.505		Yes	Detected organic	0.221	0.65	4.77	0.221	RFA	Yes	Exceeds screening level	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Naphthalene	91-20-3	1/ 2	0.039	0.039	0.0283		Yes	Detected organic	368	122	1541	122	RFC	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Phenanthrene	85-01-8	2/ 2	0.16	1.1	0.63		Yes	Detected organic				170	RSL	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO
Pyrene	129-00-0	2/ 2	0.62	2	1.31		Yes	Detected organic	207	122	3815	122	RFC	No	Below risk screening criteria	PWss-CONT2	10/28/2003	PWss-CONT2-0001-SO

^aBackground criteria for soil 0-1 ft bgs from final facility-wide background values for RVAAP, published in the Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio (USACE 2001).

RDA = Concentration associated with recommended daily allowance of essential nutrient

RFA = FWCUG for Resident Farmer Adult RFC = FWCUG for Resident Farmer Child

RSL = United States Environmental Protection Agency Residential Regional Screening Level

AOC = Area of Concern ISM = Incremental Sampling Method bgs = Below ground surface PCB = Polychlorinated Biphenyl CAS = Chemical Abstract Service SRC = Site-related Contaminant COPC = Chemical of Potential Concern -- no value available

HQ = Hazard Quotient -- = no value available
Bold = chemical is a COPC

^bFacility-Wide Cleanup Goals (FWCUGs) for Resident Farmer Adult (RFA), Resident Farmer Child (RFC), and National Guard Trainee (NGT) from Facility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition Plant (USACE 2010).
^cScreening Level Source:

Table B-4. SRC and COPC Screening for Surface Water at Paris-Windham Dump AOC

										Screening FWCUG ^b (HQ= 0.1 or Risk=1E-6)		Risk	Screening				Date Collected
	CAS	Freq of	Minimum		Average	Background	SRC?		DE.	DEG	NOT	Screening	Level	COPC?		Station at	at Max
Analyte (mg/L)	Number	Detect	Detect	Detect	Result	Criteria ^a	(yes/no)	SRC Justification	RFA	RFC	NGT	Level	Source	(yes/no)	COPC Justification	Max Detect	Detect
Inorgani																	
Aluminum	7429-90-5	7/ 7	0.042	0.28	0.104	3.37	No	Below background	63.895	14.827	73.445	14.827	RFC	No	Below background	PWsw-004	04/29/03
Arsenic	7440-38-2	7, 7	0.0028	0.0082	0.00549	0.0032	Yes	Exceeds background	0.0011	0.0012	0.0042	0.0011	RFA	Yes	Exceeds screening level	PWsw-005	04/29/03
Barium	7440-39-3	7/ 7	0.035	0.12	0.066	0.0475	Yes	Exceeds background	12.131	2.901	10.64	2.901	RFC	No	Below risk screening criteria	PWsw-004	04/29/03
Calcium	7440-70-2	7/ 7	23	60	41.4	41.4	No	Essential Nutrient	-	-	-	500	RDA	No	Essential Nutrient	PWsw-001	04/29/03
Cobalt	7440-48-4	7, 7	0.001	0.0015	0.00177	0	Yes	Exceeds background	-	-	-	0.0011	RSL	Yes	Exceeds screening level	PWsw-006	04/29/03
Copper	7440-50-8	7/ 7	0.0022	0.0039	0.0025	0.0079	No	Below background	2.788	0.614	7.199	0.614	RFC	No	Below background	PWsw-004	04/29/03
Iron	7439-89-6	7/ 7	3.6	9.4	5.04	2.56	No	Essential Nutrient	20	4.527	31.296	18	RDA	No	Essential Nutrient	PWsw-005	04/29/03
Lead	7439-92-1	7	0.0019	0.0027	0.00137	0	Yes	Exceeds background	-	-		0.015	RSL	No	Below risk screening criteria	PWsw-004	04/29/03
Magnesium	7439-95-4	7/ 7	6	12	9.4	10.8	No	Essential Nutrient	-	-	-	200	RDA	No	Essential Nutrient	PWsw-004	04/29/03
Manganese	7439-96-5	7/ 7	0.26	0.56	0.379	0.391	Yes	Exceeds background	2.476	0.633	1.449	0.633	RFC	No	Below risk screening criteria	PWsw-006	04/29/03
Mercury	7439-97-6	6/ 7	0.000072	0.0001	0.0000896	0	Yes	Exceeds background	0.0182	0.0044	0.016	0.00435	RFC	No	Below risk screening criteria	PWsw-005	04/29/03
Nickel	7440-02-0	4/ 7	0.002	0.0075	0.00487	0	Yes	Exceeds background	1.445	0.312	8.258	0.312	RFC	No	Below risk screening criteria	PWsw-006	04/29/03
Potassium	7440-09-7	7/ 7	1.7	5.4	4.17	3.17	No	Essential Nutrient	-	-		1750	RDA	No	Essential Nutrient	PWsw-004	04/29/03
Sodium	7440-23-5	7/ 7	4.2	9.9	7.76	21.3	No	Essential Nutrient				1200	RDA	No	Essential Nutrient	PWsw-004	04/29/03
Zinc	7440-66-6	1/ 7	0.013	0.024	0.0149	0.042	No	Below background	21.002	4.617	58.216	4.617	RFC	No	Below background	PWsw-004	04/29/03
		-1/						Explosi	ves						-		
Nitrocellulose	9004-70-0	1/ 1	0.094	0.094	0.094		Yes	Exceeds background				11000	RSL	No	Below risk screening criteria	PWsw-004	04/29/03

^aBackground criteria for surface water from final facility-wide background values for RVAAP, published in the Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio (USACE 2001).

RDA = Concentration associated with recommended daily allowance of essential nutrient

 $RFA = FWCUG \ for \ Resident \ Farmer \ Adult$

RFC = FWCUG for Resident Farmer Child

RSL = United States Environmental Protection Agency Residential Regional Screening Level

AOC = Area of Concern

CAS = Chemical Abstract Service

COPC = Chemical of Potential Concern

HQ = Hazard Quotient

SRC = Site-related Contaminant

-- = no value available

Bold = chemical is a COPC

^bFacility-Wide Cleanup Goals (FWCUGs) for Resident Farmer Adult (RFA), Resident Farmer Child (RFC), and National Guard Trainee (NGT) from Facility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition Plant (USACE 2010).

^cScreening Level Source:

Table B-5. COC Screening for Surface Soil (0-1 ft bgs Discrete Samples) at Paris-Windham Dump AOC: Surface Area EU Representative Receptor: Range Maintenance Soldier

									Range M	Iaintenance				
	CAS	Freq of	Minimum	Maximum	Average	UCL			Soldier	FWCUG ^a	Background	COC?		
Analyte (mg/kg)	Number	Detect	Detect	Detect	Result	95	Dist.	EPC	HQ=1	Risk=1E-5	Criteria ^b	(yes/no)	COC Justification	Ratio
						Inorg	anic Cl	hemicals						
Aluminum	7429-90-5	13/ 13	5300	18000	8350	9900	X	9900	1000000		17700	No	EPC below FWCUG	NA
Manganese	7439-96-5	13/ 13	95	1900	386	807	L	807	204672		1450	No	EPC below FWCUG	NA
					Sem	i-volatil	e Orgai	nic Compou	nds					
Benz(a)anthracene	56-55-3	1/ 1	0.25	0.25	0.25	-	D	0.25		26.2		No	EPC below FWCUG	1.E-02
Benzo(a)pyrene	50-32-8	1/ 1	0.33	0.33	0.33	-	D	0.33		2.62		No	EPC below FWCUG	1.E-01
Benzo(b)fluoranthene	205-99-2	1/ 1	0.39	0.39	0.39		D	0.39		26.2		No	EPC below FWCUG	1.E-02
Sum-of-Ratios for Ca	arcinogens	1/	•	•	•	•	-	•	• •	•		-	-	2.E-01

^aFacility-Wide Cleanup Goals (FWCUGs) for Range Maintenance Soldier from Facility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition Plant (USACE 2010).

UCL 95 = 95% upper confidence limit of the mean

Distribution Code:

D = Fewer than 5 or 50% detects, t statistic used for UCL 95 calculation

L = Lognormal distribution, Land statistic used for UCL 95 calculation

X = Distribution neither normal nor lognormal, t statistic used for UCL 95 calculation

AOC = Area of Concern

bgs = below ground surface

CAS = Chemical Abstract Service

COC = Chemical of Concern

EPC = Exposure Point Concentration

EU = Exposure Unit

HQ = Hazard Quotient

^bBackground criteria for soil 0-1 ft bgs from final facility-wide background values for RVAAP, published in the *Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio* (USACE 2001).

Table B-6. COC Screening for Subsurface Soil (>2 ft bgs Discrete Samples) at Paris-Windham Dump AOC: Fill Area EU Representative Receptor: Range Maintenance Soldier

	CAS	Evan of	Minimum	Maximum	Avonogo				U	aintenance FWCUG ^b	Background	COC?		
Analyte (mg/kg)	Number	Detect		Detect	Result	UCL 95	Dist.	EPC	HQ=1	Risk=1E-5	Criteria ^a	(yes/no)	COC Justification	Ratio
					Se	emi-volatile	Organi	c Compo	unds					
Benz(a)anthracene	56-55-3	1/ 1	1	1	1		D	1		26.2		Yes	EPC below FWCUG	4E-02
Benzo(a)pyrene	50-32-8	1/ 1	1.3	1.3	1.3		D	1.3		2.62		Yes	EPC below FWCUG	5E-01
Benzo(b)fluoranthene	205-99-2	1/ 1	1.2	1.2	1.2		D	1.2		26.2		Yes	EPC below FWCUG	5E-02
Dibenz(a,h)anthracene	53-70-3	1/ 1	0.24	0.24	0.24		D	0.24		2.62		Yes	EPC below FWCUG	9E-02
Indeno(1,2,3-cd)pyrene	193-39-5	1/ 1	0.75	0.75	0.75		D	0.75		26.2		Yes	EPC below FWCUG	3E-02
	•	1,				Pesi	ticides/I	PCBs						
PCB-1254	11097-69-1	1/ 1	0.23	0.23	0.23		D	0.23	36.7	25.7		Yes	EPC below FWCUG	9E-03
Sum-of-Ratios for Carci	nogens	1/		•				•						7E-01

^aFacility-Wide Cleanup Goals (FWCUGs) for Range Maintenance Soldier from Facility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition Plant (USACE 2010).

UCL 95 = 95% upper confidence limit of the mean

Distribution Code:

D = Fewer than 5 or 50% detects, t statistic used for UCL 95 calculation

L = Lognormal distribution, Land statistic used for UCL 95 calculation

X = Distribution neither normal nor lognormal, t statistic used for UCL 95 calculation

AOC = Area of Concern

bgs = below ground surface

CAS = Chemical Abstract Service

COC = Chemical of Concern

EPC = Exposure Point Concentration

EU = Exposure Unit

HQ = Hazard Quotient

 $PCB = Polychlorinated \ Biphenyl$

^bBackground criteria for soil >1 ft bgs from final facility-wide background values for RVAAP, published in the *Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio* (USACE 2001).

Table B-7. COC Screening for Soil at Paris-Windham Dump AOC: ISM Samples
Representative Receptor: Range Maintenance Soldier

		Analyte (mg/kg)	Benz(a)anthr	acene	Benzo(a)pyr	rene	Benzo(b)fluora	nthene	Dibenz(a,h)anth	racene	Indeno(1,2,3-cd)pyrene	
		CAS Number	56-55-3		50-32-8		205-99-2	2	53-70-3		193-39-5	5	
		Range Maintenance So	ldier FWCUG ^a :										
		HQ=1			1		-						
		Risk=1E-5	262		2 (2	2.0			2 (2		26.2	26.2	
		Background Criteria ^b	26.2 		2.62 		26.2 		2.62				
Sample ID	Date	Station	Result/COC?	Ratio	Result/COC?	Ratio	Result/COC?	Ratio	Result/COC?	Ratio	Result/COC?	Ratio	SOR
PWss-CONT1-0001-SO	9/30/2003	PWss-CONT1	No	1E-02	No No	2E-01	No	2E-02	No No	5E-02	No No	1E-02	3E-01
PWss-CONT2-0001-SO	10/28/2003	PWss-CONT2	0.36 No	4E-02	0.40 No	5E-01	0.5 No	5E-02	0.14 No	1E-01	0.51 No	3E-02	8E-01
		-			1.4		1.4		0.30		0.7		

1

Ratio = Sample concentration/FWCUG

AOC = Area of Concern

bgs = below ground surface

CAS = Chemical Abstract Service

COC = Chemical of Concern

HQ = Hazard Quotient

ISM = Incremental Sampling Method

SOR = Sum-of-Ratios

^aFacility-Wide Cleanup Goals (FWCUGs) for Range Maintenance Soldier from Facility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition Plant (USACE 2010).

^bBackground criteria for soil 0-1 ft bgs from final facility-wide background values for RVAAP, published in the Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio (USACE 2001).

Table B-8. COC Screening for Surface Soil (0-1 ft bgs Discrete Samples) at Paris-Windham Dump AOC: Surface Area EU Representative Receptor: Trespasser

									Trespasser FWCUG ^a							
Analyte (mg/kg)	CAS Number	Freq of Detect	Minimum Detect	Maximum Detect	8	UCL 95	Dist.	EPC	HQ=1	Source	Risk=1E-5	Source	L.	coc.	COC Justification	Ratio
							Inorgo	anic Chem	icals							
Aluminum	7429-90-5	13/ 13	5300	18000	8350	9900	X	9900	1000000	TJ/TA			17700	No	EPC below FWCUG	NA
Manganese	7439-96-5	13/ 13	95	1900	386	807	L	807	220293	TA			1450	No	EPC below FWCUG	NA
						Semi-v	olatile	Organic (Compoun	ds						
Benz(a)anthracene	56-55-3	1/ 1	0.25	0.25	0.25		D	0.25			11.3	TA		No	EPC below FWCUG	2.E-02
Benzo(a)pyrene	50-32-8	1/ 1	0.33	0.33	0.33		D	0.33			1.13	TA		No	EPC below FWCUG	3.E-01
Benzo(b)fluoranthene	205-99-2	1/ 1	0.39	0.39	0.39		D	0.39			11.3	TA		No	EPC below FWCUG	3.E-02
Sum-of-Ratios for Carc	inogens	1/				•					•					3.E-01

^aFacility-Wide Cleanup Goals (FWCUGs) for Adult Trespasser (TA) and Juvenile Trespasser (TJ) from Facility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition Plant (USACE 2010). FWCUG presented is the most conservative (smallest) of the Adult and Juvenile value.

UCL 95 = 95% upper confidence limit of the mean

Distribution Code:

D = Fewer than 5 or 50% detects, t statistic used for UCL 95 calculation

L = Lognormal distribution, Land statistic used for UCL 95 calculation

X = Distribution neither normal nor lognormal, t statistic used for UCL 95 calculation

AOC = Area of Concern

bgs = below ground surface

CAS = Chemical Abstract Service

COC = Chemical of Concern

EPC = Exposure Point Concentration

EU = Exposure Unit

HQ = Hazard Quotient

^bBackground criteria for soil 0-1 ft bgs from final facility-wide background values for RVAAP, published in the *Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio* (USACE 2001).

Table B-9. COC Screening for Soil at Paris-Windham Dump AOC: ISM Samples Representative Receptor: Trespasser

		Analyte (mg/kg)	Benz(a)anthra	cene	Benzo(a)py	rene	Benzo(b)fluora	nthene	Dibenz(a,h)anth	racene	Indeno(1,2,3-cd)	pyrene	
		CAS Number	Result Coc.		50-32-8		205-99-2	205-99-2			193-39-5		l
		Trespasser FWCUG ^a :											
		HQ=1									-		l
		Risk=1E-5	11.3 TA	11.3 TA		1 13 TA			1 13 TA		11.3 TA		l
		Background Criteria ^b											
Sample ID	Date	Station	Result/COC?	Ratio	Result/COC?	Ratio	Result/COC?	Ratio	Result/COC?	Ratio	Result/COC?	Ratio	SOR
PWss-CONT2-0001-SO	10/28/2003	PWss-CONT2	No	9.E-02	Yes	NA	No	1.E-01		3.E-01	No	6.E-02	6.E-01
					14		1 4		0.30		0.7		

^aFacility-Wide Cleanup Goals (FWCUGs) for Adult Trespasser (TA) and Juvenile Trespasser (TJ) from Facility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition Plant (USACE 2010). FWCUG presented is the most conservative (smallest) of the Adult and Juvenile value.

Ratio = Sample concentration/FWCUG

AOC = Area of Concern

bgs = below ground surface

CAS = Chemical Abstract Service

COC = Chemical of Concern

HQ = Hazard Quotient

ISM = Incremental Sampling Method

NA = Not applicable, sample concentration exceeds FWCUG; therefore, not included in SOR for identifying additional COCs with concentrations below FWCUGs.

SOR = Sum-of-ratios

TA = Lowest FWCUG is for the Adult Trespasser

-- = no value available

Bold = Concentration exceeds FWCUG

^bBackground criteria for soil 0-1 ft bgs from final facility-wide background values for RVAAP, published in the Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio (USACE 2001).

Table B-10. COC Screening for Surface Soil (0-1 ft bgs Discrete Samples) at Paris-Windham Dump AOC: Surface Area EU
Baseline Receptor: Resident Farmer

									Re	sident Fa	armer FWC	UG ^a				
Amaluta (ma/ka)	CAS Number	Freq of Detect	Minimum Detect	Maximum Detect	U	UCL 95	Diat.	EPC	HO-1	Source	Risk=1E-5	Source	Background Criteria ^b	COC?	COC Justification	Ratio
Analyte (mg/kg)	rumber	Detect	Detect	Detect	Kesuit	UCL 93		rganic Che	,	Source	-112 C	Source	Crneria	(yes/no)	COC Justification	Natio
							11101	gunic Che	micuis							
Aluminum	7429-90-5	13/ 13	5300	18000	8350	9900	X	9900	73798	RFC			17700	No	EPC below FWCUG	NA
Manganese	7439-96-5	13/ 13	95	1900	386	807	L	807	2927	RFC			1450	No	EPC below FWCUG	NA
						Sem	i-volat	ile Organi	c Compo	unds						
Benz(a)anthracene	56-55-3	1/ 1	0.25	0.25	0.25		D	0.25			2.21	RFA		No	EPC below FWCUG	1.E-01
Benzo(a)pyrene	50-32-8	1/ 1	0.33	0.33	0.33		D	0.33			0.221	RFA		Yes	EPC exceeds FWCUG	NA
Benzo(b)fluoranthene	205-99-2	1/ 1	0.39	0.39	0.39		D	0.39			2.21	RFA		No	EPC below FWCUG	2.E-01
Sum-of-Ratios for Car	cinogens	1/														3.E-01

^aFacility-Wide Cleanup Goals (FWCUGs) for Resident Farmer Adult (RFA) and Resident Farmer Child (RFC) from Facility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition Plant (USACE 2010).

UCL 95 = 95% upper confidence limit of the mean

Distribution Code:

D = Fewer than 5 or 50% detects, t statistic used for UCL 95 calculation

L = Lognormal distribution, Land statistic used for UCL 95 calculation

X = Distribution neither normal nor lognormal, t statistic used for UCL 95 calculation

AOC - Area of Concern

bgs = below ground

CAS = Chemical Abstract Service

COC = Chemical of Concern

EPC = Exposure Point Concentration

EU = Exposure Unit

HQ = Hazard Quotient

NA = Not applicable, sample concentration exceeds FWCUG; therefore, not included in SOR for identifying additional COCs with concentrations below FWCUGs.

SOR = Sum-of-ratios

-- = no value available

Bold = EPC exceeds FWCUG

^bBackground criteria for soil 0-1 ft bgs from final facility-wide background values for RVAAP, published in the Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio (USACE 2001).

Table B-11. COC Screening for Soil at Paris-Windham Dump AOC: ISM Samples Baseline Receptor: Resident Farmer

		Analyte (mg/kg)	Benz(a)anthr	acene	Benzo(a)pyr	ene	Benzo(b)fluorai	nthene	Dibenz(a,h)antl	hracene	Indeno(1,2,3-cd))pyrene		
		CAS Number	S Number 56-55-3		50-32-8	50-32-8 205-99-2			53-70-3		193-39-5	i	i	
		Resident Farmer FWC	lent Farmer FWCUG ^a :											
		HQ=1											l	
		Risk=1E-5	2.21 RF	A	0.221 RF	A	2.21 RFA		0.221 RF	Ά	2.21 RF	4	i	
		Background Criteria ^b	-										ĺ	
Sample ID	Date	Station	Result/COC?	Ratio	Result/COC?	Ratio	Result/COC?	Ratio	Result/COC?	Ratio	Result/COC?	Ratio	SOR	
PWss-CONT2-0001-SO	10/28/2003	PWss-CONT2	No	5.E-01	1.4 Yes	NA	1.4 No	6.E-01	0.36 Yes	NA	0.7 No	3.E-01	1.E+0	

1

Ratio = Sample concentration/FWCUG

AOC = Area of Concern

bgs = below ground surface

CAS = Chemical Abstract Service

COC = Chemical of Concern

HQ = Hazard Quotient

ISM = Incremental Sampling Method

NA = Not applicable, sample concentration exceeds FWCUG; therefore, not included in SOR for identifying additional COCs with concentrations below FWCUGs.

SOR - Sum-of-ratios

RFA = Lowest FWCUG is for the Adult Resident Farmer

^aFacility-Wide Cleanup Goals (FWCUGs) for Resident Farmer Adult (RFA) and Resident Farmer Child (RFC) from Facility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition Plant (USACE 2010).

^bBackground criteria are for soil 0-1 ft bgs from final facility-wide background values for RVAAP, published in the *Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio* (USACE 2001).

Table B-12. COC Screening for Surface Water at Paris-Windham Dump AOC Representative Receptor: Trespasser

									Trespasser FWCUG ^a						
	CAS		Minimum	Maximum	Average	UCL			HO=1	Source	Risk=1E-5	Course	Background	COC?	0007 400 4
Analyte (mg/L)	Number	Detect	Detect	Detect	Result	95	Dist.	EPC	nQ=1	Source	KISK=1E-5	Source	Criteria ^b	(yes/no)	COC Justification
							Ino	rganic Chen	iicals						
Arsenic	7440-38-2	7/ 7	0.0028	0.0082	0.00549	0.0069	N	0.00685	0.705	TJ	0.0415	TA	0.0032	No	EPC below FWCUG
Cobalt	7440-48-4	" 7	0.001	0.0015	0.00177	0.0027	т	0.0015					0	No ^c	No FWCUG available

^aFacility-Wide Cleanup Goals (FWCUGs) for Adult Trespasser (TA) and Juvenile Trespasser (TJ) from Facility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition Plant (USACE 2010). FWCUG presented is the most conservative (smallest) of the Adult and Juvenile value.

^cNo FWCUG is available for cobalt in surface water. Maximum detected concentration (0.0015 mg/L) barely exceeds the United States Environmental Protection Agency residential Regional Screening Level (0.0011 mg/L) at an HQ of 0. Distribution Code:

D = Fewer than 5 or 50% detects, t statistic used for UCL 95 calculation

L = Lognormal distribution, Land statistic used for UCL 95 calculation

N = Normal distribution, t statistic used for UCL 95 calculation

UCL 95 = 95% upper confidence limit of the mean

AOC = Area of Concern

CAS = Chemical Abstract Service

COC = Chemical of Concern

EPC = Exposure Point Concentration

HQ = Hazard Quotient

-- = no value available

^bBackground criteria for surface water from final facility-wide background values for RVAAP, published in the Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio (USACE 2001).

Table B-13. COC Screening for Surface Water at Paris-Windham Dump AOC Baseline Receptor: Resident Farmer

		Freq							Resident Farmer FWCUG ^a						
Analyte (mg/kg)	CAS Number	of Detect		Maximum Detect	Average Result		Dist.	EPC	HQ=1	Source	Risk=1E-5		Background Criteria ^b	COC? (yes/no)	
Inorganic Chemicals															
1							11101	Same Chen	nicuis						
Arsenic	7440-38-2	7/ 7	0.0028	0.0082	0.00549	0.0069		0	0.0463	RFC	0.011	RFA	0.0032	No	EPC below FWCUG

^aFacility-Wide Cleanup Goals (FWCUGs) for Resident Farmer Adult (RFA) and Resident Farmer Child (RFC) from *Facility-Wide Human Health Cleanup Goals for the Ravenna Army Ammunition Plant* (USACE 2010).

 c No FWCUG is available for cobalt in surface water. Maximum detected concentration (0.0015 mg/L) barely exceeds the United States Environmental Protection Agency residential Regional Screening Level (0.0011 mg/L) at an HQ of 0.1.

Distribution Code:

D = Fewer than 5 or 50% detects, t statistic used for UCL 95 calculation

L = Lognormal distribution, Land statistic used for UCL 95 calculation

X = Distribution neither normal nor lognormal, t statistic used for UCL 95 calculation

UCL 95 - 95% upper confidence limit of the mean

AOC = Area of Concern

CAS = Chemical Abstract Service

COC = Chemical of Concern

EPC = Exposure Point Concentration


HQ = Hazard Quotient

-- = no value available

^bBackground criteria for surface water from final facility-wide background values for RVAAP, published in the *Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio* (USACE 2001).

APPENDIX C

Ecological Risk Assessment Information and Data



1		TABLE OF CONTENTS	
2			
3		FIGURE	
4	Figure C-1.	. Ohio Rapid Assessment Method Worksheet	1
5			
6			
7		LIST OF TABLES	
8			
9	Table C-1.	Ecological Screening Values for Chemical Analytes in Soil	3
10		Ecological Screening Values for Chemical Analytes in Surface Water	
11	Table C-3.	SRC and COPEC Screening with Maximum Ratio for Shallow Surface Soil (0-1	
12		ft bgs Discrete Samples) at Dump Along Paris-Windham Road: Fill Area EU	28
13	Table C-4.	SRC and COPEC Screening with Maximum Ratio for Shallow Surface Soil (0-1	
14		ft bgs Discrete Samples) at Dump Along Paris-Windham Road: Surface Area EU	29
15	Table C-5.	SRC and COPEC Screening for Surface Water at Dump Along	
16		Paris-Windham Road	30
17	Table C-6.	Checklist of Important Ecological Places and Resources at Dump Along Paris-	
18		Windham Road	31
19	Table C-7.	Natural Resources Management Goals (OHARNG 2008)	33
20			
21			
22	REFEREN	CES	37
23			

ACRONYMS AND ABBREVIATIONS

AOC Area of Concern

1

BRAC Base Realignment and Closure
CAS Chemical Abstract Service

CERCLA Comprehensive Environmental Response, Compensation, and Liability Act

COPC Chemical of Potential Concern

COPEC Chemical of Potential Ecological Concern
DOE United States Department of Energy

DOW Department of Wildlife

EcoSSLEcological Soil Screening LevelEDQLEcological Data Quality LevelsESLEcological Screening LevelESVEcological Screening Value

EU Exposure Unit

GIS Geographic Information System

GLI Great Lakes Initiative

HTRW Hazardous, Toxic, and Radioactive Waste
INRMP Integrated Natural Resources Management Plan
NAWOC National Ambient Water Quality Criteria

OAC Ohio Administrative Code

ODNR Ohio Department of Natural Resources

OHARNG Ohio Army National Guard

Ohio EPA Ohio Environmental Protection Agency

OMZM Outside Mixing Zone Maximum

PBT Persistent, Bioaccumulative, and Toxic

PCB Polychlorinated Biphenyl PLS Planning Level Survey

PRG Preliminary Remediation Goal

Reg Region

RVAAP Ravenna Army Ammunition Plant

SRC Site-related Contaminant

T&E Threatened and Endangered Species
USACE United States Army Corps of Engineers

USEPA United States Environmental Protection Agency

2

ORAM v. 5.0 Field Form Quantitative Rating Paris - Windham Dump AOC
Site: Wetland 1 Rater(s): James Groton Date: 11 30 2011
Metric 1. Wetland Area (size).
Select one size class and assign score. So acres (>20.2ha) (6 pts) So acres (>20.2ha) (5 pts) So acres (>20.2ha) (5 pts) So acres (>20.2ha) (2 pts) So acres (10.1 to <20.2ha) (3 pts) So acres (0.12 to <1.2ha) (3 pts) So acres (0.12 to <1.2ha) (2 pts) So acres (0.04 to <0.12ha) (1 pt) So acres (0.04ha) (0 pts) So
구 용 Metric 2. Upland buffers and surrounding land use.
2a. Calculate average buffer width. Select only one and assign score. Do not double check. WIDE. Buffers average 50m (164ft) or more around welland perimeter (7) MEDIUM. Buffers average 25m to <50m (82 to <164ft) around wetland perimeter (4) NARROW. Buffers average 10m to <25m (32ft to <82ft) around wetland perimeter (1) VERY NARROW. Buffers average <10m (<32ft) around welland perimeter (0) 2b. Intensity of surrounding land use. Select one or double check and average. VERY LOW. 2nd growth or older forest, prairie, savannah, wildlife area, etc. (7) LOW. Old field (>10 years), shrub land, young second growth forest. (5) MODERATELY HIGH. Residential, fenced pasture, park, conservation tillage, new fallow field. (3) HIGH. Urban, industrial, open pasture, row cropping, mining, construction. (1)
15 23 Metric 3. Hydrology.
max 30 pts. sublotal 3a. Sources of Water. Score all that apply. High pH groundwater (5) Other groundwater (3) Precipitation (1) Seasonal/Intermittent surface water (3) Perennial surface water (lake or stream) (5) 3c. Maximum water depth. Select only one and assign score. >0.7 (27.6in) (3) 0.4 to 0.7m (15.7 to 27.6in) (2) -0.4m (<15,7in) (1) 3e. Modifications to natural hydrologic regime. Score one or double check and average. None or none apparent (12) Recovered (7) Recovering (3) Recent or no recovery (1) Recovering (3) Recent or no recovery (1)
フラスク Metric 4. Habitat Alteration and Development.
max 20 pts. substrate disturbance. Score one or double check and average. None or none apparent (4) Recovering (2) Recovering (2) Recent or no recovery (1)
None or none apparent (9) Recovered (6) Recovering (3) Recent or no recovery (1)

Figure C-1. Ohio Rapid Assessment Method Worksheet

ORAM v. 5.0 Field Form Quantitative Rating Paris - W	inlksu	Dump AOC
Site: Wetland 1 Rate	r(s): उล	mes Groton Date: 11/30/2011
Subtotat first page O 30 Metric 5. Special Wetlan		
max 10 pts. subtotal Check all that apply and score as indicated.		
Bog (10) Fen (10) Old growth forest (10) Mature forested wetland (5) Lake Erie coastal/tributary wetland- Lake Erie coastal/tributary wetland- Lake Plain Sand Prairies (Oak Ope Relict Wet Prairies (10) Known occurrence state/federal thri Significant migratory songbird/wate Category 1 Wetland. See Question	-restricted hydro nings) (10) reatened or enda r fowl habitat or n 1 Qualitative R	angered species (10)
7 34 Metric 6. Plant commun	ities, int	erspersion, microtopography.
max 20 pts. subtotal 6a. Wetland Vegetation Communities.		Community Cover Scale
Score all present using 0 to 3 scale. Aguatic bed	0	Absent or comprises <0.1ha (0.2471 acres) contiguous area
Emergent	'	Present and either comprises small part of wetland's vegetation and is of moderate quality, or comprises a
Shrub		significant part but is of low quality
2 Forest	2	Present and either comprises significant part of wetland's
Mudflats		vegetation and is of moderate quality or comprises a small
Open water		part and is of high quality
Other	3	Present and comprises significant part, or more, of wetland's
6b. horizontal (plan view) Interspersion.		vegetation and is of high quality
Select only one.		
High (5)		escription of Vegetation Quality
Moderately high(4)	low	Low spp diversity and/or predominance of nonnative or
Moderate (3)		disturbance tolerant native species
Moderately low (2) Low (1)	mod	Native spp are dominant component of the vegetation,
None (0)		although nonnative and/or disturbance tolerant native spp
6c. Coverage of invasive plants. Refer		can also be present, and species diversity moderate to moderately high, but generally w/o presence of rare
to Table 1 ORAM long form for list. Add		threatened or endangered spp
or deduct points for coverage	high	A predominance of native species, with nonnative spp
Extensive >75% cover (-5)		and/or disturbance tolerant native spp absent or virtually
Moderate 25-75% cover (-3)		absent, and high spp diversity and often, but not always,
Sparse 5-25% cover (-1)		the presence of rare, threatened, or endangered spp
Nearly absent <5% cover (0)		
Absent (1)		Open Water Class Quality
6d. Microtopography.	0	Absent <0.1ha (0.247 acres)
Score all present using 0 to 3 scale.	1	Low 0.1 to <1ha (0.247 to 2.47 acres)
Vegetated hummucks/tussucks Coarse woody debris >15cm (6in)	2 3	Moderate 1 to <4ha (2.47 to 9.88 acres)
Standing dead >25cm (10in) dbh		High 4ha (9.88 acres) or more
Amphibian breeding pools	Microtopogr	raphy Cover Scale
	0	Absent
-	1	Present very small amounts or if more common of marginal quality
	2	Present in moderate amounts, but not of highest
\cap \cap \cap	3	quality or in small amounts of highest quality
	3	Present in moderate or greater amounts
37 Category L		and of highest quality
End of Quantitative Rating.	Complete	Categorization Worksheets.

Figure C-1. Ohio Rapid Assessment Method Worksheet (continued)

1

			Soil Screening Values									
		USEPA	DOE (1997a) Preliminary Remediation Goals for Ecological Endpoints ^a		Screenin	gion 5 Ecological g Levels (2003) f 1998 EDQLs)	Preferred Ecological Screening Value (ESV) ^d					
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source			
			Inorganic C	hemicals								
Aluminum	7429-90-5	*	Al EcoSSL	50	PRGs ^b			50	PRGs			
Antimony	7440-36-0	0.27	mammalian EcoSSL for Sb	5	PRGs	0.142	USEPA Reg 5	2.70E-01	mammalian EcoSSL for Sb			
Arsenic	7440-38-2	18	plant EcoSSL for As	9.9	PRGs	5.7	USEPA Reg 5	1.80E+01	plant EcoSSL for As			
Barium	7440-39-3	330	soil invert EcoSSL for Ba	283	PRGs	1.04	USEPA Reg 5	3.30E+02	soil invert EcoSSL for Ba			
Beryllium	7440-41-7	21	mammalian EcoSSL for Be	10	PRGs	1.06	USEPA Reg 5	2.10E+01	mammalian EcoSSL for Be			
Bismuth	7440-69-9							No ESV	No Source			
Boron	7440-42-8			0.5	PRGs			5.00E-01	PRGs			
Bromine	7726-95-6			10	PRGs			1.00E+01	PRGs			
Cadmium	7440-43-9	0.36	mammalian EcoSSL for Cd	4	PRGs	0.00222	USEPA Reg 5	3.60E-01	mammalian EcoSSL for Cd			
Calcium	7440-70-2							No ESV	No Source			
Chromium	16065-83-1	26	avian EcoSSL for Cr III	0.4	PRGs	0.4	ESL for Cr+3	2.60E+01	avian EcoSSL for Cr III			
Chromium, hexavalent	18540-29-9	130	mammalian EcoSSL for Cr VI					1.30E+02	mammalian EcoSSL for Cr VI			
Cobalt	7440-48-4	13	plant EcoSSL for Co	20	PRGs	0.14	USEPA Reg 5	1.30E+01	plant EcoSSL for Co			
Copper	7440-50-8	28	avian EcoSSL for Cu	60	PRGs	5.4	USEPA Reg 5	2.80E+01	avian EcoSSL for Cu			

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

		Soil Screening Values									
		USEPA	DOE (1997a) Preliminary Remediation Goals for Ecological Endpoints ^a		USEPA Re	egion 5 Ecological g Levels (2003) of 1998 EDQLs)	Preferred Ecological Screening Value (ESV) ^d				
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source		
Cyanide	57-12-5					1.33	USEPA Reg 5	1.33E+00	USEPA Reg 5		
Fluorine	7782-41-4			200	PRGs			2.00E+02	PRGs		
Iodine	7553-56-2			4	PRGs			4.00E+00	PRGs		
Iron	7439-89-6	**	Fe EcoSSL					No ESV	No Source		
Lanthanum	7439-91-0							No ESV	No Source		
Lead	7439-92-1	11	avian EcoSSL for Pb	40.5	PRGs	0.0537	USEPA Reg 5	1.10E+01	avian EcoSSL for Pb		
Lithium	7439-93-2			2	PRGs			2.00E+00	PRGs		
Magnesium	7439-95-4							No ESV	No Source		
Manganese	7439-96-5	220	plant EcoSSL for Mn	500	PRGs ^b			2.20E+02	plant EcoSSL for Mn		
Mercury	7439-97-6			0.00051	PRGs	0.1	USEPA Reg 5	5.10E-04	PRGs		
Mercury, methyl	22967-92-6					0.00158	USEPA Reg 5	1.58E-03	USEPA Reg 5		
Molybdenum	7439-98-7			2	PRGs			2.00E+00	PRGs		
Nickel	7440-02-0	38	plant EcoSSL for Ni	30	PRGs	13.6	USEPA Reg 5	3.80E+01	plant EcoSSL for Ni		
Potassium	7440-09-7							No ESV	No Source		
Selenium	7782-49-2	0.52	plant EcoSSL for Se	0.21	PRGs	0.0276	USEPA Reg 5	5.20E-01	plant EcoSSL for Se		
Silver	7440-22-4	4.2	avian EcoSSL for Ag	2	PRGs	4.04	USEPA Reg 5	4.20E+00	avian EcoSSL for Ag		
Sodium	7440-23-5							No ESV	No Source		
Technetium	7440-26-8			0.2	PRGs			2.00E-01	PRGs		
Tellurium	13494-80-9							No ESV	No Source		

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

				1	Soil Screen	ning Values		T	
		USEPA	DOE (1997a) Preliminary Remediation Goals for Ecological Endpoints ^a		Screenin	egion 5 Ecological g Levels (2003) of 1998 EDQLs)	Preferred Ecological Screening Value (ESV) ^d		
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source
Thallium	7440-28-0			1	PRGs	0.0569	USEPA Reg 5	1.00E+00	PRGs
Tin	7440-31-5			50	PRGs	7.62	USEPA Reg 5	5.00E+01	PRGs
Titanium	7440-32-6							No ESV	No Source
Tungsten	7440-33-7							No ESV	No Source
Uranium	7440-61-1			5	PRGs			5.00E+00	PRGs
Vanadium	7440-62-2	7.8	avian EcoSSL for V	2	PRGs	1.59	USEPA Reg 5	7.80E+00	avian EcoSSL for V
Zinc	7440-66-6	46	avian EcoSSL for Zn	8.5	PRGs	6.62	USEPA Reg 5	4.60E+01	avian EcoSSL for Zn
			Anio	ns		_			
Nitrate	14797-55-8							No ESV	No Source
Sulfide	18496-25-8					0.00358	USEPA Reg 5	3.58E-03	USEPA Reg 5
			Organic Cl	hemicals		•			
Acenaphthene	83-32-9			20	PRGs	682	USEPA Reg 5	2.00E+01	PRGs
Acenaphthylene	208-96-8					682	USEPA Reg 5	6.82E+02	USEPA Reg 5
Acetone	67-64-1					2.5	USEPA Reg 5	2.50E+00	USEPA Reg 5
Acetonitrile	75-05-8					1.37	USEPA Reg 5	1.37E+00	USEPA Reg 5
Acetophenone	98-86-2					300	USEPA Reg 5	3.00E+02	USEPA Reg 5
Acetylaminofluorene[2-]	53-96-3					0.596	USEPA Reg 5	5.96E-01	USEPA Reg 5
Acrolein	107-02-8					5.27	USEPA Reg 5	5.27E+00	USEPA Reg 5
Acrylonitrile	107-13-1					0.0239	USEPA Reg 5	2.39E-02	USEPA Reg 5

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

					Soil Screen	ning Values			
		USEPA	A EcoSSLs	DOE (1997a) Preliminary Remediation Goals for Ecological Endpoints ^a		USEPA Region 5 Ecological Screening Levels (2003) (update of 1998 EDQLs)		Preferred Ecological Screening Value (ESV) ^d	
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source
Aldrin	309-00-2					0.00332	USEPA Reg 5	3.32E-03	USEPA Reg 5
2-Amino-4,6-dinitrotoluene	35572-78-2							No ESV	No Source
4-Amino-2,6-dinitrotoluene	19406-51-0							No ESV	No Source
4-Aminobiphenyl	92-67-1					0.00305	USEPA Reg 5	3.05E-03	USEPA Reg 5
Aniline	62-53-3					0.0568	USEPA Reg 5	5.68E-02	USEPA Reg 5
Anthracene	120-12-7					1480	USEPA Reg 5	1.48E+03	USEPA Reg 5
Aramite	140-57-8					166	USEPA Reg 5	1.66E+02	USEPA Reg 5
Azobenzene[p- (dimethylamino)]	60-11-7					0.04	USEPA Reg 5	4.00E-02	USEPA Reg 5
PCB-1016	12674-11-2							No ESV	No Source
Arochlor-1221	11104-28-2							No ESV	No Source
Arochlor-1232	11141-16-5							No ESV	No Source
Arochlor-1242	53469-21-9							No ESV	No Source
Arochlor-1248	12672-29-6							No ESV	No Source
PCB-1254	11097-69-1							No ESV	No Source
PCB-1260	11096-82-5							No ESV	No Source
Benzene	71-43-2					0.255	USEPA Reg 5	2.55E-01	USEPA Reg 5
Benzenemethanol	100-51-6					65.8	USEPA Reg 5	6.58E+01	USEPA Reg 5
Benz(a)anthracene	56-55-3					5.21	USEPA Reg 5	5.21E+00	USEPA Reg 5
Benzo(a)pyrene	50-32-8					1.52	USEPA Reg 5	1.52E+00	USEPA Reg 5

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

					Soil Scree	ning Values			
		USEPA	DOE (1997a) Preliminary Remediation Goals for Ecological Endpoints ^a		USEPA Region 5 Ecological Screening Levels (2003) (update of 1998 EDQLs)		Preferred Ecological Screening Value (ESV) ^d		
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source
Benzo(b)fluoranthene	205-99-2					59.8	USEPA Reg 5	5.98E+01	USEPA Reg 5
Benzo(ghi)perylene	191-24-2					119	USEPA Reg 5	1.19E+02	USEPA Reg 5
Benzo(k)fluoranthene	207-08-9					148	USEPA Reg 5	1.48E+02	USEPA Reg 5
Benzoic acid	65-85-0							No ESV	No Source
ВНС	608-73-1							No ESV	No Source
BHC, alpha	319-84-6					0.0994	USEPA Reg 5	9.94E-02	USEPA Reg 5
BHC, beta	319-85-7					0.00398	USEPA Reg 5	3.98E-03	USEPA Reg 5
BHC, delta	319-86-8					9.94	USEPA Reg 5	9.94E+00	USEPA Reg 5
BHC, gamma (Lindane)	58-89-9					0.005	USEPA Reg 5	5.00E-03	USEPA Reg 5
Biphenyl	92-52-4			60	PRGs			6.00E+01	PRGs
bis(2-chloroethoxy) methane	111-91-1					0.302	USEPA Reg 5	3.02E-01	USEPA Reg 5
bis(2-Chloroethyl) ether	111-44-4					23.7	USEPA Reg 5	2.37E+01	USEPA Reg 5
bis(2-Ethylhexyl)phthalate	117-81-7					0.925	USEPA Reg 5	9.25E-01	USEPA Reg 5
4-Bromoaniline	106-40-1							No ESV	No Source
Bromodichloromethane	75-27-4					0.54	USEPA Reg 5	5.40E-01	USEPA Reg 5
Bromoform	75-25-2					15.9	USEPA Reg 5	1.59E+01	USEPA Reg 5
Bromomethane	74-83-9					0.235	USEPA Reg 5	2.35E-01	USEPA Reg 5
4-bromophenyl-phenylether	101-55-3							No ESV	No Source
2-Butanone	78-93-3					89.6	USEPA Reg 5	8.96E+01	USEPA Reg 5

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

		Soil Screening Values									
		USEPA	DOE (1997a) Preliminary Remediation Goals for Ecological Endpoints ^a		USEPA Re	egion 5 Ecological g Levels (2003) of 1998 EDQLs)	Preferred Ecological Screening Value (ESV) ^d				
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source		
Butylbenzyl phthalate	85-68-7					0.239	USEPA Reg 5	2.39E-01	USEPA Reg 5		
N-Nitrosodi-n-Butylamine	924-16-3					0.267	USEPA Reg 5	2.67E-01	USEPA Reg 5		
Carbazole	86-74-8							No ESV	No Source		
Carbon disulfide	75-15-0					0.0941	USEPA Reg 5	9.41E-02	USEPA Reg 5		
Carbon tetrachloride	56-23-5					2.98	USEPA Reg 5	2.98E+00	USEPA Reg 5		
Chlordane	12789-03-6					0.224	USEPA Reg 5	2.24E-01	USEPA Reg 5		
alpha-Chlordane	12789-03-6					0.224	USEPA Reg 5	2.24E-01	USEPA Reg 5		
gamma-Chlordane	12789-03-6					0.224	USEPA Reg 5	2.24E-01	USEPA Reg 5		
Chloroacetamide	79-07-2			2	PRGs ^c			2.00E+00	PRGs		
3-Chloroaniline	108-42-9			20	PRGs			2.00E+01	PRGs		
4-Chloroaniline	106-47-8					1.1	USEPA Reg 5	1.10E+00	USEPA Reg 5		
Chlorobenzene	108-90-7			40	PRGs	13.1	USEPA Reg 5	4.00E+01	PRGs		
Chlorobenzilate	510-15-6					5.05	USEPA Reg 5	5.05E+00	USEPA Reg 5		
Chloroethane	75-00-3							No ESV	No Source		
Chloroform	67-66-3					1.19	USEPA Reg 5	1.19E+00	USEPA Reg 5		
Chloromethane	74-87-3					10.4	USEPA Reg 5	1.04E+01	USEPA Reg 5		
2-Chloronaphthalene	91-58-7					0.0122	USEPA Reg 5	1.22E-02	USEPA Reg 5		
2-Chlorophenol	95-57-8					0.243	USEPA Reg 5	2.43E-01	USEPA Reg 5		
3-Chlorophenol	108-43-0			7	PRGs			7.00E+00	PRGs		

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

				1	Soil Screen	ning Values			
		USEPA	DOE (1997a) Preliminary Remediation Goals for Ecological Endpoints ^a		USEPA Region 5 Ecological Screening Levels (2003) (update of 1998 EDQLs)		Preferred Ecological Screening Value (ESV) ^d		
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source
4-Chlorophenol	106-48-9							No ESV	No Source
4-Chlorophenyl-phenyl ether	7005-72-3	1					1	No ESV	No Source
4-chloro-3-methylphenol	59-50-7					7.95	USEPA Reg 5	7.95E+00	USEPA Reg 5
Chloropropene	107-05-1					0.0134	USEPA Reg 5	1.34E-02	USEPA Reg 5
Chloroprene	126-99-8	-				0.0029	USEPA Reg 5	2.90E-03	USEPA Reg 5
Chrysene	218-01-9	-				4.73	USEPA Reg 5	4.73E+00	USEPA Reg 5
m-Cresol	108-39-4					3.49	USEPA Reg 5	3.49E+00	USEPA Reg 5
2,4-D	94-75-7					0.0272	USEPA Reg 5	2.72E-02	USEPA Reg 5
4,4'-DDD	72-54-8	0.021	mammalian EcoSSL for DDT and metabolites			0.758	USEPA Reg 5	2.10E-02	mammalian EcoSSL for DDT and metabolites
4,4'-DDE	72-55-9	0.021	mammalian EcoSSL for DDT and metabolites			0.596	USEPA Reg 5	2.10E-02	mammalian EcoSSL for DDT and metabolites
4,4'-DDT	50-29-3	0.021	mammalian EcoSSL for DDT and metabolites			0.0035	USEPA Reg 5	2.10E-02	mammalian EcoSSL for DDT and metabolites
Diallate	2303-16-4					0.452	USEPA Reg 5	4.52E-01	USEPA Reg 5
Diazinon	333-41-5							No ESV	No Source
Dibenz(a,h)anthracene Dibenzofuran	53-70-3 132-64-9					18.4	USEPA Reg 5	1.84E+01 No ESV	USEPA Reg 5 No Source
1,2-Dibromo-3-Chloropropane	96-12-8					0.0352	USEPA Reg 5	3.52E-02	USEPA Reg 5
Dibromochloromethane	124-48-1					2.05	USEPA Reg 5	2.05E+00	USEPA Reg 5

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

		USEPA	Prelim Remediati for Ecol	DOE (1997a) Preliminary Remediation Goals for Ecological Endpoints ^a		egion 5 Ecological g Levels (2003) ff 1998 EDQLs)	Preferred Ecological Screening Value (ESV) ^d		
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source
Dibromoethane	106-93-4					1.23	USEPA Reg 5	1.23E+00	USEPA Reg 5
2,4-Dichloroaniline	554-00-7			100	PRGs ^c			1.00E+02	PRGs
3,4-Dichloroaniline	95-76-1			20	PRGs ^c			2.00E+01	PRGs
1,2-Dichlorobenzene	95-50-1					2.96	USEPA Reg 5	2.96E+00	USEPA Reg 5
1,3-Dichlorobenzene	541-73-1					37.7	USEPA Reg 5	3.77E+01	USEPA Reg 5
1,4-Dichlorobenzene	106-46-7			20	PRGs	0.546	USEPA Reg 5	2.00E+01	PRGs
3,3'-Dichlorobenzidine	91-94-1					0.646	USEPA Reg 5	6.46E-01	USEPA Reg 5
Cis-1,4-dichloro-2-butene	1476-11-5							No ESV	No Source
Trans-1,4-dichloro-2-butene	110-57-6							No ESV	No Source
Dichlorodifluromethane	75-71-8					39.5	USEPA Reg 5	3.95E+01	USEPA Reg 5
1,1-Dichloroethane	75-34-3					20.1	USEPA Reg 5	2.01E+01	USEPA Reg 5
1,2-Dichloroethane	107-06-2					21.2	USEPA Reg 5	2.12E+01	USEPA Reg 5
1,1-Dichloroethene	75-35-4					8.28	USEPA Reg 5	8.28E+00	USEPA Reg 5
1,2-Dichloroethene	540-59-0					0.784	USEPA Reg 5 (for trans form)	7.84E-01	USEPA Reg 5 (for trans form)
2,4-Dichlorophenol	120-83-2					87.5	USEPA Reg 5	8.75E+01	USEPA Reg 5
2,6-Dichlorophenol	87-65-0					1.17	USEPA Reg 5	1.17E+00	USEPA Reg 5
3,4-Dichlorophenol	95-77-2			20	PRGs			2.00E+01	PRGs
1,2-Dichloropropane	78-87-5			700	PRGs ^c	32.7	USEPA Reg 5	7.00E+02	PRGs
cis-1,3-Dichloropropene	10061-01-5					0.398	USEPA Reg 5	3.98E-01	USEPA Reg 5

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

					Soil Scree	ning Values			
		USEPA	DOE (1 Prelim Remediati for Eco Endpo	inary on Goals logical	Screenin	egion 5 Ecological g Levels (2003) ff 1998 EDQLs)	Preferred Ecological Screening Value (ESV) ^d		
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source
trans-1,3-Dichloropropene	10061-02-6					0.398	USEPA Reg 5	3.98E-01	USEPA Reg 5
Dieldrin	60-57-1	0.0049	mammalian EcoSSL for Dieldrin			0.00238	USEPA Reg 5	4.90E-03	mammalian EcoSSL for Dieldrin
O,O-Diethyl O-2- pyrazinylphosphorothioate	297-97-2					0.799	USEPA Reg 5	7.99E-01	USEPA Reg 5
Diethylphthalate	84-66-2			100	PRGs	24.8	USEPA Reg 5	1.00E+02	PRGs
Dimethoate	60-51-5					0.218	USEPA Reg 5	2.18E-01	USEPA Reg 5
Dimethylphthalate	131-11-3			200	PRGs ^c	734	USEPA Reg 5	2.00E+02	PRGs
3,3'-Dimethylbenzidine	119-93-7					0.104	USEPA Reg 5	1.04E-01	USEPA Reg 5
7,12'- Dimethylbenz(a)anthracene	57-97-6					16.3	USEPA Reg 5	1.63E+01	USEPA Reg 5
alpha,alpha- Dimethylphenethylamine	122-09-8					0.3	USEPA Reg 5	3.00E-01	USEPA Reg 5
2,4-Dimethylphenol	105-67-9					0.01	USEPA Reg 5	1.00E-02	USEPA Reg 5
Di-n-butyl phthalate	84-74-2			200	PRGs	0.15	USEPA Reg 5	2.00E+02	PRGs
Di-n-octylphthalate	117-84-0					709	USEPA Reg 5	7.09E+02	USEPA Reg 5
1,3-Dinitrobenzene	99-65-0					0.655	USEPA Reg 5	6.55E-01	USEPA Reg 5
2,4-Dinitrophenol	51-28-5	-		20	PRGs	0.0609	USEPA Reg 5	2.00E+01	PRGs
2,4-Dinitrotoluene	121-14-2					1.28	USEPA Reg 5	1.28E+00	USEPA Reg 5
2,6-Dinitrotoluene	606-20-2					0.0328	USEPA Reg 5	3.28E-02	USEPA Reg 5
4,6-Dinitro-2-methylphenol	534-52-1					0.144	USEPA Reg 5	1.44E-01	USEPA Reg 5
Dinoseb	88-85-7					0.0218	USEPA Reg 5	2.18E-02	USEPA Reg 5

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

		Soil Screening Values								
		USEPA	DOE (1 Prelim Remediati for Eco Endpo	1997a) inary ion Goals logical	USEPA Re	egion 5 Ecological g Levels (2003) of 1998 EDQLs)	Preferred Ecological Screening Value (ESV) ^d			
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	
1,4-Dioxane	123-91-1					2.05	USEPA Reg 5	2.05E+00	USEPA Reg 5	
Diphenylamine	122-39-4					1.01	USEPA Reg 5	1.01E+00	USEPA Reg 5	
Disulfoton	298-04-4					0.0199	USEPA Reg 5	1.99E-02	USEPA Reg 5	
Endosulfan I (alpha)	959-98-8					0.119	USEPA Reg 5	1.19E-01	USEPA Reg 5	
Endosulfan II (beta)	33213-65-9					0.119	USEPA Reg 5	1.19E-01	USEPA Reg 5	
Endosulfan, mixed isomers	115-29-7							No ESV	No Source	
Endosulfan sulfate	1031-07-8					0.0358	USEPA Reg 5	3.58E-02	USEPA Reg 5	
Endrin	72-20-8					0.0101	USEPA Reg 5	1.01E-02	USEPA Reg 5	
Endrin aldehyde	7421-93-4					0.0105	USEPA Reg 5	1.05E-02	USEPA Reg 5	
Ethyl metharcrylate	97-63-2					30	USEPA Reg 5	3.00E+01	USEPA Reg 5	
Ethylbenzene	100-41-4					5.16	USEPA Reg 5	5.16E+00	USEPA Reg 5	
Famphur	52-85-7					0.0497	USEPA Reg 5	4.97E-02	USEPA Reg 5	
Fluoranthene	206-44-0					122	USEPA Reg 5	1.22E+02	USEPA Reg 5	
Fluorene	86-73-7			30	PRGs ^c	122	USEPA Reg 5	3.00E+01	PRGs	
Furan	110-00-9			600	PRGs			6.00E+02	PRGs	
Heptane	142-82-5							No ESV	No Source	
Heptachlor	76-44-8					0.00598	USEPA Reg 5	5.98E-03	USEPA Reg 5	
Heptachlor Epoxide	1024-57-3					0.152	USEPA Reg 5	1.52E-01	USEPA Reg 5	
Hexachlorobenzene	118-74-1					0.199	USEPA Reg 5	1.99E-01	USEPA Reg 5	

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

		Soil Screening Values								
		USEPA EcoSSLs		DOE (1 Prelim Remediati for Ecol Endpo	997a) inary on Goals logical	USEPA Re	egion 5 Ecological g Levels (2003) of 1998 EDQLs)	Preferred Ecological Screening Value (ESV) ^d		
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	
Hexachlorobutadiene	87-68-3				-	0.0398	USEPA Reg 5	3.98E-02	USEPA Reg 5	
Hexachlorocyclopentadiene	77-47-4			10	PRGs	0.755	USEPA Reg 5	1.00E+01	PRGs	
Hexachloroethane	67-72-1					0.596	USEPA Reg 5	5.96E-01	USEPA Reg 5	
Hexachorophene	70-30-4					0.199	USEPA Reg 5	1.99E-01	USEPA Reg 5	
2-Hexanone	591-78-6					12.6	USEPA Reg 5	1.26E+01	USEPA Reg 5	
HMX	2691-41-0							No ESV	No Source	
Indeno(1,2,3-cd)pyrene	193-39-5					109	USEPA Reg 5	1.09E+02	USEPA Reg 5	
Isobutyl alcohol	78-83-1					20.8	USEPA Reg 5	2.08E+01	USEPA Reg 5	
Isodrin	465-73-6					0.00332	USEPA Reg 5	3.32E-03	USEPA Reg 5	
Isophorone	78-59-1					139	USEPA Reg 5	1.39E+02	USEPA Reg 5	
Isosafrole	120-58-1					9.94	USEPA Reg 5	9.94E+00	USEPA Reg 5	
Kepone	143-50-0					0.0327	USEPA Reg 5	3.27E-02	USEPA Reg 5	
Malathion	121-75-5							No ESV	No Source	
Methacrylonitrile	126-98-7					0.057	USEPA Reg 5	5.70E-02	USEPA Reg 5	
Methapyrilene	91-80-5					2.78	USEPA Reg 5	2.78E+00	USEPA Reg 5	
Methoxychlor	72-43-5					0.0199	USEPA Reg 5	1.99E-02	USEPA Reg 5	
Methyl iodide	74-88-4					1.23	USEPA Reg 5	1.23E+00	USEPA Reg 5	
Methyl methacrylate	80-62-6					984	USEPA Reg 5	9.84E+02	USEPA Reg 5	
Methyl methanesulfanate	66-27-3					0.315	USEPA Reg 5	3.15E-01	USEPA Reg 5	

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

			USEPA EcoSSLs			Screenin	egion 5 Ecological g Levels (2003) of 1998 EDQLs)	Preferred Ecological Screening Value (ESV) ^d		
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	
Methyl parathion	298-00-0					0.00029	USEPA Reg 5	2.92E-04	USEPA Reg 5	
4-Methyl-2-pentanone	108-10-1					443	USEPA Reg 5	4.43E+02	USEPA Reg 5	
3-Methylcholanthrene	56-49-5					0.0779	USEPA Reg 5	7.79E-02	USEPA Reg 5	
Methylene bromide	74-95-3					65	USEPA Reg 5	6.50E+01	USEPA Reg 5	
Methylene chloride	75-09-2					4.05	USEPA Reg 5	4.05E+00	USEPA Reg 5	
2-Methylnaphthalene	91-57-6					3.24	USEPA Reg 5	3.24E+00	USEPA Reg 5	
2-Methylphenol	95-48-7					40.4	USEPA Reg 5	4.04E+01	USEPA Reg 5	
4-Methylphenol	106-44-5					163	USEPA Reg 5	1.63E+02	USEPA Reg 5	
Mirex	2385-85-5							No ESV	No Source	
Naphthalene	91-20-3					0.0994	USEPA Reg 5	9.94E-02	USEPA Reg 5	
1,4-Naphthoquinone	130-15-4					1.67	USEPA Reg 5	1.67E+00	USEPA Reg 5	
1-Naphthylamine	134-32-7					9.34	USEPA Reg 5	9.34E+00	USEPA Reg 5	
2-Naphthylamine	91-59-8					3.03	USEPA Reg 5	3.03E+00	USEPA Reg 5	
2-Nitroaniline	88-74-4					74.1	USEPA Reg 5	7.41E+01	USEPA Reg 5	
3-Nitroaniline	99-09-2					3.16	USEPA Reg 5	3.16E+00	USEPA Reg 5	
4-Nitroaniline	100-01-6					21.9	USEPA Reg 5	2.19E+01	USEPA Reg 5	
Nitrobenzene	99-95-3			40	PRGs ^c	1.31	USEPA Reg 5	4.00E+01	PRGs	
Nitrocellulose	9004-70-0							No ESV	No Source	
Nitroglycerin	55-63-0							No ESV	No Source	

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

		Soil Screening Values								
		USEPA	DOE (1 Prelim Remediati for Eco Endpo	1997a) inary ion Goals logical	USEPA Re	egion 5 Ecological g Levels (2003) ff 1998 EDQLs)	Preferred Ecological Screening Value (ESV) ^d			
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	
Nitroguanidine	556-88-7							No ESV	No Source	
2-Nitrophenol	88-75-5					1.6	USEPA Reg 5	1.60E+00	USEPA Reg 5	
4-Nitrophenol	100-02-7			7	PRGs	5.12	USEPA Reg 5	7.00E+00	PRGs	
4-Nitroquinoline-1-oxide	56-57-5					0.122	USEPA Reg 5	1.22E-01	USEPA Reg 5	
3-Nitrotoluene	99-08-1							No ESV	No Source	
N-Nitrosodiethylamine	55-18-5					0.0693	USEPA Reg 5	6.93E-02	USEPA Reg 5	
N-Nitrosodimethylamine	62-75-9					3.2E-05	USEPA Reg 5	3.21E-05	USEPA Reg 5	
N-Nitrosodiphenylamine	86-30-6			20	PRGs ^c	0.545	USEPA Reg 5	2.00E+01	PRGs	
N-Nitrosomethylethylamine	10595-95-6					0.00166	USEPA Reg 5	1.66E-03	USEPA Reg 5	
N-Nitrosomorpholine	59-89-2					0.0706	USEPA Reg 5	7.06E-02	USEPA Reg 5	
N-Nitrosopiperidine	100-75-4					0.00665	USEPA Reg 5	6.65E-03	USEPA Reg 5	
N-Nitrosopyrrolidine	930-55-2					0.0126	USEPA Reg 5	1.26E-02	USEPA Reg 5	
N-nitroso-di-n-propylamine	621-64-7					0.544	USEPA Reg 5	5.44E-01	USEPA Reg 5	
2-Nitrotoluene	88-72-2							No ESV	No Source	
5-nitro-o-Toluidine	99-55-8					8.73	USEPA Reg 5	8.73E+00	USEPA Reg 5	
2,2'- oxybis(1-Chloropropane)	108-60-1					19.9	USEPA Reg 5	1.99E+01	USEPA Reg 5	
Parathion	56-38-2					0.00034	USEPA Reg 5	3.40E-04	USEPA Reg 5	
PCDDs	PCDD-S					2E-07	USEPA Reg 5	1.99E-07	USEPA Reg 5	
Pentachloroaniline	527-20-8			100	PRGs ^c			1.00E+02	PRGs	

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

			USEPA EcoSSLs			Screenin	gion 5 Ecological g Levels (2003) f 1998 EDQLs)	Preferred Ecological Screening Value (ESV) ^d		
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	
Pentachlorobenzene	608-93-5			20	PRGs	0.497	USEPA Reg 5	2.00E+01	PRGs	
Pentachloroethane	76-01-7					10.7	USEPA Reg 5	1.07E+01	USEPA Reg 5	
Pentachloronitrobenzene	82-68-8					7.09	USEPA Reg 5	7.09E+00	USEPA Reg 5	
Pentachlorophenol	87-86-5	2.1	avian EcoSSL for PCP	3	PRGs	0.119	USEPA Reg 5	2.10E+00	avian EcoSSL for PCP	
PETN	78-11-5							No ESV	No Source	
Phenacetin	62-44-2					11.7	USEPA Reg 5	1.17E+01	USEPA Reg 5	
Phenanthrene	85-01-8					45.7	USEPA Reg 5	4.57E+01	USEPA Reg 5	
Phenol	108-95-2			30	PRGs	120	USEPA Reg 5	3.00E+01	PRGs	
p-Phenylenediamine	106-50-3					6.16	USEPA Reg 5	6.16E+00	USEPA Reg 5	
Phorate	298-02-2					0.0005	USEPA Reg 5	4.96E-04	USEPA Reg 5	
2-Picoline	109-06-8					9.9	USEPA Reg 5	9.90E+00	USEPA Reg 5	
Polychlorinated biphenyls	1336-36-3			0.371	PRGs	0.00033	USEPA Reg 5	3.71E-01	PRGs	
Polychlorinated dibenzofurans	51207-31-9					3.9E-05	USEPA Reg 5	3.86E-05	USEPA Reg 5	
Polynuclear aromatic hydrocarbons	130498-29-2	1.1	mammalian EcoSSL for HMW PAHs					1.10E+00	mammalian EcoSSL for HMW PAHs	
Pronamide	23950-58-5					0.0136	USEPA Reg 5	1.36E-02	USEPA Reg 5	
Propionitrile	107-12-0					0.0498	USEPA Reg 5	4.98E-02	USEPA Reg 5	
4-Nitrotoluene	99-99-0							No ESV	No Source	
Pyrene	129-00-0					78.5	USEPA Reg 5	7.85E+01	USEPA Reg 5	
Pyridine	110-86-1					1.03	USEPA Reg 5	1.03E+00	USEPA Reg 5	

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

		USEPA	DOE (1997a) Preliminary Remediation Goals for Ecological Endpoints ^a		Screenin	egion 5 Ecological g Levels (2003) f 1998 EDQLs)	Preferred Ecological Screening Value (ESV) ^d		
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source
RDX	121-82-4							No ESV	No Source
Safrole	94-59-7					0.404	USEPA Reg 5	4.04E-01	USEPA Reg 5
Silvex (2,4,5-TP)	93-72-1					0.109	USEPA Reg 5	1.09E-01	USEPA Reg 5
Styrene	100-42-5			300	PRGs	4.69	USEPA Reg 5	3.00E+02	PRGs
TCDD (2,3,7,8- Tetrachlorodibenzo-p-dioxin)	1746-01-6			3.15E-06	PRGs	2E-07	USEPA Reg 5	3.15E-06	PRGs
TCDF	51207-31-9			8.40E-04	PRGs	3.9E-05	USEPA Reg 5	8.40E-04	PRGs
2,3,5,6-Tetrachloroaniline	3481-20-7			20	PRGs			2.00E+01	PRGs
1,2,4,5-Tetrachlorobenzene	95-94-3					2.02	USEPA Reg 5	2.02E+00	USEPA Reg 5
1,2,3,4-Tetrachlorobenzene	634-66-2			10	PRGs			1.00E+01	PRGs
1,1,1,2-Tetrachloroethane	630-20-6					225	USEPA Reg 5	2.25E+02	USEPA Reg 5
1,1,2,2-Tetrachloroethane	79-34-5					0.127	USEPA Reg 5	1.27E-01	USEPA Reg 5
Tetrachloroethene	127-18-4					9.92	USEPA Reg 5	9.92E+00	USEPA Reg 5
2,3,4,5-Tetrachlorophenol	4901-51-3			20	PRGs			2.00E+01	PRGs
2,3,4,6-Tetrachlorophenol	58-90-2					0.199	USEPA Reg 5	1.99E-01	USEPA Reg 5
Tetraethyl dithiopyrophosphate	3689-24-5					0.596	USEPA Reg 5	5.96E-01	USEPA Reg 5
Tetryl	479-45-8							No ESV	No Source
Toluene	108-88-3			200	PRGs	5.45	USEPA Reg 5	2.00E+02	PRGs
o-Toluidine	95-53-4					2.97	USEPA Reg 5	2.97E+00	USEPA Reg 5
4-Toluidine	106-49-0							No ESV	No Source

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

					Soil Scree	ning Values				
			USEPA EcoSSLs			Screenin	egion 5 Ecological g Levels (2003) of 1998 EDQLs)	Preferred Ecological Screening Value (ESV) ^d		
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	
Toxaphene	8001-35-2					0.119	USEPA Reg 5	1.19E-01	USEPA Reg 5	
2,4,5-Trichloroaniline	636-30-6			20	PRGs			2.00E+01	PRGs	
1,2,3-Trichlorobenzene	87-61-6			20	PRGs			2.00E+01	PRGs	
1,2,4-Trichlorobenzene	120-82-1			20	PRGs	11.1	USEPA Reg 5	2.00E+01	PRGs	
1,1,1-Trichloroethane	71-55-6					29.8	USEPA Reg 5	2.98E+01	USEPA Reg 5	
1,1,2-Trichloroethane	79-00-5					28.6	USEPA Reg 5	2.86E+01	USEPA Reg 5	
Trichloroethene	79-01-6					12.4	USEPA Reg 5	1.24E+01	USEPA Reg 5	
Trichlorofluoromethane	75-69-4					16.4	USEPA Reg 5	1.64E+01	USEPA Reg 5	
2,4,5-Trichlorophenol	95-95-4			9	PRGs	14.1	USEPA Reg 5	9.00E+00	PRGs	
2,4,6-Trichlorophenol	88-06-2			4	PRGs	9.94	USEPA Reg 5	4.00E+00	PRGs	
1,2,3-Trichloropropane	96-18-4					3.36	USEPA Reg 5	3.36E+00	USEPA Reg 5	
2,4,5-Trichlorophenoxyacetic acid	93-76-5					0.596	USEPA Reg 5	5.96E-01	USEPA Reg 5	
O,O,O-Triethyl phosphorothioate	126-68-1					0.818	USEPA Reg 5	8.18E-01	USEPA Reg 5	
1,3,5-Trinitrobenzene	99-35-4					0.376	USEPA Reg 5	3.76E-01	USEPA Reg 5	
2,4,6-Trinitrotoluene	118-96-7							No ESV	No Source	
Vinyl acetate	108-05-4					12.7	USEPA Reg 5	1.27E+01	USEPA Reg 5	
Vinyl chloride	75-01-4					0.646	USEPA Reg 5	6.46E-01	USEPA Reg 5	

Table C-1. Ecological Screening Values for Chemical Analytes in Soil (continued)

				_	Soil Scree	ning Values			
		USEPA	A EcoSSLs	DOE (1997a) Preliminary Remediation Goals for Ecological Endpoints ^a		USEPA Region 5 Ecological Screening Levels (2003) (update of 1998 EDOLs)		Preferred Ecological Screeniną Value (ESV) ^d	
Analyte	CAS Registry Number	Number (mg/kg dry soil)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source	Number (mg/kg)	Source
Xylenes (total)	1330-20-7					10	USEPA Reg 5	1.00E+01	USEPA Reg 5

- Hierarchy of values found in updated Ohio EPA Risk Assessment Guidance, section 3.3.5: http://www.epa.ohio.gov/portals/30/rules/RR-031.pdf
 - EcoSSLs: http://www.epa.gov/ecotox/ecossl/
- Ecological Screening Levels (ESLs), USEPA Region 5, 2003: http://www.epa.gov/reg5rcra/ca/edql.htm
 - ^aUnited States Department of Energy (DOE) (1997a). Preliminary Remediation Goals for Ecological Endpoints. ES/ER/TM-162/R2. August 1997.
- http://www.esd.ornl.gov/programs/ecorisk/documents/tm162r2.pdf
- bValues for which plant benchmark is lowest. According to DOE (1997a), the PRG is the lowest of three values (earthworm, plant, or wildlife). The only values shown in DOE 1997a are the ones for which the calculated value is lower than earthworm and plant values. Plant values found in: DOE 1997b. Toxicological Benchmarks for Screening Contaminants of Potential Concern for Effects on Terrestrial Plants. ES/ER/TM-85/R3. November 1997.
- ^eValues for which earthworm benchmark is lowest. According to DOE (1997a), the PRG is the lowest of three values (earthworm, plant, or wildlife). The only values shown in DOE 1997a are the ones for which the calculated value is lower than earthworm and plant values. Earthworm values found in: DOE 1997c. Toxicological Benchmarks for Screening Contaminants of Potential Concern for Effects on Soil and Litter Invertebrates and Heterotrophic Process. ES/ER/TM-126/R2.
- ^dThe Preferred Soil Value is the EcoSSLs, followed by DOE (1997a), followed by USEPA Region 5 ESLs.
- *Aluminum is identified as a chemical of potential concern (COPC) only at sites where the soil pH is less than 5.5
- 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 **In well-aerated soils between pH 5 and 8, iron is not expected to be toxic to plants. A determination of the geochemical conditions (i.e., pH and Eh at a minimum) of the environmental setting, as well as the presence of iron floc and the toxic metals, is critical to the determination of the relative importance of iron at an area of concern (AOC).
- -- = no value
- CAS = Chemical Abstract Service
- EDQL = Ecological Data Quality Level
- EcoSSL = Ecological Soil Screening Level
- Ohio EPA = Ohio Environmental Protection Agency
- PRG = Preliminary Remediation Goal
- RDX = Hexahydro-1,3,5-trinitro-1,3,5-triazine
- Reg = Region
- USEPA = United States Environmental Protection Agency

THIS PAGE INTENTIONALLY LEFT BLANK.

1

Table C-2. Ecological Screening Values for Chemical Analytes in Surface Water

						Surfa	nce Water Screening Values				
					Updated Values			USEPA	Region 5 ESLs		
					- F	T			$(2003)^{c}$		
	CAS		Ohio EPA OMZM ^a	NAW(QC 2009 Update		Tier II Values		of 1998 EDQLs)		Preferred Surface Water Value ^d
	Registry	Number		Number		Number		Number		Number	
Analyte	Number	(µg/L)	Reference	(µg/L)	Reference	(µg/L)	Reference	(µg/L)	Reference	(µg/L)	Reference
					Inorganic Ch	emicals					
Aluminum	7429-90-5			87	NAWQC 2009	100	Tier II (GLI database)		-	8.70E+01	NAWQC 2009
Ammonia	7664-41-7	500	Ohio Administrative Codetemp&pH dependent							5.00E+02	Ohio Administrative Codetemp&pH dependent
Antimony	7440-36-0	900	Ohio Administrative Code			30	Tier II (Suter & Tsao 1996)	80	USEPA Reg 5	9.00E+02	Ohio Administrative Code
Arsenic III (Diss)	7440-38-2	340	Ohio Administrative Code	150	NAWQC 2009				-	3.40E+02	Ohio Administrative Code
Arsenic	7440-38-2	340	Ohio Administrative Code					148	USEPA Reg 5	3.40E+02	Ohio Administrative Code
Arsenic V (Diss)	7440-38-2					3.1	Tier II (Suter & Tsao 1996)			3.10E+00	Tier II (Suter & Tsao 1996)
Barium	7440-39-3	2,000	Ohio Administrative Code			4.0	Tier II (Suter & Tsao 1996)	220	USEPA Reg 5	2.00E+03	Ohio Administrative Code
Beryllium	7440-41-7	93	Ohio Administrative Codehardness dependent			0.66	Tier II (Suter & Tsao 1996)	3.6	USEPA Reg 5	9.30E+01	Ohio Administrative Codehardness dependent
Boron	7440-42-8	33,000	Ohio Administrative Code			1.6	Tier II (Suter & Tsao 1996)			3.30E+04	Ohio Administrative Code
Cadmium	7440-43-9	4.5	Ohio Administrative Codehardness dependent			0.2	Tier II (Suter & Tsao 1996)	0.15	USEPA Reg 5	4.50E+00	Ohio Administrative Codehardness dependent
Cadmium (Diss)	7440-43-9	4.3	Ohio Administrative Codehardness dependent	0.25	NAWQC 2009					4.30E+00	Ohio Administrative Codehardness dependent
Calcium	7440-70-2									No ESV	No Source
Chlorine (total residual)	7782-50-5	19	Ohio Administrative Code	11	NAWQC 2009	5	Tier II (GLI database)			1.90E+01	Ohio Administrative Code
Chromium III (Diss)	7440-47-3	570	Ohio Administrative Codehardness dependent	74	NAWQC 2009	210	Tier II (Suter & Tsao 1996)	42	USEPA Reg 5	5.70E+02	Ohio Administrative Codehardness dependent
Chromium	7440-47-3	1,800	Ohio Administrative Codehardness dependent					42	USEPA Reg 5	1.80E+03	Ohio Administrative Codehardness dependent
Chromium VI (Diss)	7440-47-3	16	Ohio Administrative Code	11	NAWQC 2009	11	Tier II (Suter & Tsao 1996)			1.60E+01	Ohio Administrative Code
Cobalt	7440-48-4	220	Ohio Administrative Code			23	Tier II (Suter & Tsao 1996)	24	USEPA Reg 5	2.20E+02	Ohio Administrative Code
Copper (Diss)	7440-50-8	13	Ohio Administrative Codehardness dependent	1.45	NAWQC 2009					1.30E+01	Ohio Administrative Codehardness dependent
Copper	7440-50-8	14	Ohio Administrative Codehardness dependent					1.58	USEPA Reg 5	1.40E+01	Ohio Administrative Codehardness dependent
Cyanide	57-12-5	22	Ohio Administrative Code	5.2	NAWQC 2009			5.2	USEPA Reg 5	2.20E+01	Ohio Administrative Code
Iron	7439-89-6			1,000	NAWQC 2009	300	Tier II (GLI database)			1.00E+03	NAWQC 2009
Lead (Diss)	7439-92-1	97	Ohio Administrative Codehardness dependent	2.5	NAWQC 2009					9.70E+01	Ohio Administrative Codehardness dependent
Lead	7439-92-1	120	Ohio Administrative Codehardness dependent					1.17	USEPA Reg 5	1.20E+02	Ohio Administrative Codehardness dependent
Lithium						14	Tier II (Suter & Tsao 1996)			1.40E+01	Tier II (Suter & Tsao 1996)
Magnesium	7439-95-4									No ESV	No Source
Manganese	7439-96-5					120	Tier II (Suter & Tsao 1996)			1.20E+02	Tier II (Suter & Tsao 1996)
Mercury	7439-97-6	1.7	Ohio Administrative Code			1.3	Tier II (Suter & Tsao 1996)	0.0013		1.70E+00	Ohio Administrative Code
Mercury (Diss)	7439-97-6	1.4	Ohio Administrative Code	0.77	NAWQC 2009					1.40E+00	Ohio Administrative Code
Mercury, methyl	22967-92-6					0.0028	Tier II (Suter & Tsao 1996)	2.46E-03		2.80E-03	Tier II (Suter & Tsao 1996)
Molybdenum	7439-98-7	190,000	Ohio Administrative Code			370	Tier II (Suter & Tsao 1996)			1.90E+05	Ohio Administrative Code
Nickel (Diss)	7440-02-0	470	Ohio Administrative Codehardness dependent	52	NAWQC 2009					4.70E+02	Ohio Administrative Codehardness dependent
Nickel (TR)	7440-02-1	470	Ohio Administrative Codehardness dependent					28.9	USEPA Reg 5	4.70E+02	Ohio Administrative Codehardness dependent
Potassium	7440-09-7									No ESV	No Source
Selenium (Diss)	7782-49-2			4.6	NAWQC 2009					4.61E+00	NAWQC 2009
Selenium	7782-49-2			5	NAWQC 2009			5	USEPA Reg 5	5.00E+00	NAWQC 2009
Silver (Diss)	7440-22-4	1.4	Ohio Administrative Codehardness dependent			0.12	Tier II (Suter & Tsao 1996)			1.40E+00	Ohio Administrative Codehardness dependent
Silver	7440-22-4	1.6	Ohio Administrative Codehardness dependent			0.36	Tier II (Suter & Tsao 1996)	0.12	USEPA Reg 5	1.60E+00	Ohio Administrative Codehardness dependent
Sodium	7440-23-5									No ESV	No Source
Strontium	7440-24-6	40,000	Ohio Administrative Code			1,500	Tier II (Suter & Tsao 1996)			4.00E+04	Ohio Administrative Code
Thallium	7440-28-0	79	Ohio Administrative Code			12	Tier II (Suter & Tsao 1996)	10	USEPA Reg 5	7.90E+01	Ohio Administrative Code
Tin	7440-31-5	1,600	Ohio Administrative Code			73	Tier II (Suter & Tsao 1996)	180	USEPA Reg 5	1.60E+03	Ohio Administrative Code

Table C-2. Ecological Screening Values for Chemical Analytes in Surface Water (continued)

						Surfa	nce Water Screening Values				
					Updated Values	for Suter a	nd Tsao 1996 ^b	USEPA	Region 5 ESLs		
									(2003) ^c		
	CAS		Ohio EPA OMZM ^a		QC 2009 Update		Tier II Values	(update o	of 1998 EDQLs)		Preferred Surface Water Value ^d
Analyte	Registry Number	Number (µg/L)	Reference	Number (µg/L)	Reference	Number (µg/L)	Reference	Number (µg/L)	Reference	Number (µg/L)	Reference
Uranium						2.6	Tier II (Suter & Tsao 1996)			2.60E+00	Tier II (Suter & Tsao 1996)
Vanadium	7440-62-2	150	Ohio Administrative Code			20	Tier II (Suter & Tsao 1996)	12	USEPA Reg 5	1.50E+02	Ohio Administrative Code
Zinc (Diss)	7440-66-6	120	Ohio Administrative Code	120	NAWQC 2009					1.20E+02	Ohio Administrative Code
Zinc (TR)	7440-66-6	120	Ohio Administrative Code					65.7	USEPA Reg 5	1.20E+02	Ohio Administrative Code
Zirconium						17	Tier II (Suter & Tsao 1996)			1.70E+01	Tier II (Suter & Tsao 1996)
	'	<u>l</u>		l l	Anions	1	l	l .		1	
Chloride	16887-00-6			230,000	NAWQC 2009					2.30E+05	NAWQC 2009
Fluoride	16984-48-8					3,400	Tier II (GLI database)			3.40E+03	Tier II (GLI database)
Hydrogen Sulfide	7783-06-4			2	NAWQC 2009	2	Tier II (GLI database)			2.00E+00	NAWQC 2009
Nitrate	14797-55-8									No ESV	No Source
Nitrite	14797-65-0					20	Tier II (GLI database)			2.00E+01	Tier II (GLI database)
Sulfite	14265-45-3					200	Tier II (GLI database)			2.00E+02	Tier II (GLI database)
					Organic Che	micals		•			
Acenaphthene	83-32-9	19	Ohio Administrative Code			5.3	Tier II (GLI database)	38	USEPA Reg 5	1.90E+01	Ohio Administrative Code
Acenaphthylene	208-96-8	120	Ohio Administrative Code, Lake Erie					4,840	USEPA Reg 5	1.20E+02	Ohio Administrative Code, Lake Erie
Acetaldehyde	75-07-0					130	Tier II (GLI database)			1.30E+02	Tier II (GLI database)
Acetone	67-64-1					1,500	Tier II (Suter & Tsao 1996)	1,700	USEPA Reg 5	1.50E+03	Tier II (Suter & Tsao 1996)
Acetonitrile	75-05-8	100,000	Ohio Administrative Code			12,000	Tier II (GLI database)	12,000	USEPA Reg 5	1.00E+05	Ohio Administrative Code
Acetylaminofluorene[2-]	53-96-3							535	USEPA Reg 5	5.35E+02	USEPA Reg 5
Acrolein	107-02-8			3	NAWQC 2009	0.19	Tier II (GLI database)	0.19	USEPA Reg 5	3.00E+00	NAWQC 2009
Acrylonitrile	107-13-1	650	Ohio Administrative Code			78	Tier II (GLI database)	66	USEPA Reg 5	6.50E+02	Ohio Administrative Code
Alachlor	15972-60-8					21	Tier II (GLI database)			2.10E+01	Tier II (GLI database)
Aldrin	309-00-2					0.035	Tier II (GLI database)	0.017	USEPA Reg 5	3.50E-02	Tier II (GLI database)
2-Amino-4,6-dinitrotoluene	35572-78-2	160	Ohio Administrative Code			18	Tier II (GLI database)			1.60E+02	Ohio Administrative Code
4-Amino-2,6-dinitrotoluene	19406-51-0	98	Ohio Administrative Code			11	Tier II (GLI database)			9.80E+01	Ohio Administrative Code
Aniline	62-53-3	30	Ohio Administrative Code			4.1	Tier II (GLI database)	4.1	USEPA Reg 5	3.00E+01	Ohio Administrative Code
Anthracene	120-12-7	0.18	Ohio Administrative Code			0.73	Tier II (Suter & Tsao 1996)	0.035	USEPA Reg 5	1.80E-01	Ohio Administrative Code
Aramite	140-57-8							3.09	USEPA Reg 5	3.09E+00	USEPA Reg 5
Azobenzene[p-(dimethylamino)]	60-11-7							1.65	USEPA Reg 5	1.65E+00	USEPA Reg 5
Benzene	71-43-2	700	Ohio Administrative Code			130	Tier II (Suter & Tsao 1996)	114	USEPA Reg 5	7.00E+02	Ohio Administrative Code
Benzenemethanol	100-51-6					8.6	Tier II (Suter & Tsao 1996)	8.6	USEPA Reg 5	8.60E+00	Tier II (Suter & Tsao 1996)
Benzidine						3.9	Tier II (Suter & Tsao 1996)			3.90E+00	Tier II (Suter & Tsao 1996)
Benz(a)anthracene	56-55-3	42	Ohio Administrative Code, Lake Erie			0.027	Tier II (Suter & Tsao 1996)	0.025	USEPA Reg 5	4.20E+01	Ohio Administrative Code, Lake Erie
Benzo(a)pyrene	50-32-8	0.54	Ohio Administrative Code, Lake Erie			0.014	Tier II (Suter & Tsao 1996)	0.014	USEPA Reg 5	5.40E-01	Ohio Administrative Code, Lake Erie
Benzo(b)fluoranthene	205-99-2	23	Ohio Administrative Code, Lake Erie			2.6	Tier II (GLI database)	9.07	USEPA Reg 5	2.30E+01	Ohio Administrative Code, Lake Erie
Benzo(g,h,i)perylene	191-24-2							7.64	USEPA Reg 5	7.64E+00	USEPA Reg 5
Benzoic Acid	65-85-0					42	Tier II (Suter & Tsao 1996)			4.20E+01	Tier II (Suter & Tsao 1996)
BHC, alpha	319-84-6							12.4	USEPA Reg 5	1.24E+01	USEPA Reg 5
BHC, beta	319-85-7							0.495	USEPA Reg 5	4.95E-01	USEPA Reg 5
BHC, delta	319-86-8							667	USEPA Reg 5	6.67E+02	USEPA Reg 5
BHC, gamma (lindane)	58-89-9	0.95	Ohio Administrative Code			0.057	Tier II (GLI database)	0.026	USEPA Reg 5	9.50E-01	Ohio Administrative Code
Biphenyl	92-52-4	26	Ohio Administrative Code			6.5	Tier II (GLI database)			2.60E+01	Ohio Administrative Code

Table C-2. Ecological Screening Values for Chemical Analytes in Surface Water (continued)

						Surfa	ce Water Screening Values					
					Updated Values			USEPA	Region 5 ESLs			
					e punteu varues	lor succi a	1340 1770		(2003) ^c			
	CAS		Ohio EPA OMZM ^a	NAW(QC 2009 Update		Tier II Values		of 1998 EDQLs)		Preferred Surface Water Value ^d	
	Registry	Number		Number		Number		Number		Number		
Analyte	Number	(μg/L)	Reference	(μg/L)	Reference	(μg/L)	Reference	(μg/L)	Reference	(μg/L)	Reference	
Bis(2-chloroethyl) ether	111-44-4							19,000	USEPA Reg 5	1.90E+04	USEPA Reg 5	
Bis(2-ethylhexyl)phthalate	117-81-7	1,100	Ohio Administrative Code			3.0	Tier II (Suter & Tsao 1996)	0.3	USEPA Reg 5	1.10E+03	Ohio Administrative Code	
Bromodichloromethane	74-97-5	3,100	Ohio Administrative Code, Lake Erie							3.10E+03	Ohio Administrative Code, Lake Erie	
Bromomethane (methyl bromide)	74-83-9	38	Ohio Administrative Code			16	Tier II (GLI database)	16	USEPA Reg 5	3.80E+01	Ohio Administrative Code	
4-Bromophenyl-phenylether	101-55-3							1.5	USEPA Reg 5	1.50E+00	USEPA Reg 5	
2-Butanone (methyl ethyl ketone)	78-93-3	200,000	Ohio Administrative Code			22,000	Tier II (GLI database)	2,200	USEPA Reg 5	2.00E+05	Ohio Administrative Code	
Butylbenzylphthalate	85-68-7	130	Ohio Administrative Code			23	Tier II (GLI database)	23	USEPA Reg 5	1.30E+02	Ohio Administrative Code	
Carbofuran	1563-66-2					1	Tier II (GLI database)			1.00E+00	Tier II (GLI database)	
Carbon Disulfide	75-15-0	130	Ohio Administrative Code			15	Tier II (GLI database)	15	USEPA Reg 5	1.30E+02	Ohio Administrative Code	
Carbon Tetrachloride	56-23-5	2,200	Ohio Administrative Code			240	Tier II (GLI database)	240	USEPA Reg 5	2.20E+03	Ohio Administrative Code	
Chlordane	57-74-9			0.0043	NAWQC 2009			0.0043	USEPA Reg 5	4.30E-03	NAWQC 2009	
4-Chloroaniline	106-47-8							232	USEPA Reg 5	2.32E+02	USEPA Reg 5	
Chlorobenzene	108-90-7	420	Ohio Administrative Code			47	Tier II (GLI database)	47	USEPA Reg 5	4.20E+02	Ohio Administrative Code	
Chlorobenzilate	510-15-6							7.16	USEPA Reg 5	7.16E+00	USEPA Reg 5	
Chloroform	67-66-3	1,300	Ohio Administrative Code			140	Tier II (GLI database)	140	USEPA Reg 5	1.30E+03	Ohio Administrative Code	
2-Chloronaphthalene	91-58-7							0.396	USEPA Reg 5	3.96E-01	USEPA Reg 5	
2-Chlorophenol	95-57-8	290	Ohio Administrative Code			32	Tier II (GLI database)	24	USEPA Reg 5	2.90E+02	Ohio Administrative Code	
Chloropyrifos	2921-88-2			0.041	NAWQC 2009					4.10E-02	NAWQC 2009	
4-Chloro-3-methylphenol	59-50-7							34.8	USEPA Reg 5	3.48E+01	USEPA Reg 5	
Chrysene	218-01-9	42	Ohio Administrative Code, Lake Erie							4.20E+01	Ohio Administrative Code, Lake Erie	
Cyanazine	21725-46-2					270	Tier II (GLI database)			2.70E+02	Tier II (GLI database)	
2,4-D	94-75-7					240	Tier II (GLI database)	220	USEPA Reg 5	2.40E+02	Tier II (GLI database)	
4,4'-DDD	72-54-8									No ESV	No Source	
4,4'-DDE	72-54-8							4.51E-09	USEPA Reg 5	4.51E-09	USEPA Reg 5	
4,4'-DDT	50-29-3	11	Erie OMZA for DDT+met, Table 33-2 of OAC	0.001	NAWQC 2009	+		1.10E-05	USEPA Reg 5		Erie OMZA for DDT+met, Table 33-2 of OAC	
	8065-48-3		Effe OMZA for DD1+fflet, Table 55-2 of OAC	0.001		0.1	 T: II (CLI d-t-b)			1.10E+01	· · · · · · · · · · · · · · · · · · ·	
Demeton				0.1	NAWQC 2009	0.1	Tier II (GLI database)			1.00E-01	NAWQC 2009	
Diazinon	333-41-5			0.17	NAWQC 2009	0.08	Tier II (GLI database)			1.70E-01	NAWQC 2009	
Dibenzofuran	132-64-9	36	Ohio Administrative Code			4	Tier II (GLI database)	4	USEPA Reg 5	3.60E+01	Ohio Administrative Code	
Dibromochloromethane	124-48-1	2,900	Ohio Administrative Code, Lake Erie							2.90E+03	Ohio Administrative Code, Lake Erie	
2,2-Dibromo-3-nitrilopropionamide	10222-01-2					20	Tier II (GLI database)			2.00E+01	Tier II (GLI database)	
1,2-Dichlorobenzene	95-50-1	130	Ohio Administrative Code			23	Tier II (GLI database)	14	USEPA Reg 5	1.30E+02	Ohio Administrative Code	
1,3-Dichlorobenzene	541-73-1	79	Ohio Administrative Code			22	Tier II (GLI database)	38	USEPA Reg 5	7.90E+01	Ohio Administrative Code	
1,4-Dichlorobenzene	106-46-7	57	Ohio Administrative Code			9.4	Tier II (GLI database)	9.4	USEPA Reg 5	5.70E+01	Ohio Administrative Code	
Dichlorobenzene	25321-22-6					5	Tier II (GLI database)			5.00E+00	Tier II (GLI database)	
3,3'-Dichlorobenzidine	91-94-1							4.5	USEPA Reg 5	4.50E+00	USEPA Reg 5	
1,1-Dichloroethane	75-34-3	3,700	Ohio Administrative Code, Lake Erie			740	Tier II (GLI database)	47	USEPA Reg 5	3.70E+03	Ohio Administrative Code, Lake Erie	
1,2-Dichloroethane	107-06-2	9,600	Ohio Administrative Code			2,000	Tier II (GLI database)	910	USEPA Reg 5	9.60E+03	Ohio Administrative Code	
1,1-Dichloroethene	75-35-4	1,900	Ohio Administrative Code			210	Tier II (GLI database)	65	USEPA Reg 5	1.90E+03	Ohio Administrative Code	
1,2-Dichloroethene	540-59-0	8,800	Ohio Administrative Code			970	Tier II (GLI database)	970	USEPA Reg 5	8.80E+03	Ohio Administrative Code	
2,4-Dichlorophenol	120-83-2	110	Ohio Administrative Code			11	Tier II (GLI database)	11	USEPA Reg 5	1.10E+02	Ohio Administrative Code	
1,2-Dichloropropane	78-87-5	3,300	Ohio Administrative Code			520	Tier II (GLI database)	360	USEPA Reg 5	3.30E+03	Ohio Administrative Code	
1,3-Dichloropropene	542-75-6	15	Ohio Administrative Code			1.7	Tier II (GLI database)			1.50E+01	Ohio Administrative Code	
Dieldrin	60-57-1	0.24	Ohio Administrative Code	0.056	NAWQC 2009			7.10E-05	USEPA Reg 5	2.40E-01	Ohio Administrative Code	
Diethylphthalate	84-66-2	980	Ohio Administrative Code			220	Tier II (GLI database)	110	USEPA Reg 5	9.80E+02	Ohio Administrative Code	
7,12'-Dimethylbenz(a)anthracene	57-97-6							0.548	USEPA Reg 5	5.48E-01	USEPA Reg 5	

Table C-2. Ecological Screening Values for Chemical Analytes in Surface Water (continued)

						Surfa	ce Water Screening Values				
					Updated Values			1	Region 5 ESLs		
									(2003) ^c		
	CAS		Ohio EPA OMZM ^a	NAW	QC 2009 Update		Tier II Values		of 1998 EDQLs)		Preferred Surface Water Value ^d
	Registry	Number		Number		Number		Number		Number	
Analyte	Number	(µg/L)	Reference	(µg/L)	Reference	(µg/L)	Reference	(µg/L)	Reference	(µg/L)	Reference
2,4-Dimethylphenol	105-67-9	140	Ohio Administrative Code			15	Tier II (GLI database)	100	USEPA Reg 5	1.40E+02	Ohio Administrative Code
Dimethylphthalate	131-11-3	3,200	Ohio Administrative Code			1,100	Tier II (GLI database)			3.20E+03	Ohio Administrative Code
Di-n-butyl phthalate	84-74-2					19	Tier II (GLI database)	9.7	USEPA Reg 5	1.90E+01	Tier II (GLI database)
Di-n-octylphthalate	117-84-0							30	USEPA Reg 5	3.00E+01	USEPA Reg 5
3,5-Dinitroaniline	618-87-1	210	Ohio Administrative Code			70	Tier II (GLI database)			2.10E+02	Ohio Administrative Code
1,3-Dinitrobenzene	99-65-0	100	Ohio Administrative Code			22	Tier II (GLI database)	22	USEPA Reg 5	1.00E+02	Ohio Administrative Code
2,4-Dinitrophenol	51-28-5							19	USEPA Reg 5	1.90E+01	USEPA Reg 5
2,3-Dinitrotoluene	602-01-7	21	Ohio Administrative Code			2.3	Tier II (GLI database)			2.10E+01	Ohio Administrative Code
2,4-Dinitrotoluene	121-14-2	390	Ohio Administrative Code			44	Tier II (GLI database)	44	USEPA Reg 5	3.90E+02	Ohio Administrative Code
2,5-Dinitrotoluene	619-15-8	50	Ohio Administrative Code			5.6	Tier II (GLI database)			5.00E+01	Ohio Administrative Code
2,6-Dinitrotoluene	606-20-2	730	Ohio Administrative Code			81	Tier II (GLI database)	81	USEPA Reg 5	7.30E+02	Ohio Administrative Code
3,5-Dinitrotoluene	618-85-9	860	Ohio Administrative Code			95	Tier II (GLI database)			8.60E+02	Ohio Administrative Code
4,6-Dinitro-2-methylphenol	534-52-1							23	USEPA Reg 5	2.30E+01	USEPA Reg 5
Dinoseb	88-85-7							0.48	USEPA Reg 5	4.80E-01	USEPA Reg 5
1,4-Dioxane	123-91-1							22,000	USEPA Reg 5	2.20E+04	USEPA Reg 5
Diphenylamine	122-39-4							412	USEPA Reg 5	4.12E+02	USEPA Reg 5
1,2-Diphenylhydrazine	122-66-7					1.1	Tier II (GLI database)			1.10E+00	Tier II (GLI database)
Disulfoton	298-04-4							0.0402	USEPA Reg 5	4.02E-02	USEPA Reg 5
Endosulfan	115-29-7					0.009	Tier II (GLI database)			9.00E-03	Tier II (GLI database)
Endosulfan I (alpha)	959-98-8			0.056	NAWQC 2009			0.056	USEPA Reg 5	5.60E-02	NAWQC 2009
Endosulfan II (beta)	33213-65-9			0.056	NAWQC 2009			0.056	USEPA Reg 5	5.60E-02	NAWQC 2009
Endosulfan Sulfate	1031-07-8							2.22	USEPA Reg 5	2.22E+00	USEPA Reg 5
Endrin	72-20-8	0.086	Ohio Administrative Code	0.036	NAWQC 2009			0.036	USEPA Reg 5	8.60E-02	Ohio Administrative Code
Endrin Aldehyde	7421-93-4							0.15	USEPA Reg 5	1.50E-01	USEPA Reg 5
Ethylbenzene	100-41-4	550	Ohio Administrative Code			61	Tier II (GLI database)	14	USEPA Reg 5	5.50E+02	Ohio Administrative Code
Ethylene Glycol	107-21-1	1,300,000	Ohio Administrative Code			140,000	Tier II (GLI database)			1.30E+06	Ohio Administrative Code
					NAWQC (Suter &						
Fluoranthene	206-44-0	3.7	Ohio Administrative Code	6.16	Tsao 1996)	0.8	Tier II (GLI database)	1.9	USEPA Reg 5	3.70E+00	Ohio Administrative Code
Fluorene	86-73-7	110	Ohio Administrative Code			19	Tier II (GLI database)	19	USEPA Reg 5	1.10E+02	Ohio Administrative Code
Formaldehyde	50-00-0					74	Tier II (GLI database)			7.40E+01	Tier II (GLI database)
Guthion	86-50-0			0.01	NAWQC 2009	0.005	Tier II (GLI database)			1.00E-02	NAWQC 2009
Heptachlor	76-44-8			0.0038	NAWQC 2009			3.80E-03	USEPA Reg 5	3.80E-03	NAWQC 2009
Heptachlor Epoxide	1024-57-3			0.0038	NAWQC 2009			3.80E-03	USEPA Reg 5	3.80E-03	NAWQC 2009
Hexachlorobenzene	118-74-1							3.00E-04	USEPA Reg 5	3.00E-04	USEPA Reg 5
Hexachlorobutadiene	87-68-3					1	Tier II (GLI database)	0.053	USEPA Reg 5	1.00E+00	Tier II (GLI database)
Hexachlorocyclopentadiene	77-47-4					0.45	Tier II (GLI database)	77	USEPA Reg 5	4.50E-01	Tier II (GLI database)
Hexachloroethane	67-72-1							8	USEPA Reg 5	8.00E+00	USEPA Reg 5

Table C-2. Ecological Screening Values for Chemical Analytes in Surface Water (continued)

						Surfa	ce Water Screening Values				
					Updated Values	for Suter a	nd Tsao 1996 ^b	USEPA	Region 5 ESLs		
								(2003) ^c			
	CAS		Ohio EPA OMZM ^a	NAW(C 2009 Update	Tier II Values		(update of 1998 EDQLs)			Preferred Surface Water Value ^d
	Registry	Number		Number		Number		Number		Number	
Analyte	Number	(µg/L)	Reference	(µg/L)	Reference	(µg/L)	Reference	(µg/L)	Reference	(µg/L)	Reference
Hexachlorophene	70-30-4							0.228	USEPA Reg 5	2.28E-01	USEPA Reg 5
2-Hexanone	591-78-6							99	USEPA Reg 5	9.90E+01	USEPA Reg 5
HMX	2691-41-0	1,200	Ohio Administrative Code			220	Tier II (GLI database)			1.20E+03	Ohio Administrative Code
Hydroquinone	123-31-9					2.2	Tier II (GLI database)			2.20E+00	Tier II (GLI database)
Indeno(1,2,3-cd)pyrene	193-39-5							4.31	USEPA Reg 5	4.31E+00	USEPA Reg 5
Isodecyl diphenyl phosphate	29761-21-5					1.73	Tier II (GLI database)			1.73E+00	Tier II (GLI database)
Isodrin	465-73-6							0.0309	USEPA Reg 5	3.09E-02	USEPA Reg 5
Isophorone	78-59-1	7,500	Ohio Administrative Code			920	Tier II (GLI database)	920	USEPA Reg 5	7.50E+03	Ohio Administrative Code
Isopropylbenzene	98-82-8	43	Ohio Administrative Code			4.8	Tier II (GLI database)			4.30E+01	Ohio Administrative Code
4-Isopropyltoluene	99-87-6	150	Ohio Administrative Code			16	Tier II (GLI database)			1.50E+02	Ohio Administrative Code
Kepone	143-50-0							0.132	USEPA Reg 5	1.32E-01	USEPA Reg 5
Malathion	121-75-5			0.1	NAWQC 2009	0.1	Tier II (GLI database)			1.00E-01	NAWQC 2009
MBAS (foaming agents, aesthetic					-						
criteria)		500	Ohio Administrative Code							5.00E+02	Ohio Administrative Code
Methanol	67-56-1					330	Tier II (GLI database)			3.30E+02	Tier II (GLI database)
Methoxychlor	72-43-5			0.03	NAWQC 2009	0.03	Tier II (GLI database)	0.019	USEPA Reg 5	3.00E-02	NAWQC 2009
Methyl Metharcrylate	80-62-6							2,800	USEPA Reg 5	2.80E+03	USEPA Reg 5
4-Methyl-2-pentanone	108-10-1							170	USEPA Reg 5	1.70E+02	USEPA Reg 5
Methyl tert-butyl ether	1634-04-4	6,500	Ohio Administrative Code	51,000	NAWQC 2009	730	Tier II (GLI database)			6.50E+03	Ohio Administrative Code
Methylamine	74-89-5					860	Tier II (GLI database)			8.60E+02	Tier II (GLI database)
3-Methylcholanthrene	56-49-5							0.0891	USEPA Reg 5	8.91E-02	USEPA Reg 5
Methylene Chloride											
(dichloromethane)	75-09-2	11,000	Ohio Administrative Code			2,200	Tier II (Suter & Tsao 1996)	940	USEPA Reg 5	1.10E+04	Ohio Administrative Code
Methylene Dithiocyanate	6317-18-6					1	Tier II (GLI database)			1.00E+00	Tier II (GLI database)
2-Methylnaphthalene	91-57-6					4.7	Tier II (GLI database)	330	USEPA Reg 5	4.70E+00	Tier II (GLI database)
2-Methylphenol	95-48-7	600	Ohio Administrative Code			67	Tier II (GLI database)	67	USEPA Reg 5	6.00E+02	Ohio Administrative Code
3-Methylphenol	108-39-4	560	Ohio Administrative Code			62	Tier II (GLI database)	62	USEPA Reg 5	5.60E+02	Ohio Administrative Code
4-Methylphenol	106-44-5	480	Ohio Administrative Code			53	Tier II (GLI database)	25	USEPA Reg 5	4.80E+02	Ohio Administrative Code
Mirex	2385-85-5			0.001	NAWQC 2009	0.001	Tier II (GLI database)			1.00E-03	NAWQC 2009
Naphthalene	91-20-3	170	Ohio Administrative Code			21	Tier II (GLI database)	13	USEPA Reg 5	1.70E+02	Ohio Administrative Code
Nitrilotriacetic Acid	139-13-9					5000	Tier II (GLI database)			5.00E+03	Tier II (GLI database)
Nitrobenzene	99-95-3	2,000	Ohio Administrative Code			380	Tier II (GLI database)	220	USEPA Reg 5	2.00E+03	Ohio Administrative Code
Nitroglycerin	55-63-0	160	Ohio Administrative Code			18	Tier II (GLI database)			1.60E+02	Ohio Administrative Code
2-Nitrophenol	88-75-5	650	Ohio Administrative Code			73	Tier II (GLI database)			6.50E+02	Ohio Administrative Code
4-Nitrophenol	100-02-7					58	Tier II (GLI database)	60	USEPA Reg 5	5.80E+01	Tier II (GLI database)
N-Nitrosodiethylamine	55-18-5							768	USEPA Reg 5	7.68E+02	USEPA Reg 5
N-Nitrosodiphenylamine	86-30-6					25	Tier II (GLI database)			2.50E+01	Tier II (GLI database)
2-Nitrotoluene	88-72-2	640	Ohio Administrative Code			71	Tier II (GLI database)			6.40E+02	Ohio Administrative Code
3-Nitrotoluene	99-08-1	380	Ohio Administrative Code			42	Tier II (GLI database)			3.80E+02	Ohio Administrative Code
4-Nitrotoluene	99-99-0	410	Ohio Administrative Code			46	Tier II (GLI database)			4.10E+02	Ohio Administrative Code
Nonylphenol	84852-15-3			28	NAWQC 2009					2.80E+01	NAWQC 2009
Oil & Grease (aesthetic criteria)		10,000	Ohio Administrative Code							1.00E+04	Ohio Administrative Code

Table C-2. Ecological Screening Values for Chemical Analytes in Surface Water (continued)

						Surfa	ce Water Screening Values				
					Updated Values	for Suter an	nd Tsao 1996 ^b	USEPA	Region 5 ESLs		
									(2003) ^c		
	CAS		Ohio EPA OMZM ^a	NAW	QC 2009 Update		Tier II Values	(update o	f 1998 EDQLs)		Preferred Surface Water Value ^d
	Registry	Number		Number		Number		Number		Number	
Analyte	Number	(µg/L)	Reference	(µg/L)	Reference	(µg/L)	Reference	$(\mu g/L)$	Reference	(µg/L)	Reference
Parathion	56-38-2	0.065	Ohio Administrative Code	0.013	NAWQC 2009			0.013	USEPA Reg 5	6.50E-02	Ohio Administrative Code
PCDDs	PCDD-S							2.78E-07	USEPA Reg 5	2.78E-07	USEPA Reg 5
Pentachlorobenzene	608-93-5					3.1	Tier II (GLI database)	0.019	USEPA Reg 5	3.10E+00	Tier II (GLI database)
Pentachloroethane	76-01-7							56.4	USEPA Reg 5	5.64E+01	USEPA Reg 5
Pentachlorophenol	87-86-5	5.3	Ohio Administrative CodepH dependent	15	NAWQC 2009			4	USEPA Reg 5	5.30E+00	Ohio Administrative CodepH dependent
Perchlorate	14797-73-0	20,000	Ohio Administrative Code							2.00E+04	Ohio Administrative Code
Phenanthrene	85-01-8	31	Ohio Administrative Code		-	2.3	Tier II (GLI database)	3.6	USEPA Reg 5	3.10E+01	Ohio Administrative Code
Phenol	108-95-2	4,700	Ohio Administrative Code		-	400	Tier II (GLI database)	180	USEPA Reg 5	4.70E+03	Ohio Administrative Code
Phenol (cold water and salmon											
spawning habitat)	108-95-2	4,600	Ohio Administrative Code							4.60E+03	Ohio Administrative Code
Phorate	298-02-2				1			3.62	USEPA Reg 5	3.62E+00	USEPA Reg 5
PCBs	1336-36-3	120	Erie OMZA, Table 33-2 of OAC	0.014	NAWQC 2009			1.20E-04	USEPA Reg 5	1.20E+02	Erie OMZA, Table 33-2 of OAC
Propylene Glycol	57-55-6	640,000	Ohio Administrative Code		1	71,000	Tier II (GLI database)		-	6.40E+05	Ohio Administrative Code
Pyrene	129-00-0	42	Ohio Administrative Code		1	4.6	Tier II (GLI database)	0.3	USEPA Reg 5	4.20E+01	Ohio Administrative Code
Pyridine	110-86-1				1			2,380	USEPA Reg 5	2.38E+03	USEPA Reg 5
RDX	121-82-4	520	Ohio Administrative Code		1	79	Tier II (GLI database)		-	5.20E+02	Ohio Administrative Code
SAS-310		5	Ohio Administrative Code		1	0.61	Tier II (GLI database)		-	5.00E+00	Ohio Administrative Code
Silvex (2,4,5-TP)	93-72-1				-			30	USEPA Reg 5	3.00E+01	USEPA Reg 5
Simazine	122-34-9					9	Tier II (GLI database)			9.00E+00	Tier II (GLI database)
Styrene	100-42-5	290	Ohio Administrative Code			32	Tier II (GLI database)	32	USEPA Reg 5	2.90E+02	Ohio Administrative Code
1,2,4,5-Tetrachlorobenzene	95-94-3					8.3	Tier II (GLI database)	3	USEPA Reg 5	8.30E+00	Tier II (GLI database)
2,3,7,8-Tetrachlorodibenzo-p-dioxin											
(TCDD)	1746-01-6							3.00E-09	USEPA Reg 5	3.00E-09	USEPA Reg 5
1,1,1,2-Tetrachloroethane	630-20-6	770	Ohio Administrative Code			85	Tier II (GLI database)			7.70E+02	Ohio Administrative Code
1,1,2,2-Tetrachloroethane	79-34-5	910	Ohio Administrative Code		-	260	Tier II (GLI database)	380	USEPA Reg 5	9.10E+02	Ohio Administrative Code
Tetrachloroethene	127-18-4	430	Ohio Administrative Code		-	53	Tier II (GLI database)	45	USEPA Reg 5	4.30E+02	Ohio Administrative Code
2,3,4,6-Tetrachlorophenol	58-90-2							1.2	USEPA Reg 5	1.20E+00	USEPA Reg 5
Tetraethyl Dithiopyrophosphate	3689-24-5							13.9	USEPA Reg 5	1.39E+01	USEPA Reg 5
Tetrahydrofuran	109-99-9	74,000	Ohio Administrative Code			11,000	Tier II (GLI database)			7.40E+04	Ohio Administrative Code
Tetryl	479-45-8									No ESV	No Source
Toluene	108-88-3	560	Ohio Administrative Code			62	Tier II (GLI database)	253	USEPA Reg 5	5.60E+02	Ohio Administrative Code
Toxaphene	8001-35-2			0.0002	NAWQC 2009	0.005	Tier II (GLI database)	1.40E-04	USEPA Reg 5	2.00E-04	NAWQC 2009
Tribromomethane (Bromoform)	75-25-2	1,100	Ohio Administrative Code			230	Tier II (GLI database)	230	USEPA Reg 5	1.10E+03	Ohio Administrative Code
2,4,6-Tribromophenol	118-79-6	50	Ohio Administrative Code			5.6	Tier II (GLI database)			5.00E+01	Ohio Administrative Code
Tributyltin (TBT)	688-73-3			0.072	NAWQC 2009		<u>′</u>			7.20E-02	NAWQC 2009
Trichlorobenzene	12002-48-1					5	Tier II (GLI database)			5.00E+00	Tier II (GLI database)
1,2,4-Trichlorobenzene	120-82-1							30	USEPA Reg 5	3.00E+01	USEPA Reg 5

Table C-2. Ecological Screening Values for Chemical Analytes in Surface Water (continued)

						Surfa	ce Water Screening Values				
					Updated Values	for Suter ar	nd Tsao 1996 ^b	USEPA	Region 5 ESLs		
	CAS		Ohio EPA OMZM ^a	NAWQ	C 2009 Update	Tier II Values			(2003) ^c of 1998 EDQLs)		Preferred Surface Water Value ^d
	Registry	Number		Number		Number		Number		Number	
Analyte	Number	(µg/L)	Reference	(μg/L)	Reference	(µg/L)	Reference	(µg/L)	Reference	(µg/L)	Reference
1,1,1-Trichloroethane	71-55-6	690	Ohio Administrative Code			76	Tier II (GLI database)	76	USEPA Reg 5	6.90E+02	Ohio Administrative Code
1,1,2-Trichloroethane	79-00-5	3,300	Ohio Administrative Code			740	Tier II (GLI database)	500	USEPA Reg 5	3.30E+03	Ohio Administrative Code
Trichloroethene	79-01-6	2,000	Ohio Administrative Code			220	Tier II (GLI database)	47	USEPA Reg 5	2.00E+03	Ohio Administrative Code
2,4,5-Trichlorophenol	95-95-4					1.9	Tier II (GLI database)			1.90E+00	Tier II (GLI database)
2,4,6-Trichlorophenol	88-06-2	39	Ohio Administrative Code			4.9	Tier II (GLI database)	4.9	USEPA Reg 5	3.90E+01	Ohio Administrative Code
2,4,5-Trichlorophenoxyacetic Acid	93-76-5							686	USEPA Reg 5	6.86E+02	USEPA Reg 5
O,O,O-Triethyl Phosphorothioate	126-68-1							58.2	USEPA Reg 5	5.82E+01	USEPA Reg 5
Trimethylbenzene	25551-13-7					15	Tier II (GLI database)			1.50E+01	Tier II (GLI database)
1,2,4-Trimethylbenzene	95-63-6	140	Ohio Administrative Code			15	Tier II (GLI database)			1.40E+02	Ohio Administrative Code
1,3,5-Trimethylbenzene	108-67-8	230	Ohio Administrative Code			26	Tier II (GLI database)			2.30E+02	Ohio Administrative Code
1,3,5-Trinitrobenzene	99-35-4	27	Ohio Administrative Code			11	Tier II (GLI database)			2.70E+01	Ohio Administrative Code
2,4,6-Trinitrotoluene	118-96-7	120	Ohio Administrative Code			13	Tier II (GLI database)			1.20E+02	Ohio Administrative Code
Triphenyl Phosphate	115-86-6					4	Tier II (GLI database)			4.00E+00	Tier II (GLI database)
Urea	57-13-6	150,000	Ohio Administrative Code			17,000	Tier II (GLI database)			1.50E+05	Ohio Administrative Code
Vinyl Acetate	108-05-4							248	USEPA Reg 5	2.48E+02	USEPA Reg 5
Vinyl Chloride	75-01-4	8,400	Ohio Administrative Code			930	Tier II (GLI database)	930	USEPA Reg 5	8.40E+03	Ohio Administrative Code
Xylenes (total)	1330-20-7	240	Ohio Administrative Code			27	Tier II (GLI database)	27	USEPA Reg 5	2.40E+02	Ohio Administrative Code

^aOhio Environmental Protection Agency (Ohio EPA), Division of Surface Water. 1999. Ohio Administrative Code (OAC), Chapters 3745-1, 3745-2, May 11 (Ohio River Basin). Where Ohio River Basin is unavailable, Lake Erie is used (as noted).

GLI = Great Lakes Initiative Clearinghouse database, contains Tier II secondary chronic values; http://epa.gov/gliclear/. Values used as supplement to original Suter and Tsao values because of scholarship and methodology shown in Suter and Tsao. Ohio EPA Tier II values used where available; otherwise lowest or most recent value, as appropriate.

NAWQC = National Ambient Water Quality Criteria, originally found in Suter and Tsao 1996 and updated 2009 as National Recommended Water Quality Criteria; values are freshwater chronic. http://epa.gov/waterscience/criteria/wqctable/

NAWQC 2009 value for copper can be found at http://www.epa.gov/waterscience/criteria/copper/2007/criteria-full.pdf NAWQC 2009 value for methyl tert-butyl ether can be found at http://www.epa.gov/waterscience/criteria/mtbe/#findings

-- = no value

CAS = Chemical Abstract Service

Diss = dissolved

ID = Insufficient data available to calculate criterion

ESV = Ecological Screening Value

HMX = Octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazocine

OMZM = Outside Mixing Zone Maximum

PCB = Polychlorinated Biphenyl

RDX = Hexahydro-1,3,5-trinitro-1,3,5-triazine

bSuter, G. W. and C.L. Tsao, Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Aquatic Biota: 1996 Revision, ES/ER/TM-96/R2 Lockheed Martin Energy Systems, Oak Ridge National Laboratory. See notes below for NAWQC and GLI.

^{*}United States Environmental Protection Agency (USEPA) 2003. Ecological Screening Levels (ESLs). Formerly Ecological Data Quality Levels (EDQLs). http://www.epa.gov/reg5rcra/ca/edql.htm

^dThe preferred surface water value is the hierarchy of Chapters 3745-1 and 3745-2 of the Ohio Administrative Code for the Ohio River Basin (1999), Suter and Tsao 1996 (NAWQC followed by Tier II), and EDQLs from USEPA Region 5 (USEPA 2003).

Table C-3. SRC and COPEC Screening with Maximum Ratio for Shallow Surface Soil (0-1 ft bgs Discrete Samples) at Dump Along Paris-Windham Road: Fill Area EU

	CAS	Freq of	Minimum	Maximum	Average	Background	PBT ^b Compound?	SRC?				COPEC?		Ratio of Max to	
Analyte (mg/kg)	Number	Detect	Detect	Detect	Result	Criteria ^a	(yes/no)	(yes/no)	SRC Justification	ESV	ESV Source ^c	(yes/no)	COPEC Justification	ESV	
							Inorgo	anic Chemi	icals						
Aluminum	7429-90-5	5/ 5	6500	11000	8240	17700	No	No	Below background	50	PRGs	No	Below background	220	
Arsenic	7440-38-2	5/ 5	9.2	13	11.2	15.4	No	No	Below background	18	EcoSSL	No	Below background	0.72	
Barium	7440-39-3	5/ 5	47	180	76.6	88.4	No	Yes	Exceeds background	330	EcoSSL	No	Below ESV	0.55	
Beryllium	7440-41-7	5/ 5	0.34	1.2	0.576	0.88	No	Yes	Exceeds background	21	EcoSSL	No	Below ESV	0.06	
Calcium	7440-70-2	5/ 5	1500	39000	9860	15800	No	No	Essential Nutrient	No ESV	No Source	No	Essential Nutrient	No ESV	
Chromium	7440-47-3	5/ 5	8.3	11	9.96	17.4	No	No	Below background	26	EcoSSL	No	Below background	0.42	
Cobalt	7440-48-4	5/ 5	4.3	7.1	5.98	10.4	No	No	Below background	13	EcoSSL	No	Below background	0.55	
Copper	7440-50-8	5/ 5	9.3	19	14	17.7	No	Yes	Exceeds background	28	EcoSSL	No	Below ESV	0.68	
Iron	7439-89-6	5/ 5	14000	22000	18000	23100	No	No	Essential Nutrient	No ESV	No Source	No	Essential Nutrient	No ESV	
Lead	7439-92-1	5/ 5	14	19	16.2	26.1	No	No	Below background	11	EcoSSL	No	Below background	1.73	
Magnesium	7439-95-4	5/ 5	1500	6100	2580	3030	No	No	Essential Nutrient	No ESV	No Source	No	Essential Nutrient	No ESV	
Manganese	7439-96-5	5/ 5	390	880	540	1450	No	No	Below background	220	EcoSSL	No	Below background	4	
Mercury	7439-97-6	5/ 5	0.025	0.048	0.036	0.036	Yes	Yes	Exceeds background	0.00051	PRGs	Yes	Exceeds ESV, PBT Compound	94.12	
Nickel	7440-02-0	5/ 5	10	21	14.2	21.1	No	No	Below background	38	EcoSSL	No	Below background	0.55	
Potassium	7440-09-7	5/ 5	740	1100	892	927	No	No	Essential Nutrient	No ESV	No Source	No	Essential Nutrient	No ESV	
Sodium	7440-23-5	5/ 5	130	380	202	123	No	No	Essential Nutrient	No ESV	No Source	No	Essential Nutrient	No ESV	
Vanadium	7440-62-2	5/ 5	10	15	12.2	31.1	No	No	Below background	7.8	EcoSSL	No	Below background	1.92	
Zinc	7440-66-6	5/ 5	50	100	66.8	61.8	No	Yes	Exceeds background	46	EcoSSL	Yes	Exceeds ESV	2.17	
							Semi-volatile								
Acenaphthylene	208-96-8	1/ 1	0.13	0.13	0.13	0	No	Yes	Detected organic	682	USEPA Reg 5	No	Below ESV	1.91E-04	
Anthracene	120-12-7	1/ 1	0.12	0.12	0.12	0	No	Yes	Detected organic	1480	USEPA Reg 5	No	Below ESV	8.11E-05	
Benz(a)anthracene	56-55-3	1/ 1	1	1	1	0	No	Yes	Detected organic	5.21	USEPA Reg 5	No	Below ESV	0.19	
Benzo(a)pyrene	50-32-8	1/ 1	1.3	1.3	1.3	0	No	Yes	Detected organic	1.52	USEPA Reg 5	No	Below ESV	0.86	
Benzo(b)fluoranthene	205-99-2	1/ 1	1.2	1.2	1.2	0	No	Yes	Detected organic	59.8	USEPA Reg 5	No	Below ESV	0.02	
Benzo(ghi)perylene	191-24-2	1/ 1	0.75	0.75	0.75	0	No	Yes	Detected organic	119	USEPA Reg 5	No	Below ESV	0.01	
Benzo(k)fluoranthene	207-08-9	1/ 1	1.4	1.4	1.4	0	No	Yes	Detected organic	148	USEPA Reg 5	No	Below ESV	0.01	
Chrysene	218-01-9	1/ 1	1.1	1.1	1.1	0	No	Yes	Detected organic	4.73	USEPA Reg 5	No	Below ESV	0.23	
Dibenz(a,h)anthracene	53-70-3	1/ 1	0.24	0.24	0.24	0	No	Yes	Detected organic	18.4	USEPA Reg 5	No	Below ESV	0.01	
Fluoranthene	206-44-0	1/ 1	1.7	1.7	1.7	0	No	Yes	Detected organic	122	USEPA Reg 5	No	Below ESV	0.01	
Indeno(1,2,3-cd)pyrene	193-39-5	1/ 1	0.75	0.75	0.75	0	No	Yes	Detected organic	109	USEPA Reg 5	No	Below ESV	0.01	
Phenanthrene	85-01-8	1/ 1	0.32	0.32	0.32	0	No	Yes	Detected organic	45.7	USEPA Reg 5	No	Below ESV	0.01	
Pyrene	129-00-0	1/ 1	1.4	1.4	1.4	0	No	Yes	Detected organic	78.5	USEPA Reg 5	No	Below ESV	0.02	
								ticides/PCI	Bs						
PCB-1254 ^d	11097-69-1	1/ 1	0.23	0.23	0.23	0	Yes	Yes	Detected organic	0.371	PRGs	Yes	PBT Compound	0.62	

^aBackground criteria for soil 0-1 ft bgs from final facility-wide background values for RVAAP, published in the Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio (USACE 2001). ^bPersistent, Bioaccumulative, and Toxic (PBT) chemicals are defined by Ohio EPA 2008 as: aldrin/dieldrin, chlordane,1,1'-(2,2,2trichloroethylidene)bis[4-chlorobenzene] (DDT) and metabolites (DDD+DDE), hexachlorobenzene, hexachlorobutadiene (hexachloro-1,3-butadiene), hexachlorocyclohexanes (BHCs, alpha-BHC, beta-BHC, delta-BHC), lindane (gammahexachlorocyclohexane), alkyl-lead, mercury and its compounds, mirex, photomirex, octachlorostyrene, PCBs, 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), dioxin, PCDF (furans), 1,2,3,4-tetrachlorodibenzo-p-dioxin (TCDD), 1,2,3,4-tetrachlorodibenzo-p-dioxin (TCDD), 1,2,3,4-tetrachlorodibenzo-p-dio tetrachlorobenzene, 1,2,4,5-tetrachlorobenzene, toxaphene, and other chemicals that are reasonably anticipated to bioaccumulate in animal tissues.

bgs = below ground surface

CAS = Chemical Abstract Service

COPEC = Chemical of Potential Ecological Concern

DOE = United States Department of Energy

EcoSSL = Ecological Soil Screening Level

ESV = Ecological Screening Value

EU = Exposure Unit

Max = Maximum concentration

Ohio EPA = Ohio Environmental Protection Agency

PCB = Polychlorinated biphenyl

PRG = Preliminary Remediation Goal

SRC = Site-related Contaminant

USEPA = United States Environmental Protection Agency

^cScreening Level Source: See table C-1. Hierarchy of values according to Ohio EPA Risk Assessment Guidance is EcoSSLs, followed by DOE 1997a (Preliminary Remediation Goals for Ecological Endpoints. ES/ER/TM-162/R2. August 1997), followed by Region 5 ESLs. ^dESV is for total PCBs

Table C-4. SRC and COPEC Screening with Maximum Ratio for Shallow Surface Soil (0-1 ft bgs Discrete Samples) at Dump Along Paris-Windham Road: Surface Area EU

Analyte (mg/kg)	CAS Number	Freq of Detect	Minimum Detect	Maximum Detect	Average Result	Background Criteria ^a	PBT ^b Compound? (yes/no)	SRC? (yes/no)	SRC Justification	ESV	ESV Source ^c	COPEC? (yes/no)	COPEC Justification	Ratio of Max to ESV
, , , , ,		l .	l					nic Chemi				, ,	l	
Aluminum	7429-90-5	13/ 13	5300	18000	8350	17700	No	Yes	Exceeds background	50	PRGs	Yes	Exceeds ESV	360
Antimony	7440-36-0	2/ 13	0.49	0.6	0.31	0.96	No	No	Below background	0.27	EcoSSL	No	Below background	2.22
Arsenic	7440-38-2	13/ 13	2.6	13	7.53	15.4	No	No	Below background	ε		No	Below background	0.72
Barium	7440-39-3	13/ 13	40	150	74.8	88.4	No	Yes	Exceeds background	330	EcoSSL EcoSSL	No	Below ESV	0.45
Beryllium	7440-41-7	13/ 13	0.33	1.9	0.566	0.88	No	Yes	Exceeds background	21	EcoSSL	No	Below ESV	0.09
Cadmium	7440-43-9	4/ 13	0.1	0.59	0.204	0	No	Yes	Exceeds background	0.36	EcoSSL	Yes	Exceeds ESV	1.64
Calcium	7440-70-2	13/ 13	1700	55000	6650	15800	No	No	Essential Nutrient	No ESV	No Source	No	Essential Nutrient	No ESV
Chromium	7440-47-3	13/ 13	7.9	17	12.4	17.4	No	No	Below background	26	EcoSSL	No	Below background	0.65
Cobalt	7440-48-4	13/ 13	4.7	7.5	5.85	10.4	No	No	Below background	13	EcoSSL	No	Below background	0.58
Copper	7440-50-8	13/ 13	9.4	27	18.5	17.7	No	Yes	Exceeds background	28	EcoSSL	No	Below ESV	0.96
Iron	7439-89-6	13/ 13	12000	18000	14800	23100	No	No	Essential Nutrient	No ESV	No Source	No	Essential Nutrient	No ESV
Lead	7439-92-1	13/ 13	15	29	19.5	26.1	No	Yes	Exceeds background	11	EcoSSL	Yes	Exceeds ESV	2.64
Magnesium	7439-95-4	13/ 13	1300	10000	2700	3030	No	No	Essential Nutrient	No ESV	No Source	No	Essential Nutrient	No ESV
Manganese	7439-96-5	13/ 13	95	1900	386	1450	No	Yes	Exceeds background	220	EcoSSL	Yes	Exceeds ESV	8.64
Mercury	7439-97-6	13/ 13	0.045	0.08	0.0631	0.036	Yes	Yes	Exceeds background	0.00051	PRGs	Yes	Exceeds ESV, PBT Compound	156.86
Nickel	7440-02-0	13/ 13	9.9	37	19.1	21.1	No	Yes	Exceeds background	38	EcoSSL	No	Below ESV	0.97
Potassium	7440-09-7	13/ 13	730	1900	1180	927	No	No	Essential Nutrient	No ESV	No Source	No	Essential Nutrient	No ESV
Silver	7440-22-4	1/ 13	0.39	0.39	0.396	0	No	Yes	Exceeds background	4.2	EcoSSL	No	Below ESV	0.09
Sodium	7440-23-5	11/ 13	120	480	185	123	No	No	Essential Nutrient	No ESV	No Source	No	Essential Nutrient	No ESV
Vanadium	7440-62-2	13/ 13	9.3	18	13	31.1	No	No	Below background	7.8	EcoSSL	No	Below background	2.31
Zinc	7440-66-6	13/ 13	51	120	81.7	61.8	No	Yes	Exceeds background	46	EcoSSL	Yes	Exceeds ESV	2.61
								Explosives						
Nitrocellulose	9004-70-0	1/ 1	2	2	2	0	No	Yes	Detected organic	No ESV	No Source	Yes	Detected organic	No ESV
							Semi-volatile	Organic C	ompounds					
Benz(a)anthracene	56-55-3	1/ 1	0.25	0.25	0.25	0	No	Yes	Detected organic	5.21	USEPA Reg 5	No	Below ESV	0.05
Benzo(a)pyrene	50-32-8	1/ 1	0.33	0.33	0.33	0	No	Yes	Detected organic	1.52	USEPA Reg 5	No	Below ESV	0.22
Benzo(b)fluoranthene	205-99-2	1/ 1	0.39	0.39	0.39	0	No	Yes	Detected organic	59.8	USEPA Reg 5	No	Below ESV	0.01
Benzo(k)fluoranthene	207-08-9	1/ 1	0.33	0.33	0.33	0	No	Yes	Detected organic	148	USEPA Reg 5	No	Below ESV	0.002
Chrysene	218-01-9	1/ 1	0.33	0.33	0.33	0	No	Yes	Detected organic	4.73	USEPA Reg 5	No	Below ESV	0.07
Fluoranthene	206-44-0	1/ 1	0.44	0.44	0.44	0	No	Yes	Detected organic	122	USEPA Reg 5	No	Below ESV	0.004
Pyrene	129-00-0	1/ 1	0.44	0.44	0.44	0	No	Yes	Detected organic	78.5	USEPA Reg 5	No	Below ESV	0.01
								ticides/PCB	Ss					
PCB-1254 ^d	11097-69-1	1/ 1	0.086	0.086	0.086	0	Yes	Yes	Detected organic	0.371	PRGs	Yes	PBT Compound	0.23
								rganic Com	pounds					
Acetone	67-64-1	1/ 1	0.041	0.041	0.041	0	No	Yes	Detected organic	2.5	USEPA Reg 5	No	Below ESV	0.02

^aBackground criteria for soil 0-1 ft bgs from final facility-wide background values for RVAAP, published in the Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio (USACE 2001). ^bPBT Persistent, Bioaccumulative, and Toxic (PBT) chemicals are defined by Ohio EPA 2008 as: aldrin/dieldrin, chlordane, 1,1'-(2,2,2trichloroethylidene) bis [4-chlorobenzene] (DDT) and metabolites (DDD+DDE), hexachlorobenzene, hexachlorobutadiene (hexachloro-1,3-butadiene), hexachlorocyclohexanes (BHCs, alpha-BHC, beta-BHC, delta-BHC), lindane (gammahexachlorocyclohexane), alkyl-lead, mercury and its compounds, mirex, photomirex, octachlorostyrene, polychlorinated biphenyls (PCBs), 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), dioxin, PCDF (furans), 1,2,3,4-tetrachlorobenzene, 1,2,4,5-tetrachlorobenzene, toxaphene, and other chemicals that are reasonably anticipated to bioaccumulate in animal tissues.

^cScreening Level Source: See table C-1. Hierarchy of values according to Ohio EPA Risk Assessment Guidance is EcoSSLs, followed by DOE 1997a (Preliminary Remediation Goals for Ecological Endpoints. ES/ER/TM-162/R2. August 1997), followed by Region 5 ESLs.

^dESV is for total PCBs

bgs = below ground surface

CAS = Chemical Abstract Service

COPEC = Chemical of Potential Ecological Concern

DOE = United States Department of Energy

EcoSSL = Ecological Soil Screening Level

ESV = Ecological Screening Value

EU = Exposure Unit

Max = Maximum concentration

Ohio EPA = Ohio Environmental Protection Agency

PRG = Preliminary Remediation Goal

SRC = Site-related Contaminant

USEPA = United States Environmental Protection Agency

Table C-5. SRC and COPEC Screening for Surface Water at Dump Along Paris-Windham Road

Analyte (mg/L)	CAS Number	Freq of Detect	Minimum Detect	Maximum Detect	Average Result	Background Criteria ^a	PBT ^b Compound? (yes/no)	SRC? (yes/no)	SRC Justification	ESV	ESV Source ^c	COPEC? (yes/no)	COPEC Justification	Ratio of Max to ESV
				1	Γ	T	Inorganic	Chemicals		ı			T	
Aluminum	7429-90-5	7/ 7	0.042	0.28	0.104	3.37	No	No	Below background	0.087	NAWQC 2009	No	Below background	3.22
Arsenic	7440-38-2	7/ 7	0.0028	0.0082	0.00549	0.0032	No	Yes	Exceeds background	0.34	Ohio Administrative Code	No	Below ESV	0.02
Barium	7440-39-3	7/ 7	0.035	0.12	0.066	0.0475	No	Yes	Exceeds background	2	Ohio Administrative Code	No	Below ESV	0.06
Calcium	7440-70-2	7/ 7	23	60	41.4	41.4	No	No	Essential Nutrient	No ESV	No Source	No	Essential Nutrient	No ESV
Cobalt	7440-48-4	4/ 7	0.001	0.0015	0.00177	0	No	Yes	Exceeds background	0.22	Ohio Administrative Code	No	Below ESV	0.01
Copper	7440-50-8	7/ 7	0.0022	0.0039	0.0025	0.0079	No	No	Below background	0.014	Ohio Administrative Code ^d	No	Below background	0.28
Iron	7439-89-6	7/ 7	3.6	9.4	5.04	2.56	No	No	Essential Nutrient	1	NAWQC 2009	No	Essential Nutrient	9.40
Lead	7439-92-1	2/ 7	0.0019	0.0027	0.00137	0	No	Yes	Exceeds background	0.12	Ohio Administrative Code ^d	No	Below ESV	0.02
Magnesium	7439-95-4	7/ 7	6	12	9.4	10.8	No	No	Essential Nutrient	No ESV	No Source	No	Essential Nutrient	No ESV
Manganese	7439-96-5	7/ 7	0.26	0.56	0.379	0.391	No	Yes	Exceeds background	0.12	Tier II (Suter & Tsao 1996)	Yes	Exceeds ESV	4.67
Mercury	7439-97-6	6/ 7	0.000072	0.0001	0.0000896	0	Yes	Yes	Exceeds background	0.0017	Ohio Administrative Code	Yes	PBT Compound	0.06
Nickel	7440-02-0	4/ 7	0.002	0.0075	0.00487	0	No	Yes	Exceeds background	0.47	Ohio Administrative Code ^d	No	Below ESV	0.02
Potassium	7440-09-7	7/ 7	1.7	5.4	4.17	3.17	No	No	Essential Nutrient	No ESV	No Source	No	Essential Nutrient	No ESV
Sodium	7440-23-5	7/ 7	4.2	9.9	7.76	21.3	No	No	Essential Nutrient	No ESV	No Source	No	Essential Nutrient	No ESV
Zinc	7440-66-6	4/ 7	0.013	0.024	0.0149	0.042	No	No	Below background	0.12	Ohio Administrative Code	No	Below background	0.20
							Miscell	laneous						
Asbestos (MFL)	1332-21-4	1/ 6	0.0001	0.0001	6.73	0	No	Yes	Exceeds background	No ESV	No Source	Yes	Exceeds background	No ESV
					T	1	Explo	osives		T				
Nitrocellulose	9004-70-0	1/ 1	0.094	0.094	0.094	0	No	Yes	Exceeds background	No ESV	No Source	Yes	Exceeds background	No ESV

^aBackground criteria from final facility-wide background values for RVAAP, published in the Final Phase II Remedial Investigation Report for Winklepeck Burning Grounds at Ravenna Army Ammunition Plant, Ravenna, Ohio (USACE 2001).

CAS = Chemical Abstract Service

COPEC = Chemical of Potential Ecological Concern

ESL = Ecological Screening Level

ESV = Ecological Screening Value

Max = Maximum concentration

MFL = Million Fibers per Liter

NAWQC = National Ambient Water Quality Criteria

Ohio EPA = Ohio Environmental Protection Agency

PBT = Persistent, Bioaccumulative, and Toxic

RVAAP = Ravenna Army Ammunition Plant

SRC = Site-related Contaminant

Bold = Chemical is a COPEC

bPBT chemicals are defined by Ohio EPA 2008 as: aldrin/dieldrin, chlordane, 1,1'-(2,2,2trichloroethylidene)bis[4-chlorobenzene] (DDT) and metabolites (DDD+DDE), hexachloro-butadiene (hexachloro-1,3-butadiene), hexachlorocyclohexanes (BHCs, alpha-BHC, beta-BHC, delta-BHC), lindane (gammahexachlorocyclohexane), alkyl-lead, mercury and its compounds, mirex, photomirex, octachlorostyrene, polychlorinated biphenyls (PCBs), 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), dioxin, PCDF (furans), 1,2,3,4-tetrachlorobenzene, 1,2,4,5-tetrachlorobenzene, 1,2,4,5-tetrachlorobenzene toxaphene, and other chemicals that are reasonably anticipated to bioaccumulate in animal tissues.

Screening Level Source: See table C-2. Hierarchy of values according to Ohio EPA Risk Assessment Guidance and letter from Ohio EPA oMZM (Outside Mixing Zone Max), followed by NAWQC or Tier II values, followed by Region 5 ESLs. ^dValue is hardness dependent

	Army	Ohio EPA	Dump Along Paris- Windham Road	
Resource	(2005)	(2008)	Absent	Present
National Park	X	X	X	
Designated Federal Wilderness Area	X	X	X	
National Lakeshore Recreational Area	X	X	X	
Habitat known to be used by federal designated or proposed threatened or endangered species	X	X	X	
National or State Wildlife Refuge	X	X	X	
Federal land designated for protection of natural ecosystems	X	X	X	
Habitat known to be used by state designated threatened or endangered species	X	X	X	
Federally-designated Scenic or Wild River	X	X	X	
State land designated for wildlife or game management	X	X	X	
State-designated Scenic or Wild River	X	X	X	
Wetlands and waters of the State ^a	X	X		X
National preserve	X	X^{b}	X	
State-designated Natural Areas	X	X^{b}	X	
Spawning areas critical for the maintenance of fish/shellfish species within river, lake, or coastal tidal waters	X	X^{c}	X	
Migratory pathways and feeding areas critical for maintenance of anadromous fish species ^d	X	X ^c	X	
Terrestrial areas used for breeding by large or dense aggregations of animals	X	X^{c}	X	
Particular areas, relatively small in size, important to maintenance of unique biotic communities	X	X^{c}	X	
Locally important ecological place ^e	X		X	
Critical habitat for federal designated threatened or endangered species	X		X	
Marine Sanctuary	X		X	
Areas identified under the Coastal Zone Management Act	X		X	
Sensitive Areas identified under the National Estuary Program or Near Coastal Waters Program	X		X	
Critical areas identified under the Clean Lakes Program	X		X	
National Monument	X		X	
National Seashore Recreational Area	X		X	
Unit of Coastal Barrier Resources System	X		X	
Coastal Barrier (undeveloped)	X		X	
Coastal Barrier (partially developed)	X		X	
Administratively Proposed Federal Wilderness Area	X		X	
National river reach designated as Recreational	X		X	

Table C-6. Checklist of Important Ecological Places and Resources at Dump Along Paris-Windham Road (continued)

	Army	Ohio EPA	Dump Along Paris- Windham Road	
Resource	(2005)	(2008)	Absent	Present
Habitat known to be used by species under review as to its Federal threatened or endangered status	X		X	
State-designated areas for protection or maintenance of aquatic life	X		X	
Fragile landscapes, land sensitive to degradation if vegetative habitat or cover diminishes	X		X	
State, local, or private land designated for protection of natural ecosystems		X	X	
Federal land designated for wildlife or game management		X	X	
Surface water, as that term is used in Chapter 3745-1 of the OAC		X	X	
Federally-listed or state-listed threatened or endangered species		X	X	
State of Ohio special interest or declining species and its associated habitat		X	X	
State Park		X	X	

- U.S. Army Biological Technical Assistance Group, Technical Document for Ecological Risk Assessment: Process for
 Developing Management Goals. August 2005.
 Ohio EPA. Guidance for Conducting Ecological Risk Assessments (Ohio EPA). Division of Emergency and Remedial
 - Ohio EPA. Guidance for Conducting Ecological Risk Assessments (Ohio EPA). Division of Emergency and Remedial Response. April 2008.
- ^aFor Ohio EPA 2008, as qualified by "regulated under federal law and state of Ohio's water quality laws."
- 8 bOhio EPA does not restrict preserves and natural areas to National or State.
- 9 °Ohio EPA lists "wildlife populations and their associated important nesting areas and food resources, taking into consideration land use and the quality and extent of habitat on and in the vicinity of the site."
- 11 dWithin river reaches or areas in lakes or coastal tidal waters in which fish spend extended periods of time.
- 12 °Identified by the Integrated Natural Resource Management Plan (INRMP), Base Realignment and Closure (BRAC)
- Cleanup Plan or Redevelopment Plan, or other official land management plans.
- The Ohio Army National Guard (OHARNG 2008) has five special interest areas (important resources) at RVAAP: mixed mature woods, Hemlock Ravine-Wadsworth Glen, mixed swamp forest, mixed valuable communities, and oak/maple
- swamp forest. Also, the OHARNG recognizes the importance of federal and state-listed threatened and endangered plant and animal species.
 - x = designated as important and when bolded there are possible qualifiers
- 19 OAC = Ohio Administrative Code
- 20 Ohio EPA = Ohio Environmental Protection Agency

21

18

6

Goals and Objectives of	Comments on Goals
Ohio Army National Guard Goal 1. Manage natural resources in a manner that is compatible with and supports the military mission while complying with applicable Federal and State laws and Army regulations and policies. Objective 1.1: Initiate programs and projects that enhance the training land and training opportunities and/or do not unnecessarily limit training land availability.	U.S. Army committed to natural resources management in a manner that is compatible with and supports the military mission and complies with Federal and State laws and Army regulations and policies.
Objective 1.2: Continue to educate Camp Ravenna users regarding the natural resources at the Camp Ravenna and their part in ensuring sustainable use of the site in perpetuity.	
Goal 2. Maintain and foster positive working relationships with the U.S. Fish and Wildlife Service, the ODNR DOW, and other federal, state and local natural resources management agencies and organizations for the benefit of the military mission, the natural resources being managed, and the citizens of Ohio and the nation.	The U.S. Army works and coordinates with other federal and state agencies as necessary if mission or projects have the potential to impact natural resources.
Objective 2.1: Effectively communicate mission needs to cooperating agencies and solicit input/review on projects with the potential to impact natural resources, especially in areas of regulatory primacy.	
Objective 2.2: Provide copies of biological surveys to interested cooperating agencies.	
Objective 2.3: Facilitate cooperative management programs and projects that are compatible with the military mission and within the capabilities of the Camp Ravenna staff.	
Goal 3. Monitor the condition of the natural resources and the implied impacts from training and the natural resources management program on the natural resources at the Camp Ravenna.	The U.S. Army conducts natural resource management activities at the facility to monitor potential impacts from training or other disturbance activities.
Objective 3.1: Maintain current species inventories and other PLSs through periodic reoccurring surveys and inventories.	
Goal 4. Protect and maintain populations of rare plant and animal species on the Camp Ravenna in compliance with Federal and State laws and regulations.	The U.S. Army protects and maintains populations of rare plant and animal species by implementing a natural resource management plan at the facility and by avoiding and/or not disturbing areas with rare species.
Objective 4.1: Avoid negative impacts to federally listed species and avoid/minimize impacts to State listed and otherwise rare species.	

Goals and Objectives of	Comments on Goals
Ohio Army National Guard	Relative to HTRW Work at RVAAP
Goal 5. Sustain usable training lands and native	The U.S. Army sustains usable training lands and
natural resources by managing non-native and	native natural resources by implementing a natural
invasive species, vegetation and plant communities, and nuisance wildlife species.	resource management plan which incorporates invasive species and nuisance species management
Objective 5.1: Manage populations of invasive plant species where they hinder training and/or habitat management objectives.	and by utilizing native species mixes for re-vegetation after ground disturbance activities.
Objective 5.2: Manage non-native and invasive insect species that pose a threat to forest resources.	
Objective 5.3: Manage terrestrial vegetation to support training, encourage native plant communities, and prevent damage to training site facilities and infrastructure.	
Objective 5.4: Manage the beaver population to prevent damage to training site facilities and infrastructure and to maintain the quality warm water habitats of Hinkley Creek, Sand Creek, and South Fork Eagle Creek.	
Objective 5.5: Manage other nuisance animals that negatively impact the ecosystem.	
Goal 6. Manage wildlife resources in a manner compatible with the military mission and within the limits of the natural habitat.	The U.S. Army minimizes habitat disturbance during HTRW activities and utilizes sustainability practices when disturbance is required in order to properly manage and maintain wildlife populations and
Objective 6.1: Cooperatively manage wildlife resources with the Ohio DOW.	resources.
Objective 6.2: Provide opportunity for wildlife recreation to the public that is compatible with the military mission.	
Objective 6.3: Maintain wildlife population without augmenting the habitat with artificial food plots.	

Goals and Objectives of Ohio Army National Guard	Comments on Goals Relative to HTRW Work at RVAAP
Goal 7. Manage forest resources to the benefit of the military mission, to perpetuate the ecosystem functions, to support regional ecosystem needs, and for the production of forest products. Objective 7.1: Maintain current forest resource data.	The U.S. Army sustains and manages forest resources by implementing a natural resource management plan. During HTRW activities, efforts are made by the Army to minimize impacts to forest communities.
Objective 7.2: Implement forest management strategies identified in the Camp Ravenna INRMP.	
Goal 8. Manage wetlands and other surface waters in accordance with applicable Federal, State, and local regulations and to protect water quality and ecological functions while facilitating the military mission. Objective 8.1: Avoid wetland fills.	Wetlands and other surface waters are to be protected during disturbance activities in accordance with federal, state, and local regulations. Avoidance measures will be implemented as practical. Some AOCs have wetlands.
Objective 8.2: Minimize and mitigate unavoidable wetland fills.	
Objective 8.3: Maintain healthy aquatic ecosystems in ponds.	
Objective 8.4: Restore, enhance and create wetlands when possible and compatible with the military mission.	
Goal 9. Manage soil to maintain productivity and prevent and repair erosion in accordance with State and Federal laws and regulations so that the Camp Ravenna can support doctrinally required military training in perpetuity.	Management of soil relevant to remedial activities under CERCLA. Appropriate storm water and erosion controls are to be utilized during activities that require ground disturbance.
Objective 9.1: Conduct training and other activities in locations with soil most suitable for supporting the activity.	
Objective 9.2: Rehabilitate, repair, and maintain areas damaged by training and other activities.	

Goals and Objectives of Ohio Army National Guard	Comments on Goals Relative to HTRW Work at RVAAP
Goal 10. Manage cultural resources on the Camp Ravenna in accordance with State and Federal laws and regulations while implementing the natural resources management program. Objective 10.1: Comply with Federal, State, and local laws and regulations pertaining to cultural resources found on the training site.	The U.S. Army utilizes a cultural resource management plan to manage and protect cultural resources at the facility. Coordination with state and federal agencies regarding cultural resources is conducted as necessary. Restoration contractors are also advised to utilize the Camp Ravenna Policy for Inadvertent Discoveries for reporting purposes should they come upon a cultural item.
Goal 11. Develop, maintain, and manage data regarding natural resources at the Camp Ravenna through the use of GIS for efficient data storage, retrieval, analysis, and presentation. Objective 11.1: Develop accurate and usable natural resources GIS data.	Natural resource data is collected and managed by the OHARNG. This data may be utilized during restoration activities in order to provide an accurate portrait of natural resources at an AOC.

- 23456789 OHARNG. Integrated Natural Resources Management Plan and Environmental Assessment for the Ravenna Training and Logistics Site, Portage and Trumbull Counties, Ohio. March 2008.
- AOC = Area of Concern
- CERCLA = Comprehensive Environmental Response, Compensation, and Liability Act
- DOW = Department of Wildlife
- GIS = Geographic Information System
- HTRW = Hazardous, Toxic and Radioactive Waste
- INRMP = Integrated Natural Resources Management Plan
- 10 ODNR = Ohio Department of Natural Resources
- 11 OHARNG = Ohio Army National Guard
- 12 Ohio EPA = Ohio Environmental Protection Agency
- 13 PLS = Planning Level Survey (Wetland)
- RVAAP = Ravenna Army Ammunition Plant 14
- 15 T & E = Threatened and Endangered Species
- 16 USACE = United States Army Corps of Engineers

1	REFERENCES
2	U.S. Army Biological Technical Assistance Group (BTAG) 2005. Technical Document for
4	Ecological Risk Assessment: Process for Developing Management Goals. August 2005.
5	Note that the state of the stat
6	National Recommended Water Quality Criteria (NAWQC) 2009.
7	http://water.epa.gov/scitech/swguidance/waterquality/standards/current/index.cfm. 2009
8	Obia Assess National Count (OHADNC) 2000 Language I.N. and I.D. Market I.D. Ma
9	Ohio Army National Guard (OHARNG) 2008. Integrated Natural Resources Management Plan and
10 11	Environmental Assessment for the Ravenna Training and Logistics Site, Portage and Trumbull Counties, Ohio. March 2008.
12	Trumbun Counties, Onto. March 2006.
13	Ohio Environmental Protection Agency (Ohio EPA) 2001. Ohio Rapid Assessment Method for
14	Wetlands v. 5.0, User's Manual and Scoring Forms. Division of Surface Water, 401/Wetland
15	Ecology Unit. February 2001.
16	
17	Ohio EPA 2003. Guidance for Conducting Ecological Risk Assessments (Ohio EPA). Division of
18	Emergency and Remedial Response. February 2003.
19	
20	Ohio EPA 2008. Guidance for Conducting Ecological Risk Assessments (Ohio EPA). Division of
21	Emergency and Remedial Response. April 2008.
22	
23	Suter, G.W. II, and C.L. Tsao 1996. Toxicological Benchmarks for Screening Potential Constituents
24	of Concern for Effects on Aquatic Biota: 1996 Revision, ES/ER/TM 96/R2, Lockheed Martin
25	Energy Systems, Oak Ridge National Laboratory, Oak Ridge, TN. June 1996.
26 27	United States Army Corns of Engineers (USACE) 2001. Phase II Pamedial Investigation Penert for
28	United States Army Corps of Engineers (USACE) 2001. Phase II Remedial Investigation Report for the Winklepeck Burning Grounds at the Ravenna Army Ammunition Plant, Ravenna, Ohio.
20 29	April 2001.
30	11pm 2001.
31	United States Environmental Protection Agency (USEPA) 1998. Technical Approach for Developing
32	EDQLs for Resource Conservation and Recovery Act (RCRA) Appendix IX Constituents and
33	Other Significant Contaminants of Ecological Concern. USEPA, Region 5. April 1998.
34	
35	USEPA 2003. Ecological Screening Levels. USEPA, Region 5. August 2003.
36	
37	USEPA 2010. Ecological Soil Screening Levels (Eco-SSLs). Office of Solid Waste and Emergency
38	Response, Washington, D.C. 2010. http://www.epa.gov/ecotox/ecossl/ 2010.

THIS PAGE INTENTIONALLY LEFT BLANK.

1

APPENDIX D

Detailed Cost Estimate

Focused Feasibility Study for Soil and Dry Sediment Dump Along Paris-Windham Road - Ravenna Army Ammunition Plant (RVAAP), Ravenna, Ohio Summary of Alternatives

	Dump Along Paris-Windham Road Alternatives		Non Discounted Cost Soil and Dry Sediment			
			Capital Cost	O&M Cost	Total	
1	No Action	30 years	\$0	\$0	\$0	
2	Land Use Controls	30 years	\$16,024	\$142,015	\$158,039	

Dump Along Paris-Windham Road Alternatives		Duration	Discounted Cost (4.125%)			
			Soil and Dry Sediment			
			Capital Cost	O&M Cost	Total	
1	No Action	30 years	\$0	\$0	\$0	
2	Land Use Controls	30 years	\$16,024	\$77,360	\$93,384	

Notes:

- 1. The base year of comparison and cost data will be CY2010. The discounted rates used to calculate present values will be based on Economic Guidance Memorandum, 11-01, Federal Interest Rates for Corps of Engineers Projects for Fiscal Year 2011.
- 2. Costs were estimated for comparison purposes only and are believed to be accurate within a range of -30% to +50%. Use of these costs for other purposes, including but not limited to, budgetary or construction cost estimating is not appropriate.

1

Dump Along Paris-Windham Road - Ravenna Army Ammunition Plant (RVAAP), Ravenna, Ohio Alternative 2 - Land Use Controls Key Parameters and Assumptions

Key Parameters and Assumptions:

Item	Unit	Value	Notes
Capital Cost			
Land Use Controls			
Base Master Planning Documents	hrs	80	Accuracy CO has to accious and accion DMD decuments. Included
Legal/Technical Labor	\$/hr	120	Assume 80 hrs to review and revise BMP documents. Included deed and groundwater restrictions.
Site Work	Ψ/	0	
Site Area	sf	12,000	
Civil Survey	day	1.0	
Civil Survey	\$/day	950	Survey AOC for land use controls. RSMeans 017123131100.
As Built Drawings	hours	8	Develop record drawings.
As Built Drawings As Built Drawings	\$/hr	60	Develop record drawings.
Install Signs on Posts	ea	6	Assume warning signs located around AOC perimeter at 300 ft
Install Signs on Posts	\$/ea	209.00	centers. RSMeans 028907000100 & 1500. Add 25% for custom
motali digno di 1 doto	φ/σα	200.00	letters. Furnish, place, and install.
Plans and Reports			
Corrective Action Completion Report	hrs	80	Includes documentation of corrective action and report.
Technical Labor	\$/hr	80	
	**		
O&M Cost (Years 0 to 30)			
Site Inspection and Maintenance	years	30	
Site Inspection	events	60	
Site Inspections	hrs	4	Inspect site semi-annually for disturbance/erosion, warning signs,
Field Labor	\$/hr	60	and complete checklist for annual report.
Site Maintenance	events	30	Assume signs are replaced every 10 years. Assume AOC area
Site Maintenance	\$/yr	290	is overseeded and fertilized every 5 years. Costs have been
Assessed Dansert			annualized.
Annual Report		00	
Annual O&M Report	event	30	
Annual O&M Report	\$/year	640	Assume 8 hours @ \$80/hr for letter report.
CERCLA Reviews			
CERCLA 5-Year Reviews	events	6	Assume 5 year reviews for 30 years.
CERCLA 5-Year Reviews	\$/event	7,400	Assume 80 hours/review @ \$80/hr. Add \$1,000 misc expenses.
		·	

Dump Along Paris-Windham Road - Ravenna Army Ammunition Plant (RVAAP), Ravenna, Ohio Alternative 2 - Land Use Controls Cost Estimate

CAPITAL COST	\$16,024

Activity (unit)	Quantity	Unit Cost	Total
Land Use Controls			
Base Master Planning Documents (hr)	80	\$120.00	\$9,600
Site Work			
Civil Survey (day)	1.0	\$950.00	\$950
As Built Drawings (hrs)	8	\$60.00	\$480
Install Signs on Posts (ea)	6	\$209.00	\$1,254
Plans and Reports			
Corrective Action Completion Report (ea)	80	\$80.00	\$6,400
Subtotal			\$9,084
Design		20%	\$1,817
Office Overhead		5%	\$454
Field Overhead		15%	\$1,363
Subtotal			\$12,718
Profit		6%	\$763
Contingency		20%	\$2,544
Total			\$16,024

OPERATION AND MAINTENANCE

\$142,015

Activity (unit)	Quantity	Unit Cost	Total Cost	Present Value (4.125%)
Site Inspection and Maintenance				
Site Inspection (ea)	60	\$240	\$14,400	\$8,176
Site Maintenance (ea)	30	\$290	\$8,700	\$4,939
Annual Report				
Annual O&M Report (ea)	30	\$640	\$19,200	\$10,901
CERCLA Reviews				
CERCLA 5-Year Reviews (ea)	6	\$7,400	\$44,400	\$23,213
Subtotal O&M			\$86,700	\$47,228
Design		10%	\$8,670	\$4,723
Office Overhead		5%	\$4,335	\$2,361
Field Overhead		15%	\$13,005	\$7,084
Subtotal			\$112,710	\$61,397
Profit		6%	\$6,763	\$3,684
Contingency		20%	\$22,542	\$12,279
Total			\$142,015	\$77,360

TOTAL ALTERNATIVE CAPITAL AND O&M COST (Non Discounted Cost)

\$158,039